



Bioaccessibility of Trace Elements in Different Oyster Mushroom Varieties Grown in Kenya

Rachel Nambafu^a, Judith Waudu^a, Margaret Nganga^a,
Peter Lusi^a and Hudson Nyambaka^{a*}

^a Chemistry Department, Kenyatta University, P.O. Box-43844 - 00100, Nairobi, Kenya.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AIR/2023/v24i4945

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96473>

Original Research Article

Received: 17/12/2022

Accepted: 22/02/2023

Published: 23/03/2023

ABSTRACT

Trace elements, especially chromium (Cr), vanadium (V) and selenium (Se) have potential beneficial effects on glucose metabolism in people with type 2 diabetes. Food products incorporating mushrooms are not only a good source of such nutrients but are thought to have readily bioavailable nutrients. Nutritional efficacy of food products may be ensured by accessing bioaccessibility of nutrients, which provides valuable information on matrix and appropriate dosage. The study determined bioaccessible Cr, V and Se in four varieties of oyster mushrooms *Pleurotus ostreatus* (PO), *Pleurotus sajor-caju* (PS), *Pleurotus pulmonarius* (PP) and *Pleurotus citrinopileatus* (PC) grown in Kenya. Bioaccessibility was estimated using in vitro simulated gastrointestinal procedure, while nutrient levels were determined using atomic absorption procedure. Bioaccessible levels of chromium ranged from 26.56% in PS to 78.50% in PC; selenium from 92.52% in PC to ND in PS and PP; and vanadium from 92.46% in PC to 69.95% in PP. Vanadium was the most bio accessible than the other elements in the four oyster mushrooms, while chromium was the most bio accessible in the PC variety.

*Corresponding author: E-mail: nyambaka.hudson@ku.ac.ke;

Keywords: Bioaccessibility; oyster mushrooms varieties; vanadium; chromium; selenium.

1. INTRODUCTION

“Bioaccessibility can be defined as quantity or fraction which is released from the food matrix in the gastrointestinal tract and becomes available for absorption” [1-3]. “Bioavailability on the other hand is the ingested fraction available at the site of action for utilization in normal physiological functions; usually determined through *In vivo* assays” [4,5]. “It is the result of three main steps: digestibility and solubility of the element in the gastrointestinal tract; absorption of the element by the intestinal cells and transport into the circulation; and incorporation from the circulation to the functional entity or target” [6,7]. “Bioaccessibility is usually evaluated by *in vitro* digestion procedures generally simulating gastric and small intestinal digestion, sometimes followed by Caco-2-cell uptake” [8]. “*In vitro* methods are developed to simulate the physiological conditions (temperature, agitation, pH, enzyme, and chemical composition) and the sequence of events that occur during digestion in the human gastrointestinal tract” [9]. The bioaccessible level of trace elements after consumption is not necessarily related to the levels in foods since food processing and cooking affect intestinal transit time of nutrients as well as their enteric formation of mixed micelles [3,10]. In this study a static method was used which involved *in vitro* enzymolysis procedure simulating human gastro intestinal involving two steps (gastric and intestinal).

Mushrooms offer tremendous applications as they can be used as food and medicines besides their key ecological roles. Bano [11] suggested that “food value of mushrooms lies between meat and vegetables”. “The fruiting bodies of mushrooms are characterized by a high level of well assimilated mineral elements. Major mineral constituents in mushrooms are K, P, Na, Ca, Mg and elements like Cu, Zn, Fe, Mo, Cd form minor constituents” [12-14]. “The present use of mushrooms is totally different from the traditional use because a lot of research has been done on the chemical composition of mushrooms, which has revealed that mushrooms can be used as a diet to combat diseases. The early history regarding the use of mushrooms in different countries has been reviewed by a number of workers” [13,15-17,13,14]. “The mineral proportions vary according to the species, age, and the diameter of the fruiting body. It also

depends upon the type of the substratum” [18]. “The mineral content of wild edible mushrooms has been found higher than cultivated ones” [19-21]. “The minerals have been found to boost the immune system, have anti-cancerous properties, and act as anti-hypercholesterolaemic and hepatoprotective properties. Some mushrooms such as *Pleurotus* species are excellent food for the people suffering from hypertension and cardiovascular diseases due to high potassium and sodium content” [22]. “Chromium is generally recognized to play an important role in glucose and lipid metabolism. One effect of chromium is that it is very helpful in preventing and reversing type 2 diabetes because plasma glucose is more effectively regulated in the presence of chromium” [23].

“Vanadium compounds have been demonstrated to mimic the action of insulin in isolated cell systems, animal models and diabetic patients. This has brought the use of V compounds as potential sources of diabetes therapy into focus” [24,25]. “Selenium have been shown to reduce the risk of developing dysglycemia- a broad term that refers to any abnormalities in blood glucose levels leading to disease” [26]. “It is also thought to work by exerting insulin-like actions *in vitro*” [27].

In Kenya hypertension and diabetes are becoming the major causes of morbidity and mortality with their prevalence surpassing that of HIV/AIDs [28]. Various mushrooms species are used in the treatment and prevention of diabetes, due to their rich nutritional and medicinal properties such as the elements (K, Ca, Na, P, Fe, Cr, Se, and V) [29]. The management of diabetes through use of nutraceuticals requires that the substrate not only contain high levels nutrients but also bio available nutrients. Since the levels of the nutrients depend on growing medium, variety, maturity of tissue and climatic condition it is important to quantify the levels in various species of mushrooms. Mushrooms have a chitin structure that is likely to have effect on the micronutrients bio available. Bioavailability of nutrients depends on various factors such as food tissue and on host related factors and is studied through intervention (*in vitro* procedures), or algorithms and simulated gastrointestinal digestion (*in vivo* methods). Bioavailability is improved by formulating a food product containing high levels of nutrients such Cr, V and Se that are important in management of

diabetes. This study was conducted to determine the bioaccessibility of trace elements (chromium, vanadium, and selenium) in the fresh mushrooms commonly consumed in Kenya.

2. MATERIALS AND METHODS

2.1 Experimental

2.1.1 Samples and sample preparation

Four oyster mushroom that is *Pleurotus ostreatus* (PO), *Pleurotus sajor-caju* (PS), *Pleurotus pulmonarius* (PP) and *Pleurotus citrinopileatus* (PC) varieties were collected from nearby farms in Jomo Kenyatta University and identified by a taxonomist. The freshly harvested mushrooms were cleaned, cut into pieces, and dried in the shade for four days and then powdered before keeping in a labeled special glass container awaiting analysis.

2.1.2 Reagents and apparatus

All reagents were of analytical reagent grade. Double deionized water was used in dilution of reagent solutions. HNO₃ and H₂O₂ were of supra pure quality (Merck). Gastric juice, Pepsin (1% w/v), Pcreatin (3% w/v), Amylase (1% w/v), Bile salt, selenium, chromium and vanadium as standards were purchased from Sigma Andrich. All plastics and glassware were cleaned, soaked in dilute HNO₃ (10%) and rinsed with distilled water prior to use.

2.1.3 Instrumentation

Atomic Absorption Spectrophotometer (AAS), Thermo Jarell Ash (model AAS S11) was used for analysis of trace elements. The most appropriate wavelength, flow rate, slit width and other AAS instrument parameters for metals, minerals were selected as given in the instrument user's manual and background correction was used during determination of metals/minerals.

2.2 Procedures of Analysis

2.2.1 Method validation

Calibration of AAS was done using the working standard prepared from commercially available metal/mineral standard solutions (1000µg/ml). The stock solutions were kept under refrigeration conditions to be used for analysis. The working

solutions of different concentrations were prepared daily by serial dilution of the standard. Various volumes of the stock solutions in each case (2, 4, 6, 8 ml) were further diluted to 100 ml to obtain the working solutions of concentrations 50 ppm, 100 ppm, 150 ppm and 200 ppm. After filtering absorbencies were obtained and used to draw calibration curves. All analyses were done in triplicates.

2.2.2 Determination of trace elements

Each mushroom sample was air-dried at 105 °C overnight and crushed using a mortar and pestle into powder. Digestion of the mushroom samples was performed using a mixture of HNO₃:H₂O₂ (10:1, 12 ml g⁻¹ of sample) and heated at 100 °C for between 10-15 min. After cooling, the solution was made to 50 ml with deionized water after filtration. The amounts of Cr, Se and V were determined using an atomic absorption spectrophotometer. All analyses were done in triplicates.

2.2.3 *In vitro* simulated gastrointestinal digestion

The *in vitro* enzymolysis procedure simulating human gastrointestinal digestion was carried out in triplicate. Procedural blanks were run to check the presence of Se, Cr and V in the reagents. Mushroom samples (0.5 g) in a flask were incubated with 5 ml of gastric juice (1% w/v pepsin in 0.15 M NaCl, adjusted to pH 2 with HCl (37% v/v) and, after 1 min of vigorous shaking for initial degassing, the flask was placed in a mixing water bath at 37⁰ C for 4 h [30]. The solution was then adjusted to pH 6.8 with NaHCO₃. After adding 5 ml of intestinal juice (3% w/v pancreatin, 1.5% w/v amylase, 1% w/v bile salts in 0.15 M NaCl), solution was vigorously shaken for 1 min, degassed, and further incubated for 4 h at 37⁰ C under gentle shaking. The solution was then centrifuged at 8000 g and 4⁰ C for 15 min, the supernatant collected, filtered through 0.45 µm membranes, and stored at -80⁰ C until analysis. The amount of solubilized chromium, selenium, and vanadium in the supernatant was measured using atomic absorption spectrophotometer as a measure of their bioaccessibility and data analyzed using SPSS.

2.2.4 Stastical analysis

The data was analysed using SPSS program. ANOVA was used to determine whether there is significant difference in the

bioaccessibility of the trace elements. In this study the between sample estimate of variance is greater than within sample that is Chromium between group(BG) variation was 2.808 while the within group variation(WG) was 0.472;Vanadium BG(0.013) and WG(0.001) and selenium BG(116.691) and WG(44.222) to be explained by random error hence a one-tailed F-test was calculated to test whether it is significantly greater. From the table of critical values of F for a one tailed test (p=0.05) the critical value of F is 8.845. The calculated value of F (5.946 for chromium) and 2.639 is smaller than this the null hypothesis is accepted the sample means do not differ significantly while that for Vanadium is 9.094 is greater than the critical value. A significant result in one way ANOVA can arise for several reasons: one mean may differ from others, all the means may differ from each other, and the means may fall into two distinct groups. A simple way of deciding the reason for a significant result is to arrange the means in increasing order and compare the difference between adjacent values with a quantity called

least significant difference [31]. “The least significant difference method is not entirely rigorous, it can be shown that it leads to rather too many significant difference. However it is a simple follow up test when ANOVA has indicated that there is a significant difference between means” (Harris, 2007).In this study the lowest mean was 0.450 and the highest was 3.501.The least significant difference test is used in the context of ANOVA when the F-ratio suggests rejection of the null hypothesis. Turkey, Students-Newman-Keuls (SNK) and Bonferroni Post Hoc test are used for reporting the range tests and pairwise multiple comparisons to determine means that differ in ANOVA. Range tests identify homogeneous subsets of means that are not different from each other. Pairwise multiple comparisons test the difference in each pair of means and yield a matrix where asterics indicates significantly different group means at alpha level of 0.05.In this study Bonferroni post Hoc test was used (Table 1). The PO variety in chromium showed a significant difference (Table 1).

Table 1. Bioaccessible levels of Chromium, Vanadium and Selenium

Mushroom Variety	Level of Chromium Mean (Number) ug/g		Level of Vanadium Mean (Number) ug/g		Level of Selenium Mean (Number) ug/g	
	Bio	Raw	Bio	Raw	Bio	Raw
	PC	1.1221(3)	0.9506(3)	0.4159(3)	0.4498(3)	14.8466(3)
PO*	1.0025(3)*	3.510(3)*	0.4138(3)	0.4880(3)	8.91368(3)	1.1652(3)
PP	0.6789(3)	0.7478(3)	0.4237(3)	0.6057(3)	-1.3883(3)	0.7757(3)
PSC	0.4119(3)	0.9559(3)	0.4011(3)	0.4710(3)	-5.1000(3)	2.3377(3)

*. The mean difference is significant at the 0.05 level. Based on post hoc Turkey HSD

3. RESULTS AND DISCUSSION

3.1 Levels of Trace Elements in Oyster Mushrooms

Table 2. Mean levels of trace elements in Oyster mushrooms

Mushroom variety	Concentration (µg/g)		
	Chromium	Vanadium	Selenium
	Mean±SD	Mean±SD	Mean
PSC	1.551±1.661	0.471±1.481	2.338±0.055
PP	1.798±1.472	0.606±3.341	0.776±0.066
PO	3.501±0.857	0.488±0.946	1.165±0.01
PC	1.430±0.657	0.450±2.472	1.110±0.06

PO: *Pleurotus ostreatus*, PSC: *Pleurotus Saju-sajur Caju*, PP: *Pleurotus pulmonarius*, PC: *Pleurotus cintrinopileatus*

The mean trace element concentrations ($\mu\text{g/g}$) in dry weight in the mushroom samples were Chromium PSC(1.551),PP(1.798),PO(3.501),and PC(1.430).Vanadium concentrations in the four species were PS(2.338),PP(0.776),PO(1.165) and PC(1.110).Selenium levels were PSC (0.471), PP(0.606),PO(0.488) and PC(0.450) (Table 2). The precision of the results was evaluated based on the standard deviation of the results of triplicate samples($n=3$) analyzed under the same conditions $\text{mean}\pm\text{SD}$. The mean concentration of chromium was highest in PO and lowest in PP. Selenium concentration was highest in PSC and lowest in PP. Vanadium concentration was highest in PC and lowest in PP. The values are comparable to those discussed by authors [32-38], (Bano et al.1981), [39,40].

3.2 Trace elements Bioaccessibility

“Absorption of selenium occurs throughout gastrointestinal tract, with duodenum as a major site” [41]. Selenium absorption is very efficient, normally in the range of 26.3% - 97% [42,43], with higher retention of organic selenium compound than inorganic selenium compounds [44]. “Very little is known about mechanism of selenium absorption, it may involve both carrier mediated process for organic compound and diffusion-controlled process for inorganic selenium compound” [45].

The amounts of selenium bioaccessible in the four oyster mushrooms are as shown in Table 3 are PO-86.94%, PC-92.52% which is high and is comparable to literature value [42,43] and thus is potentially bioavailable. The other varieties, PS and PP had undetected levels. The soluble extract obtained after *In vitro* simulated gastrointestinal digestion of the selenized mushroom contained about 106gSe dry weight. This means that 75% of the Se taken up by the mushroom was solubilized in conditions simulating human gastrointestinal digestion and thus was potentially bioavailable [46]. Se content of mushrooms is generally higher than that of most vegetables [47] but it is very variable.

“Se that is not solubilized after gastrointestinal digestion might be present in form of indigestible Se-containing polysaccharides. For instance, it has been shown that part of the Se in Se-enriched mycelia of *Pleurotus ostreatus* is

associated with chitin-containing structures in cell walls” [48]. “Formation of Se-containing polysaccharides might explain the low Se bioavailability found elsewhere for other species” [49,50]. Limited bioavailability of Se in these species might be the result of a low bioaccessibility due to a larger incorporation of Se in mushroom polysaccharides compared to *Pleurotus ostreatus* in the conditions of the present study. Bioavailability of selenium from PO, PF, PSC, and PE was found to range between 0.011 and 0.512mg/100 g [37,51].

The bioaccessible vanadium in the four oyster mushrooms in this study was PS-85.16%,PP-69.95%,PO-84.80% and PC-92.46% (Table 3).Many studies confirm that of the total dietary vanadium ingested, less than 5% is absorbed by the gastrointestinal tract (GI) [52-54].Other studies claim greater than 10% of ingested vanadium may have resulted in greater than normal absorption efficiencies [55]. The majority of the 5% dietary vanadium absorbed is taken up by the upper GI tract [56].

The bioaccessible chromium levels in the four oyster mushrooms were PS-26.56%,PP-37.59%,PO-28.63% and PC-78.50% (Table 3).These values are comparable to those in literature *in vitro* and *in vivo* studies in rats have shown that about 80% Cr in the blood is associated with transferring [57]. “Chromium is absorbed together with other metal ions in the gut through the unsaturated passive transport. The absorption process depends on the Cr content in the diet and on the chemical form of this element and other food components” [58]. “The efficiency of this process is very low with the average absorption ranging from 0.4-2.5%” [59]. “The absorption process depends on the Cr content in the diet and on the chemical form of this element and other food components” [60]. “Organic sources of Cr (picolinate, or propionate-methionine salt) are much better absorbed than inorganic forms (oxides), and lead to the increase of these compounds’ concentration in tissues” [61-63]. “However, other factors present in the diet show a significant impact on the amounts of Cr absorbed from the gastrointestinal tract. Starch, simple sugars, ascorbic acid, oxalic acid, nicotinic acid, some amino acids, aspirin increase absorption of this element” [64-67] while “high concentrations of phosphate, calcium, magnesium, titanium, zinc, vanadium and iron reduce the rate of this process” [64,68].

Table 3. Bioaccessibility of elements in four oyster mushrooms

Mushroom variety	Bioaccessibility (%)		
	Chromium	Vanadium	Selenium
PSC	26.56	85.16	ND
PP	37.59	69.95	ND
PO	28.63	84.80	86.94
PC	78.50	92.46	92.52

ND: Not Detected

“More than 80% of Cr is removed from the body in the form of urine, while the remaining part of this element is excreted via faeces and sweat” [69]. “In humans, consumption of large amounts of sugar, exhaustive physical exercise, pregnancy, and lactation leads to increased Cr excretion in the urine” [70-81].

4. CONCLUSIONS

The study indicates that the oyster mushroom contains fairly high levels of the elements, but their bio accessibility differs in all the varieties. The extent of bio accessibility of the elements was affected by the variety although selenium remained least bio accessible in all varieties. Vanadium was better bio accessible in the four oyster mushrooms, while chromium was better bio accessible in the PC variety.

ACKNOWLEDGEMENT

I am grateful to Africa Development Bank (ADB) and National Research Fund (NRF) for funding my study and the technical support in Jomo Kenyatta University.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Heaney RP. Factors influencing the measurement of bioavailability taking calcium as a model. *J Nutr.* 2001;131(4);Suppl:1344S-8S.
2. Onyambu ZM, Nawiri MP, Nyambaka HN, Noah NM. In vitro bioaccessibility of the vitamin B series from thermally processed leafy African indigenous vegetables. *J Food Qual.* 2021;2021:1-8. Available: <https://doi.org/10.1155/2021/5540724>
3. Nambafu R, Swaleh S, Nyambaka H. Bioavailability studies of vitamin A and E in

indigenous vegetables and their potential use in the management of HIV and AIDS. *Adv Res.* 2021;22(2):36-44.

4. Guerra A, Etienne-Mesmin L, Livrelli V, Denis S, Blanquet-Diot S, Alric M. Relevance and challenges in modeling human gastric and small intestinal digestion. *Trends Biotechnol.* 2012;30(11): 591-600.
5. Lusi P, Nyambaka H, Mbakaya CF, Maseta E, Bwete V, Murungi J. Bioavailability studies of trace elements in a potential food formulation for use in the management of HIV and AIDS. *Int J Pure Appl Chem.* 2013;8(1):47-53.
6. Wienk KJ, Marx JJ, Beynen AC. The concept of iron bioavailability and its assessment. *Eur J Nutr.* 1999;38(2):51-75.
7. Etcheverry P, Grusak MA, Fleige LE. Application of in vitro bioaccessibility and bioavailability methods for calcium, carotenoids, folate, iron, magnesium, polyphenols, zinc, and vitamins B6, B12, D, and E. *Front Physiol.* 2012;3:1-21.
8. Courraud J, Berger J, Cristol JP, Avallone S. Stability and Bioaccessibility of different forms of carotenoids and vitamin A during in-vitro digestion. *Food Chem.* 2013;1362: 871-7.
9. Fernández-García E, Carvajal-Lérida I, Pérez-Gálvez A. In vitro bioaccessibility assessment as a prediction tool of nutrient efficiency. *Nutr Res.* 2009;29(11):751-60.
10. Nawiri MP, Nyambaka HN, Murungi JI. Sundried cowpeas and Amaranthus leaves recipe improves beta-carotene and retinol in serum and hemoglobin concentration among preschool children. *Eur J Nutr.* 2013;52(2):582-9.
11. Bano Z. Nutritive value of Indian mushrooms and medicinal practices. *Eco Bot.* 1976;31:367-71.
12. Samsudin NP, Abdullah N. Edible mushrooms from Malaysia: a literature review on their nutritional and medicinal properties. *Int Food Res J.* 2019;26(1): 11-31.

13. Kalač P. A review of chemical composition and nutritional value of wild growing and cultivated mushrooms. *J Sci Food Agric.* 2013;93(2):209-18.
14. Li H, Tian Y, Menolli J, Ye L, Karunarathna SC, Perezmoreno J, Rahmn MM, Rashid MH, Phengsintham P, Rizal L, Kasuya T. Reviewing the worlds edible mushroom species: a new evidence-based classification system. *Compre Rev Food Sci.* 2021;20(2):1982-2014.
15. Murugesan S. Sustainable food security-edible and medicinal mushroom. *Sustainable Agriculture towards food security.* Singapore. Springer. 2017: 185-96.
16. Chang S, Wasser S. The cultivation and environmental impact of mushroom. Vol. P43. New York: Oxford University Press; 2017.
17. Ho LH, Zulkifli NA, Tan TC. Edible mushrooms: nutritional properties potential nutraceutical values in food product development. An introduction to mushroom. London: In tech Open. 2020;19-36.
18. Kalac P, Svoboda L. A review of trace element concentrations in edible mushrooms. *Food Chem.* 2000;69(3): 273-81.
19. Aletor V. Compositional studies on edible tropical species of mushrooms. *Food Chem.* 1995;54(3):265-8.
20. Mattila P, Könkö K, Eurola M, Pihlava JM, Astola J, Vahteristo L et al. Contents of vitamins, mineral elements, and some phenolic compounds in cultivated mushrooms. *J Agric Food Chem.* 2001;49(5):2343-8.
21. Rudawska M, Leski T. Macro and micro elemental contents in fruiting bodies of wild mushrooms from the Netecka forest in west – central Poland. *Food Chem.* 2005;92(3):499-506.
22. Ganesan K, Xu B. Anti-obesity effects of medicinal and edible mushrooms. *Molecules.* 2018;23(11):2880.
23. Anderson RA, Cheng N, Bryden NA, Polansky MM, Cheng N, Chi J et al. Elevated intakes of supplemental chromium improve glucose and insulin variables in individuals with type 2 diabetes. *Diabetes.* 1997;46(11):1786-91.
24. Shechter Y. Insulin-mimetic effects of vanadate. Possible implications for future treatment of diabetes. *Diabetes.* 1990; 39(1):1-5.
25. Shamberger RJ. The insulin-like effects of vanadium. *J Adv Med.* 1996;9:121-31.
26. Akbaraly TN, Arnaud J, Rayman MP, Hininger-Favier I, Roussel AM, Berr C et al..A. Plasma selenium and risk of dysglycemia in an elderly French population: results from the prospective epidemiology of Vascular Ageing Study. *Nutr Metab Lond.* 2010;18: 71-121.
27. Becker DJ, Reul B, Ozcelikay AT, Buchet JP, Henquin JC, Brichard SM. Oral selenate improves glucose homeostasis and partly reverses abnormal expression of liver glycolytic and gluconeogenic enzymes in diabetic rats. *Diabetologia.* 1996;39(1):3-11.
28. WHO. Geneva: World Health Organization [guideline]: sodium intake for adults and children; 2012.
29. Rajarathnam S, Shashirekha MN. Mushroom nutraceuticals," in *Advances in preservation and processing technologies of fruits and vegetables.* New Delhi, India: New India Publishing Agency; 2011.
30. Crews HM, Clarke PA, Lewis DJ, Owen LM, Strutt PR, Izquierdo A. Investigation of selenium speciation in vitro gastrointestinal extracts of cooked cod by high-performance liquid chromatography-inductively coupled plasma mass spectrometry. *J Anal At Spectrom.* 1996;11(12):1177-82.
31. Miller JC, Miller JN. *Statistic for Analytical chemistry.* 2nd ed, Ellis Horwood limited publisher. Chichester; 1988.
32. Patil SS, Ahmed SA, Telong SM, Baig MMV. The nutritive values of pleurotusostreatuscultivaters of different agro wastes. *Innov Romenian Food Biotechnol.* 2010;7:66-76.
33. Yang JH, Lin HC, Mau JL. Non-volatile taste components of several commercial mushrooms. *Food Chem.* 2001;72(4):465-71.
34. Ahmed SA, Kadam JA, Mane VP. Biological efficiency and nutritional contents of *Pleurotus florida*. Singer cultivated on different agro-wastes. *Nat Sci.* 2009;7(1):44-8.
35. Alam N, Amin R, Khan A, Ara I, Shim MJ, Lee MW et al. Nutritional analysis of cultivated mushrooms in Bangladesh - *Pleurotusostreatus*, *Pleurotussajorcaju*, *Pleurotus florida* and *Calocybe indica*. *Mycobiology.* 2008;36(4):228-32.

36. Gosh N, Chakravarty DK. Predictive analysis of the protein quality of *Pleurotuscitrinopileatus*. J Food Sci Technol. 1990;27(4):236-8.
37. Gąsecka M, Mleczek M, Siwulski M, Niedzielski P. Phenolic composition and antioxidant properties of *Pleurotus ostreatus* and *Pleurotus eryngii* enriched with selenium and zinc. Eur Food Res Technol. 2016;242(5):723-32.
38. Kortei NK, Wiafe-Kwagyan M. Comparative appraisal of the total phenolic content, flavonoids, free radical scavenging activity and nutritional qualities of *Pleurotus ostreatus* (EM-1) and *Pleurotostreous*(P-31) cultivated on rice (*Oryzae sativa*) straw in Ghana. J Advbiol Biotechnol. 2015;3(4):153-64.
39. Jegadeesh R, Lakshmanan H, Kabyeul J. Cultivation of pink oyster mushroom *Pleurotus djamor* var. *roseus* on various agro-residues by low cost technique. J Mycopathol Res. 2018;56(3): 213-20.
40. Ijeh I, Okwujiako IA, Nwosu PC. Phytochemical composition of *Pleurotus tuber regium* and effect of its dietary incorporation on body weights and serum triacylglycerols in albino mice. J Med Plants Res. 2009;3(11):939-43.
41. Humaloja T, Mykkänen HM. Intestinal absorption of ⁷⁵Se-labeled sodium selenite and Selenomethionine in chicks: effects of time segment selenium concentration and method of measurement. J Nutr. 1986;116(1): 142-8.
42. Durcos V, Arnanda J, Tahiri M, Coudray C, Baratt.F., Bouteloipdemange C, Brown F, Rayssiguier Y, Roussel AM. Influence of Short- Chain Fracto Oligosaccharides on Absorption of Cu, Zn and Se in Healthy Postmenopausal Women. J. Am. Coll. Nutr. 2005;24:30-37.
43. Thompson CD, Burton CE, Robinson MF. On supplementing the selenium intake of New Zealander. Short experiments with large doses of selenite or selenomethionine. Br J Nutr. 1978;39: 587-97.
44. Finley JW. The retention and distribution by healthy young men of stable isotopes of selenium consumed as selenite, selenate or hydroponically, brown broccoli dependent on the isotopic form. J Nutr. 1999;129(4):865-71.
45. Ardüser F, Wolfram S, Scharrer E. Active absorption of selenate by rat ileum. J Nutr. 1985;115(9):1203-8.
46. Hur SJ, Lim BO, Decker EA, McClements DJ. In vitro human digestion models for food applications. Food Chem. 2011;125(1):1-12.
47. Rayman MP, Infante HG, Sargent M. Food-chain selenium and human health: spotlight on speciation. Br J Nutr. 2008;100(2):238-53.
48. Serafin Muñoz AH, Kubachka K, Wrobel K, Gutierrez Corona JF, Yathavakilla SK, Caruso JA et al. Se-enriched mycelia of *Pleurotus ostreatus*: distribution of selenium in cell walls and cell membranes/cytosol. J Agric Food Chem. 2006;54(9):3440-4.
49. Chansler MW, Mutanen M, Morris VC, Levander OA. Nutritional bioavailability to rats of selenium in Brazil nuts and mushrooms. Nutr Res. 1986;6(12):1419-28.
50. Mutanen M. Bioavailability of selenium in mushrooms, *Boletus edulis*, to young women. Int J Vitam Nutr Res. 1986;56(3):297-301. PMID 3781756.
51. Tang C, Hoo PC, Tan LT. Golden needle mushroom: a culinary medicine with evidenced-based biological activities and health promoting properties. Front Pharmacol. 2006;7:474.
52. Curran GL, Azarnoff DL, Bolinger RE. J Clin Invest. 1990;38:1251-126.
53. Byrne AR, Kosta L. Vanadium in foods and in human body fluids and tissues. Sci.J. Nutr. 1978;115:1203-8.
54. Nielsen FH. Trace Minerals in foods. New York: Marcel Dekker; 1988. p. 357-428.
55. Nielsen FH, Uthus EO 1990. Vanadium in biological systems. Physiology and biochemistry. Kluwer. London: Academic Press.51-62.
56. Patterson BW, Hansard II, Ammerman R, Henry LA, Zech, Fisher WR. Am J Physiol. 1986;251:325-32.
57. Feng W, Li B, Liu J, Chai Z, Zhang P, Gao Y et al. Study of chromium-containing proteins in sub cellular fractions of rat liver by enriched stable isotopic tracer technique and gel filtration chromatography. Anal Bioanal Chem. 2003; 375(3):363-8.
58. Dowling HJ, Offenbacher EG, Pi-Sunyer FX. Absorption of inorganic trivalent chromium from the vascular perfused rat

- small intestine. J Nutr. 1989;119(8): 1138-45.
59. European Commission. Opinion of the scientific committee on food on the tolerable; 2003. Upper Intake Level of Trivalent Chromium (expressed on 4 April 2003). Available:<http://ec.europa.eu/food/fs/sc/scf/out197-en.pdf> (access: 2010.04.29).
 60. Anderson RA, Bryden NA, Polansky MM, Gautschi K. Dietary chromium effects on tissue chromium concentrations and chromium absorption in rats. J Trace Elem Exp Med. 1996;9(1):11-25.
 61. Ohh SJ, Lee JY. Dietary chromium-methionine chelate supplementation and animal performance. Asian Australas J Anim Sci. 2005;18(6):898-907.
 62. Wang MQ, He YD, Lindemann MD, Jiang ZG. Efficacy of Cr (III) supplementation on growth, carcass composition, blood metabolites, and endocrine parameters in finishing pigs. Asian Australas J Anim Sci. 2009;22(10):1414-9.
 63. Zha LY, Xu ZR, Wang MQ, Gu LY. Effects of chromium nanoparticle dosage on growth, body composition, serum hormones and tissue chromium in Sprague-Dawley rats. J Zhejiang Univ Sci B. 2007;8(5):323-30.
 64. Chen NS, Tsai A, Dyer IA. Effect of chelating agents on chromium absorption in rats. J Nutr. 1973;103(8):1182-6.
 65. Davis ML, Seaborn CD, Stoecker BJ. Effects of over-the-counter drugs on chromium retention and urinary excretion in rats. Nutr Res. 1995;15(2):201-10.
 66. Offenbacher EG. Promotion of chromium absorption by ascorbic acid. Trace Elem Elect. 1994;11:178-81.
 67. Samanta S, Haldar S, Ghosh TK. Production and carcass traits in broiler chickens given diets supplemented with inorganic trivalent chromium and an organic acid blend. Br Poult Sci. 2008;49(2):155-63.
 68. Hill CH. Mineral interrelationships. Trace elements in human health and disease. New York: Academic Press. 1976;281-300.
 69. Ducros V. Chromium metabolism, a literature review. Biol Trace Elem Res. 1992;32:65-77.
 70. Anderson RA, Bryden NA, Polansky MM, Richards MP. Chromium supplementation of turkeys: effects on tissue chromium. J Agric Food Chem. 1989;37(1):131-4.
 71. Dennis Chasteen ND. The biochemistry of vanadium. Struct Bond. 1983;53:105-38.
 72. Chen J, Gaikwad V, Holmes M, Murray B, Povey M, Wang Y et al. Development of a simple model device for in vitro gastric digestion investigation. Food Funct. 2011;2(3-4):174-82.
 73. Combs JR. Selenium in global food systems. Br J Nutr. 2001;85:1517-547.
 74. Gao Y, Walder K, Sunderland T, Kantham L, Feng HC, Quick M et al. Elevation in Tanis expression alters glucose metabolism and insulin sensitivity in H4IIE cells. Diabetes. 2003;52(4):929-34.
 75. Kustin K, Macara IG. The new biochemistry of vanadium. Comments Inorg Chem. 1982;2(1-2):1-22.
 76. Silva SO, Costa SMG, Clemente E. Chemical composition of *Pleurotus pulmonarius*, substrates and residue after cultivation. Braz Arch Biol Technol. 2002;45(4):531-5.
 77. Suzuki KT, Doi C, Suzuki N. Metabolism of ⁷⁶Se-methylselenocysteine compared with that of ⁷⁷Se-selenomethionine and ⁸²Se-selenite. Toxicol Appl Pharmacol. 2006; 217(2):185-95.
 78. Thomson CD. Selenium speciation in human body fluids. Analyst. 1998;123(5): 827-31.
 79. Whanger PD. Selenocompounds in plants and animals and their biological significance. J Am Coll Nutr. 2002;21(3): 223-32.
 80. Whanger PD. Selenium and its relationship to cancer: an update. Br J Nutr. 2004;91(1):11-28.
 81. Yoneda S, Suzuki KT. Detoxification of mercury by selenium by binding of equimolar. Toxicol Appl Pharmacol. 1997;143(2): 274-80.

© 2023 Nambafu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96473>