

Hormonal Contraceptive Uptake and Socio-Demographic Pattern of Acceptors in a Family Planning Clinic of a Tertiary Health Facility in Rivers State, Nigeria

P. A. Awoyesuku^{1*} and B. O. A. Altraide¹

¹Department of Obstetrics and Gynaecology, Rivers State University Teaching Hospital, 6-8 Harley Street, Old G.R.A, Port-Harcourt, P.M.B. 5064, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Author PAA designed the study, performed the statistical analyses and wrote the first draft of the manuscript. Author BOAA assisted in data collection, managed the analyses of the study and literature searches. Both authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Abdelmonem Awad M. Hegazy, Professor, Department of Anatomy and Embryology, Faculty of Medicine, Zagazig University, Egypt.

Reviewers:

(1) Dr. Mohammed Ismail Khan, ESIC Medical College, India.

(2) M. V. Chandramathi, Symbiosis Law School, India.

(3) Suzy Reiter, Mid-County Health Center, USA.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50322>

Original Research Article

**Received 10 April 2019
Accepted 24 June 2019
Published 02 August 2019**

ABSTRACT

Background: Hormonal methods of birth control are a safe and reliable way to prevent pregnancy for most women. Their uptake rate in comparison to other contraceptive methods in our environment has not been well documented.

Objective: To determine the uptake of hormonal contraceptives and assess socio-demographic characteristics related to the choice, among acceptors in a tertiary health facility in Rivers State.

Methodology: A hospital-based cross-sectional study was adopted. A sample size of 124 was used. New clients were consecutively recruited over a 12-month period. Demographic data (age, parity, educational level, marital status) and contraceptive-related data (choice of contraceptive method, reasons for use) were obtained and analyzed using SPSS version 20.0.

Results: A total of 124 new female clients were recruited with median age of 34 years. Of these, 92(74.2%) accepted hormonal methods while 32 (25.8%) chose non-hormonal methods. Among the hormonal contraceptive acceptors, 94.6% (n=87) used implants, 4.3% (n=4) used injectable and 1.1% (n=1) used oral contraceptive pill. Bivariate analysis of socio-demographic factors and hormonal uptake among the acceptors was statistically significant for marital status, educational level and reason for contraception.

Conclusion: We found a substantial uptake of hormonal contraceptives, mainly implants. The uptake was particularly pronounced among married women with higher educational level and whose reason for contraception is completed family size.

Keywords: Hormonal contraceptives; contraceptive uptake; new acceptors; tertiary hospital; Rivers State.

1. INTRODUCTION

Hormonal methods of birth control contain estrogen and progestin, or progestin only; they are a safe and reliable way to prevent pregnancy for most women. Hormonal methods include subdermal implants, injectable, pills, Mirena IUCS, vaginal ring or skin patch. Non hormonal methods include copper IUCD, condoms, cervical cap, diaphragm, and sponge.

Hormonal contraception, particularly oral pills and injectable (depot medroxyprogesterone acetate), is the most popular and widely available form of contraception in Kenya [1]. Hormonal contraceptives are a safe and effective form of contraception for women <40 years, women with superficial venous thrombosis / varicose veins, later postpartum period, breastfeeding mothers after 6 months and HIV-infected women [2].

Contraceptive implants are one of the most effective family planning methods available and well-accepted worldwide [3,4]. They are long term hormonal contraceptives and a better option for women in sub-Saharan Africa due to its effectiveness and convenience [5,6]. Jadelle is a set of two flexible cylindrical implants, each containing 75 micrograms of progestin-levenorgestrel, while the Implanon contains 68 micrograms of Etonogestrel [7]. They are sealed with adhesives, sterilized and inserted in a superficial plane beneath the skin of the upper arm under aseptic condition and can be effective for five years in the case of Jadelle but three years for Implanon [8,9]. Unlike other hormonal delivery systems, they do not cause unnecessary peaks in progestin levels and do not use estrogen, and thus their health risks are minimal [10,11].

The injectable contraceptive method was the most preferred method of birth control among

women of reproductive age group in Warri (South-South), South-Eastern and Kano (Northern) Nigeria [12,13,14]. Injectable contraceptive was also the method of choice among women seeking terminal fertility control in South-Western Nigeria [15]. Contrary to the forgoing, intrauterine contraceptive device was the most chosen method in 74.6% of the married women attending family planning of a tertiary institution in Oshogbo, Nigeria [16].

The contraceptive CHOICE project, one of the largest prospective cohort studies of women in the US seeking reversible contraception, found that when the barriers of cost, access and knowledge are removed, women chose the most effective and least user-dependent methods [17]. In their study, the uptake of long-acting reversible contraceptives increased from 5% at the baseline to 75% during the study.

Studies elsewhere have reported high levels of hormonal contraceptive utilization in various groups [18,19,20]. The uptake rate of hormonal contraceptives in comparison to other contraceptive methods in our environment has not been well documented. This study seeks to determine the uptake of hormonal contraceptives and assess socio-demographic characteristics related to the choice, among acceptors in a tertiary health facility in Rivers State.

2. METHODOLOGY

A hospital-based cross-sectional study was adopted. Using the formula for cross-sectional studies [21], a sample size of 124 was attained based on the alpha level of 0.05, contraceptive uptake of 15% in Nigeria from 2013 Demographic Health Survey (DHS) [22], precision of 7.5% and non-response rate of 30%.

New clients at the family planning clinic were consecutively recruited over a 12-month period.

Demographic data (age, parity, educational level, and marital status) and contraceptive-related data (choice of contraceptive method and reasons for use) were obtained in the study. Data was collected by the authors from the records directly and analyzed using the Statistical Package for Social Sciences (SPSS) version 20.0.

The hormonal contraceptive uptake among the new clients at the family planning clinic was expressed in percentages. Bivariate analysis employed Chi square/Fisher's exact statistics in determining significant differences in the demographic pattern. Statistical significance was set at $P < 0.25$ for bivariate analysis.

Statistically significant variables on bivariate analysis were entered into a multivariate analysis model. Multivariate analysis was done using unconditional binary logistic regression model. The dependent variable was hormonal contraceptive uptake (categorized as Yes/No) while the demographic characteristics and reasons for contraceptive use comprised the independent variables. Odds ratio and 95% confidence intervals were determined and $P < 0.05$ following multivariate analysis were considered statistically significant.

3. RESULTS

The study comprised of a total of 124 new female clients with median age of 34 years and an age range of 20 to 49 years. Of these, 92(74.2%) accepted hormonal methods while 32 (25.8%)

chose non-hormonal methods (Fig. 1). Among the hormonal contraceptive acceptors, 94.6% (n=87) used implants, 4.3% (n=4) used injectable and 1.1% (n=1) used oral contraceptives (Fig. 2).

Table 1 shows the socio-demographic profile of clients, majority of whom had secondary education & above (90.3%); were married (93.4%) and had parity of 3 & above (71.8%). Majority of the clients (55.7%) also had completed family size as their reason for contraception. Bivariate analysis of socio-demographic factors and hormonal uptake among the acceptors was statistically significant ($p < 0.25$) for marital status, educational level and reason for contraception; however, these were not statistically significant ($p < 0.05$) on multivariate analysis as shown in Table 2.




Fig. 1. Distribution of type of contraceptive accepted by the clients




Fig. 2. Distribution of hormonal contraceptive accepted by the clients

Table 1. Socio-demographic characteristics and uptake of hormonal contraceptive (Bivariate analysis)

Variables	Hormonal uptake		Total n (%)
	Yes n (%)	No n (%)	
Age category			
≤ 34 years	51 (72.9)	19 (27.1)	70 (100.0)
> 34 years	41 (75.9)	13 (24.1)	54 (100.0)
	<i>Chi-Square = 0.150; P = .699</i>		
Educational level			
Below secondary	9 (75.0)	3 (25.0)	12 (100.0)
Secondary & higher	83 (74.1)	29 (25.9)	112 (100.0)
	<i>Fisher's exact P = 1.000</i>		
Marital status			
Married	85 (72.6)	32 (27.4)	117 (100.0)
Single	7 (100.0)	0 (0.0)	7 (100.0)
	<i>Fisher's exact P = .189*</i>		
Parity			
Para ≤ 2	29 (82.9)	6 (17.1)	35 (100.0)
Para > 2	63 (70.8)	26 (29.2)	89 (100.0)
	<i>Chi-Square = 1.912; P = .167*</i>		
Completed family size			
Yes	48 (69.6)	21 (30.4)	69 (100.0)
No	44 (80.0)	11 (20.0)	55 (100.0)
	<i>Chi-Square = 1.740; P = .187*</i>		
Child spacing			
Yes	42 (79.2)	11 (20.8)	53 (100.0)
No	50 (70.4)	421 (29.6)	71 (100.0)
	<i>Chi-Square = 1.234; P = .267</i>		

*Statistically significant $P < 0.25$ **Table 2. Socio-demographic characteristics and uptake of hormonal contraceptive (Multivariate analysis)**

Variables*	Coefficient (B)	Odds ratio (OR)	95% CI	p value
Parity				
Para > 2	0.477	1.61	0.45 – 5.83	0.467
Para ≤ 2 ^R		1	1	
Completed family size				
Yes	0.287	1.33	0.45 – 3.96	0.605
No ^R		1	1	

Multivariate analysis of socio-demographic variables ($P < 0.25$ on bivariate analysis) and hormonal uptake among family planning acceptors

*Marital status was removed from the multivariate analysis since the column for singles that did not use hormonal was zero. The significant findings on bivariate analysis ($p < 0.25$), following logistic regression (multivariate analysis) were not significant @ $p < 0.05$

4. DISCUSSION

Our study shows that the uptake of hormonal contraceptives at our Centre as a percentage of all acceptors was high (74.2%). This is similar to the findings by Abaasa A. et al. [19] of 51.1% and Balkus J. et al. [20] of 72%. Of the new hormonal contraceptive users, a huge majority, 87 (94.6%) accepted the subdermal implantable

methods above the rest. This is contrary to the findings of the above studies where a majority of their acceptors chose the injectable, followed by the contraceptive pills, above the implants. Other studies [18,23] also recorded a low uptake of implants.

The high uptake of the implants in our study may be attributed to the fact that it is readily available

in our Centre; there is no lack of medical personnel skilled at insertion and the reason for contraception, which is mainly, completed family size, demanding long-acting methods. The non-availability of the commodity at the time of counseling and sometimes lack of medical personnel that is skilled at implant insertion when needed makes its need to be unmet in some Centers [18].

Uptake of hormonal contraceptive was higher among the younger age group just like in other studies [19]; however, this was not statistically significant when compared to the non-acceptors. Majority of our clients were highly educated with 90.3% haven attained secondary education and above. Both younger age group and lower educational status have been associated with lower contraceptive use in Uganda [24].

Bivariate analysis of socio-demographic factors and hormonal uptake among the acceptors was statistically significant ($p < 0.25$) for marital status, educational level and reason for contraception; however these were not statistically significant ($p < 0.05$) on multivariate analysis. Like this study, marital status was associated with hormonal contraceptive use in the study by Balkus J et al. [20].

5. CONCLUSION

We found a substantial uptake of hormonal contraceptives, mainly implants, among women attending our family planning clinic. The uptake was particularly pronounced among married women with higher educational level and whose reason for contraception was completed family size. Promotion and provision of hormonal contraception greatly increases the proportion of women using a reliable method of contraception.

CONSENT

As per international standard, patient's written consent has been collected and preserved by the authors.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kenya Demographic and Health Survey; 2003.
Available:<http://www.cbs.go.ke>
2. WHO. Medical Eligibility Criteria for Contraceptive use. 5TH ed. Geneva. World Health Organization; 2015.
3. National Population Commission, NDHS (2008) National Population Commission. ORC Marco, Calverton.
4. Engender Health. SEED Assessment Guide for Family Planning Programming. Engender Health, New York; 2011. Available:<http://www.engenderhealth.org/pubs/family-planning/seed-assessment-guide-for-family-planning.php>
5. Hubacher D, Olawo A, Manduka C, Kiare J. Factors associated with uptake of subdermal contraceptive implants in a young Kenyan Population. *Contraception*. 2011;84:413-417.
Available:<http://dx.doi.org/10.1016/j.contraception.2011.02.007>
6. Ladipo OA, Akinso SA. Contraceptive implants. *African Journal of Reproductive Health*. 2005;9:16-30.
Available:<http://dx.doi.org/10.2307/3583156>
7. Ali MM, Cleland J, Shah IH. Causes and consequences of contraceptive discontinuation: Evidence from 60 demographic and health surveys. World Health Organization, Geneva; 2012.
8. Darney PD. Hormonal implants: Contraception for a new century. *American Journal of Obstetrics and Gynecology*. 1994;170:1536.
Available:[http://dx.doi.org/10.1016/s0002-9378\(94\)05016-7](http://dx.doi.org/10.1016/s0002-9378(94)05016-7)
9. Bangladesh Institute of Research for Promotion of Essential & Reproductive Health and Technologies, Dhaka. 1993; 47:569-582.
10. Jacobstein R. Fostering change in medical settings: Some considerations for family planning programmes. *IPPF Medical Bulletin*. 2009;43:3-4.
11. Wickstrom J, Jacobstein R. Contraceptive security: Incomplete without long-acting and permanent methods of family planning. *Studies in Family Planning*. 2011;42:291-298.
Available:<http://dx.doi.org/10.1111/j.1728-4465.2011.00292.x>

12. Oghenekaro AV. Contraceptive choice amongst women in Warri, Nigeria. *Int. J Life Sci Pharma Res.* 2012;2:35-9.
13. Chigbu B, Onwere S, Aluka C, Kamanu C, Okoro O, Feyi-Waboso P. Contraceptive choices of women in rural South-Eastern Nigeria. *Niger. J Clin Pract.* 2010;13:195-9.
14. Yakasai IA, Yusuf AM. Contraceptive choices amongst women in Kano, Nigeria: A five-year review. *J Med Trop.* 2013;15: 113-6.
15. Adekunle DA, Afolabi AF, Adeyemi AS. Terminal fertility control: Clients own reason for the choice of the contraceptive method. *Open J Obstet Gynecol.* 2013;3: 706-9.
16. Adeyemi AS, Adekunle DA, Komolafe JO. Pattern of contraceptives choice among the married women attending the family planning clinic of a tertiary health institution. *Niger J Med.* 2008;17:67-70.
17. McNichollas C, Tessa M, Peipert JF. The contraceptive CHOICE project round up; what we did and what we learned. *Clin Obstet Gynecol.* 2014;57(4):635-643. Available: <https://dx.doi.org/10.1097/GFR.0000000000000070>
18. Madugu NH, Abdul MA, Bawa U, Kolawole B. Uptake of hormonal implants contraceptive in Zaria, Northern Nigeria. *Open Journal of Obstetrics and Gynecology.* 2015;5:268-273. Available: <http://dx.doi.org/10.4236/ojog.2015.55039>
19. Abaasa A, Gafos M, Anywaine Z, Nunn A, Crook A, Lewin J, et al. Uptake of Hormonal Contraceptives and correlates in a phase III clinical trial in rural South Western Uganda. *Reproductive Health.* 2017;14:36. Available: <http://dx.doi.org/10.1186/s12978-017-0296-3>
20. Balkus J, Bosire R, John-Stewart G, Mbori-Ngacha D, Schiff MA, Wamalwa D, et al. High uptake of postpartum hormonal contraception among HIV-1-seropositive women in Kenya. *Sex Transm Dis.* 2017; 34(1):25-29. Available: <http://dx.doi.org/10.1097/01.olq.000218880.88179.36>
21. Kirkwood BR, Sterne JAC. Sample size determination. *Essential Medical Statistics 2nd Edition.* Blackwell Publishers, United Kingdom. 155.
22. Maryland, USA. National Population Commission (NPC) Nigeria and ICF International. *Nigeria Demographic and Health Survey 2013.* Abuja, Nigeria, and Rockville; 2014.
23. Hubacher D, Mavranezouli I, McGinn E. Unintended Pregnancy in Sub-Saharan Africa: Magnitude of the problem and Potential Role of Contraceptive Implants to alleviate it. *Contraception.* 2008;78:73-78.
24. Uganda MOH. ICF, Centers for Disease Control and Prevention Uganda, USAID, WHO Uganda, Uganda Bureau of Statistics, Uganda Virus Research Institute. *Uganda AIDS indicator survey.* Kampala: MOH and ICF. 2011;1-238.

© 2019 Awoyesuku and Altraide; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50322>