



Causes of Neonatal Morbidity and Mortality in Bingham University Teaching Hospital, Jos, Plateau State, Nigeria

M. M. Ihekaike^{a*}, M. Shehu^a, A. O. Jimoh^a, U. Onyenuche^a,
A. Uhunmwangho-Courage^a and A. M. Yakubu^a

^a Department of Paediatrics, Bingham University/ Bingham University Teaching Hospital, Jos, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author MMI participated in conceptualization of the study, literature search, study design, data acquisition, data and statistical analysis, manuscript preparation, editing and review. Author MS participated in study design, data and statistical analysis, manuscript editing and review. Author AOJ participated in study design, data and statistical analysis, and manuscript editing and review. Author UO participated in definition of intellectual content and manuscript editing and review. Author AUC participated in definition of intellectual content, data and statistical analysis, and manuscript editing and review. Author AMY participated in definition of intellectual content and manuscript editing and review. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJPR/2023/v13i2267

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103601>

Original Research Article

Received: 24/05/2023

Accepted: 28/07/2023

Published: 05/08/2023

ABSTRACT

Introduction: Neonatal morbidity and mortality continue to be serious public health issues in developing countries, where it also has a considerable impact on the under-five mortality rate.

Aims: To determine the causes of morbidity and mortality in neonates admitted to the inborn Special Care Baby Unit (SCBU) at Bingham University Teaching Hospital (BHUTH), Jos.

*Corresponding author: Email: marciaihekaike@yahoo.co, marcia.ihekaike@binghamuni.edu.ng;

Study Design: Cross Sectional Descriptive study.

Place and Duration of Study: Inborn Neonatology Unit of the Department of Paediatrics, Bingham University teaching Hospital, Jos, Plateau State, Nigeria.

Methodology: A retrospective analysis of the records of all the babies admitted into the inborn SCBU of BHUTH over an 18-month period of January 2021 to June 2022 was done. We included 195 babies (99 males, 96 females)

Results: A total of 196 patients were admitted into the inborn SCBU of BHUTH during the period reviewed and 195 (99.5%) had complete records. There were 99 (50.8%) males and 96 (49.2%) females with M: F of 1.03:1. The leading causes of admission were prematurity (51.3%), asphyxia (21.5%) and neonatal jaundice (10.8%). Nine (4.6%) infants had severe birth asphyxia, while 33 (16.9%) had moderate birth asphyxia. Twenty-eight deaths were recorded over the 18-month period giving a mortality rate of 14.4%. Prematurity (71.4%) and severe birth asphyxia (17.9%) were the leading causes of death. Of all the deaths that occurred, 9 (32.1%) occurred within the first 24 hours of hospitalization, while 3 (10.7%) of them died after 7 days of admission.

Conclusion: In our community, the neonatal mortality rate is still very high. This death rate can be decreased with the support of concerted efforts to guarantee appropriate antenatal care, close monitoring of labour, and enhanced neonatal unit facilities for newborn care.

Keywords: Morbidity; mortality; neonatal; prematurity.

1. INTRODUCTION

The most crucial time for a child's survival is the first month [1]. In 2020, 2.4 million children died in their first month of life worldwide. Every day, over 6700 newborns die accounting for nearly half (47%) of all under-five mortality, up from 40% in 1990 [2].

Neonatal morbidity and mortality continue to be serious public health issues in developing countries, where it also has a considerable impact on the under-five mortality rate because worldwide under-5 mortality is dropping faster than neonatal mortality [2,3].

Sub-Saharan Africa has the world's highest neonatal mortality rate (27 deaths per 1000 live births), accounting for 43% of worldwide newborn deaths, followed by Central and Southern Asia (23 deaths per 1000 livebirths), accounting for 36% of global newborn deaths [2].

Nigeria had a neonatal mortality rate of 34.9 deaths per 1,000 live births in 2021. Nigeria's neonatal mortality decreased considerably over time, going from 63.7 deaths per 1,000 live births in 1972 to 34.9 deaths per 1,000 live births in 2021 [3].

The majority of neonatal deaths (75%) occur during the first week of life, and almost 1 million babies died during the first 24 hours in 2019. The majority of neonatal deaths in 2019 were caused by preterm delivery, birth asphyxia, infections, and birth abnormalities [2].

Periodic neonatal auditing is essential since disease patterns vary from place to place and over time, even within the same location [4]. Assessing the pattern of morbidity and mortality among newborns coming to this facility is very desirable and crucial in determining the quality of medical care offered and identifying inadequacies in their overall management. It will also help policymakers plan more effectively.

Therefore, this study aims to ascertain the causes of morbidity and mortality in neonates admitted to the inborn Special Care Baby Unit (SCBU) at Bingham University Teaching Hospital (BHUTH), Jos between January 2021 and June 2022.

2. MATERIALS AND METHODS

This is a retrospective study of all newborns admitted into the inborn SCBU of BHUTH between January 2021 and June 2022. Being a tertiary centre, the hospital serves as a referral centre for patients from neighbouring states of Kaduna and Bauchi and even from other secondary and tertiary facilities within the state. The newborn unit is located within the maternity complex and caters for the needs of babies born in the hospital's delivery suite. The SCBU is equipped with 4 incubators, 2 cots, 4 phototherapy units, and one radiant warmer. There is a nursing room attached to the unit where mothers of the babies on admission can breast feed their babies without having to enter into the main ward. Basic facilities for care in the form of intravenous fluid therapy, phototherapy,

naso-gastric feeding and respiratory support in the form of intermittent AMBU bagging and oxygen therapy as well as non-invasive respiratory support [bubble continuous positive airway pressure (BCPAP) are available. The hospital has a separate outborn SCBU which is located within the paediatric complex.

The medical records of these babies were reviewed and the following data were abstracted: the age at presentation, the gender, the reason for admission, duration of admission, and the outcome of admission.

Socioeconomic classification of the study population was not ascertained as most of the case record files did not capture information on parental educational backgrounds and occupation/employment status.

Ethical clearance for the study was obtained from Health Research Ethics Committee of Bingham University Teaching Hospital with reference number NHREC/21/05/2005/00971.

The data retrieved were entered into a personal computer and analyzed with SPSS Version 21.

3. RESULTS

During the period reviewed, a total of 196 patients were admitted into the unit of which 195 (99.5%) had complete records and were included in the study. There were 99 (50.8%) males and 96 (49.2%) females with M: F of 1.03:1. Table 1 shows the diagnostic criteria for some common neonatal problems seen in our hospital.

The causes of neonatal admissions are as shown in Table 2.

The other causes of morbidity accounting for 9.7% of the cases are as listed in Table 3.

A total of 152 (77.9%) babies were discharged home and 15 (7.7%) were Discharged Against Medical Advice (DAMA). Twenty-eight deaths were recorded over the 18-month period giving a mortality rate of 14.4%. Of these, 19 (67.9%) were males and 9 (32.1%) females giving a male to female ratio of 2.1:1. The sex specific mortality rate was 19.2% for males and 9.4% for females. The case fatality rate (CFR) was highest for severe birth asphyxia as shown in Table 4.

The duration of hospital admission for babies that died are as shown in Table 5.

Table 1. Diagnostic criteria for some common neonatal problems

Disease	Criteria
Prematurity	Babies born before 37 completed weeks of gestation
Severe birth asphyxia	Apgar score 0-3, [*] As well as clinical signs of Central nervous system depression
Moderate birth asphyxia	Apgar score 4-6
Jaundice	Clinical jaundice plus serum bilirubin greater than the norm on the normogram for the babies gestational age and weight
Neonatal sepsis	Maternal or neonatal risk factors for sepsis with laboratory evidence of infection [†]

^{*}Facilities for blood gases and PH are not available

[†]Complete blood count, blood and surface swab cultures, urine culture

Table 2. Causes of admission into inborn SCBU of Bingham University Teaching Hospital

Causes of Admission	Frequency (%)
Prematurity	100 (51.3)
Moderate birth asphyxia	33 (16.9)
Severe birth asphyxia	9 (4.6)
Neonatal Jaundice	21 (10.8)
Neonatal sepsis	13 (6.7)
Others	19 (9.7)
Total	195 (100.0)

Table 3. Other causes of admission into SCBU

Causes of Morbidity	N (%)
Meconium Aspiration Syndrome	3 (15.8)
Aspiration syndrome	1 (5.3)
Congenital Pneumonia	2 (10.5)
Transient tachypnoea of the newborn	5 (26.3)
Congenital anomalies	2 (10.5)
Hypoglycaemia	4 (21.0)
Haemorrhagic disease of the newborn	1 (5.3)
Dehydration Fever	1 (5.3)
Total	19 (100.0)

Table 4. Causes of death and case fatality rate of morbidities

Morbidity	Total No. Admitted	Mortality (%)	CFR (%)
Prematurity	100	20 (71.4)	20.0
Severe Birth Asphyxia	9	5 (17.9)	55.6
Respiratory disorders	13	2 (7.1)	15.4
Congenital Malformations	2	1 (3.6)	50.0
Total	124	28	

CFR = case fatality rate

Table 5. Duration of hospitalization of newborns that died

Duration of Hospitalization	Frequency (%)
< 24 hours	9 (32.1)
1 – 3 days	9 (32.1)
4 -7 days	6 (21.4)
> 7 days	3 (10.7)
Missing record	1 (3.7)
Total	28 (100)

4. DISCUSSION

Neonatal death remains a major public health issue in underdeveloped nations, like Nigeria, despite technological advancement and interventions for improving the quality of life of newborns. The rates of neonatal morbidity and mortality reflect a country's socioeconomic status as well as the quality of its healthcare systems. These critical factors can help plan for better healthcare delivery [5].

This study showed a male preponderance in both the morbidity and mortality patterns which is consistent with findings from previous studies done both within Nigeria [4, 6-7] and in other countries including Ethiopia, Bangladesh and Pakistan [1,5,8-9]. The sex specific mortality rate for males in this study was as high as 19.2%, more than twice that of their female counterparts which was 9.4%. The increased risk of mortality in male neonates is not well understood, but

could be as a result of genetic differences. A study in Pakistan on gender variability in neonatal mortality had shown that the risks of stillbirths, and early neonatal mortality were higher among male infants than their female counterparts [10].

Prematurity accounted for over half of the admissions in our study. An otherwise well, term baby would not be admitted. Preterm infants have underdeveloped organs that can't support the child to live independently, making them vulnerable to several issues like hypoglycemia, infections, and temperature instability. As a result, they are more likely to be kept in an incubator or given kangaroo mother care (KMC) until the child is old enough to be sent home. Other studies have also shown that the commonest cause of morbidity is prematurity [6,8,11].

Prematurity, birth asphyxia and neonatal jaundice were the three most prevalent

morbidities recorded in this study. This pattern is similar to that of another study carried out in Abuja, Nigeria [6]. Other studies carried out in northern Nigeria reported birth asphyxia, neonatal sepsis and prematurity as the commonest indications for hospital admissions in newborns [7,12]. A number of studies have reported neonatal sepsis as the leading morbidity. This was however not the case in this study possibly because we recruited only inborn infants most of whom were admitted from birth as a result of prematurity or asphyxia [13-16]. The burden of neonatal sepsis in this study may have been reduced by the fact all the participants were delivered in the hospital.

The mortality rate of 14.4% in this study is high and comparable to findings from studies done in other parts of Nigeria that ranged between 13.3% and 16.9% [4,7,13,17-19] though some authors have reported mortality rate as high as 20.4% in Gusau [20] and 25.9% in Azare [15]. However, certain studies have reported lower mortality rate as low as 6.8% in Pakistan [8], 2.1% in China [21]. This pattern of mortality further highlights the need for concerted efforts to improve the quality of health care delivery both during the antenatal and perinatal period to reduce the rate of deaths from preventable causes.

Close to 90% of the neonates that died in this study died within the first week of life. This finding is similar to that from other studies [16,22] as well as the WHO report on newborn mortality [2]. Early neonatal death is usually linked to maternal health, thus emphasizing the need for close monitoring of pregnancies and skilled supervision of deliveries.

The DAMA rate in our study as 7.7% which is much higher than the 1.5% reported by Also et al [16] in Dutse, 1.7% by Onazi et al [23] in Gusau and 5.2% by Imoudu et al [15] in Azare. The reason for the high rate of DAMA in our facility may not be unrelated to the higher cost of assessing health care there, being that ours is a private University Teaching Hospital unlike these other studies [15,16,23] that were carried out in Government funded hospitals.

While acknowledging the restricted generalizability due to a small sample size and the exclusion of neonates born and died at home, the study provided valuable insights into institution-based neonatal deaths during the study period. Like other retrospective studies, the cost-effectiveness and time efficiency of utilizing

existing records contribute to the study's strengths as hospital records offer data for in-depth analysis of medical history, treatment, and outcomes. It also offers valuable insights into neonatal mortality within the hospital setting and serves as a foundation for future research.

5. CONCLUSION

In conclusion, neonatal morbidity and mortality have remained high in our community, and the primary causes, which are essentially preventable has remained the same over the years. There is need for the government and other relevant agencies to examine the factors responsible for this and implement measures to reduce neonatal morbidity and mortality in our community.

CONSENT

As per international standard, parental written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

Ethical clearance for the study was obtained from Health Research Ethics Committee of Bingham University Teaching Hospital with reference number NHREC/21/05/2005/00971.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Desalew A, Sintayehu Y, Teferi N, Amare F, Geda B, Worku T, et al. Cause and predictors of neonatal mortality among neonates admitted to neonatal intensive care units of public hospitals in eastern Ethiopia: A facility-based prospective follow-up study. *BMC Pediatr* 2020; 20(1):160. Available: <https://doi.org/10.1186/s12887-020-02051-7>
2. World Health Organization Newborn Mortality. Accessed 18th June, 2023. Available: <https://www.who.int/news-room/fact-sheets/detail/levels-and-trends-in-child-mortality-report-2021>
3. World Data Atlas. Nigeria – Neonatal Mortality Rate.

- Accessed 18th June, 2023.
Available:<https://knoema.com/atlas/Nigeria/Neonatal-mortality-rate>.
4. Ekwochi U, Ndu IK, Nwokoye IC, Ezenwosu OU, Amadi OF, Osuorah D. Pattern of morbidity and mortality of newborns admitted into the sick and special care baby unit of Enugu State University Teaching Hospital, Enugu state. *Niger J Clin Pract*. 2014;17(3):346-51. DOI: 10.4103/1119-3077.130238. PMID: 24714015.
 5. Ali MB, Maruf AA, Rashid MM. Pattern and outcome of admissions in a neonatal high dependency unit of a medical college hospital in Khulna, Bangladesh. *Mediscope* 2018;5(2):1-7. DOI:10.3329/mediscope.v5i2.37141.
 6. Audu LI, Otuneye AT, Mairami AB, Mukhtar-Yola M, Mshelia LJ. Determination of neonatal case-specific fatality rates in a tertiary health institution in North Central Nigeria. *BMC Pediatr* 2021;21(1):302. Available:<https://doi.org/10.1186/s12887-021-02778-x>.
 7. Mukhtar-Yola M, Iliyasu Z. A review of neonatal morbidity and mortality in Aminu Kano Teaching Hospital, northern Nigeria. *Trop Doct* 2007;37(3):130-2. DOI:10.1258/004947507781524683. PMID: 17716492.
 8. Ali SR, Ahmed S, Lohana H. Disease patterns and outcomes of neonatal admissions at a secondary care hospital in Pakistan. *Sultan Qaboos Univ Med J* 2013;13(3):424-8. PMID: 23984028;
 9. Asghar RM, Sharif M, Saheel K, Ashraf RR, Hussain A. An analysis of five years neonatal mortality in NICU of a Tertiary Care Hospital of Rawalpindi 2014-2019. *JRMC* 2020;24(4):328-33. Available:<https://doi.org/10.37939/jrnc.v24i4.1394>
 10. Aghai ZH, Goudar SS, Patel A, Saleem S, Dhaded DM, Kavi A, et al. Gender variations in neonatal and early infant mortality in India and Pakistan: A secondary analysis from the Global Network Maternal Newborn Health Registry. *Reprod Health*. 2020;17 Suppl 3:S178-89. Available:<https://doi.org/10.1186/s12978-020-01028-0>
 11. Shehu M, Dusu C, Davou K, Habila M, Mava Y, Eseigbe E, et al. Neonatal morbidity and mortality in Bingham University Teaching Hospital Jos, Plateau, North-Central Nigeria. *PMJG* 2022;9(2):95–8. DOI:<https://doi.org/10.60014/pmjpg.v9i2.233>
 12. Abdullahi UI. Neonatal morbidity and mortality in a Rural Tertiary Hospital in Nigeria. *CHRISMED J Health Res* 2018; 5(1):8-10
 13. Omoigberale AI, Sadoh WE, Nwaneri DU. A 4-year review of neonatal outcome at the University of Benin Teaching Hospital. Benin City. *Niger J Clin Pract* 2010; 13(3):321-5. PMID: 20857794.
 14. Ochoga MO, Abah RO, Michael A, Yaguoide LE, Onalo R, Idoko A, et al. Retrospective assessment of neonatal morbidity and mortality in the special care baby unit of a private health facility in Benue state, North Central Nigeria. *Niger J Paediatr* 2020;47(4):353-7. DOI:10.4314/njp.v47i4.9
 15. Imoudu IA, Ahmad H, Yusuf MO, Makarfi HU, Umar T. An analysis of neonatal morbidity and mortality in Azare North Eastern Nigeria. *J Dent Med Sci* 2014; 13(3):25-8. DOI:10.9790/0853-13312528
 16. Also U, Gwarzo GD. Pattern of morbidity and mortality among neonates seen in a tertiary hospital. *Sahel Med J* 2020; 23(1):47-50. DOI: 10.4103/smj.smj_21_19
 17. Eke BC, Ezomike UO, Chukwu BF, Chinawa JM, Korie FC, Chukwudi N. et al. Pattern of Neonatal Mortality in a Tertiary Health Facility in Umuahia, South East, Nigeria. *Int J Trop Dis & Health* 2014;4(2):136-46. DOI: 10.9734/IJTDH/2014/4985
 18. Ugwu GI. Pattern of morbidity and mortality in the newborn special care unit in a tertiary institution in the Niger delta region of Nigeria: A prospective study. *Glob Adv Res J Med Sci* 2012;1(6):133-8.
 19. Okechukwu AA, Achonwa A. Morbidity and mortality patterns of admissions into the Special Care Baby Unit of University of Abuja Teaching Hospital, Gwagwalada, Nigeria. *Niger J Clin Pract* 2009;12(4):389-94. PMID: 20329678.
 20. Garba BI, Muhammad AS, Mohammed BA, Obasi BO, Adeniji AO. A study of neonatal mortality in a specialist hospital in Gusau, Zamfara, North-Western Nigeria. *Int J Trop Dis* 2017;28(2):1-6. DOI: 10.9734/IJTDH/2017/38337

21. Xu Y, Zhu X, Wang H, Pan Z, Li X, Guo X, et al. Prevalence of major morbidities and outcome of all hospitalized neonates. A retrospective cohort study of Huai'an neonatal survivals. *J Matern Fetal Neonatal Med.* 2022;35(25):9800-10. DOI: 10.1080/14767058.2022.2054320. PMID: 35341440.
22. Udo JJ, Anah MU, Ochigbo SO, Etuk IS, Ekanem AD. Neonatal morbidity and mortality in Calabar, Nigeria: a hospital-based study. *Niger J Clin Pract.* 2008; 11(3):285-9. PMID: 19140371.
23. Onazi SO, Akeredolu FD, Yakubu M, Jiya MN, Hano IJ. Neonatal morbidity and mortality: A 5-year analysis at Federal Medical Centre, Gusau, Zamfara State, Northwest Nigeria. *Ann Med Res Pract* 2021;2:1-5. DOI:10.25259/ANMRP_8_2021

© 2023 Ihekaïke et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/103601>