



# **Reaction of Maize Genotypes against Fall Armyworm, *Spodoptera frugiperda* (Smith) in Madhya Pradesh, India**

**Shivani Suman <sup>a++\*</sup>, N. S. Bhaduria <sup>a#</sup>, S. P. S. Tomar <sup>a†</sup> and Pradyumn Singh <sup>b‡</sup>**

<sup>a</sup> Department of Entomology, RVSKV Gwalior, Madhya Pradesh, India.

<sup>b</sup> B.M. College of Agriculture, Khandwa, Madhya Pradesh, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. Authors SS and NSB conceived, designed the research and conducted experiment. Authors SS, SPST and SP contributed to analytical tools. Author SS wrote the manuscript. All authors read and approved the manuscript.*

## **Article Information**

DOI: 10.9734/IJECC/2023/v13i102633

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/104947>

**Original Research Article**

**Received: 04/06/2023**

**Accepted: 09/08/2023**

**Published: 14/08/2023**

## **ABSTRACT**

**Aims:** To identify the less susceptible varieties, allowing for targeted pest management strategies, reduced pesticide use, increased resilience to pest outbreaks, and long-term sustainable pest control solutions.

**Study Design:** Randomized Block Design.

**Place and Duration of Study:** Research Farm, College of Agriculture, Gwalior (Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior) during two consecutive years i.e., Kharif, 2021-22 and Kharif 2022-23.

<sup>++</sup> Ph.D. Scholar;

<sup>#</sup> Professor;

<sup>†</sup> Senior Scientist;

<sup>‡</sup> Scientist Entomology;

\*Corresponding author: E-mail: shivanisuman1996@gmail.com;

**Methodology:** In the experiment, twelve different varieties were sown on 17<sup>th</sup> July 2021 and 17<sup>th</sup> July 2022 during both the consecutive years respectively. Observations on the number of fall armyworm, *Spodoptera frugiperda* larvae and plant damage per cent were recorded from randomly selected ten plants at weekly intervals. Observations for FAW was also be recorded with visual scoring method. A numerical scale (0-9), also known as the Davis scale, was used to evaluate leaf damage.

**Results:** The result of both the year indicated that not all varieties were found to be completely resistant to fall armyworm. For the purpose of interpreting the results, all the varieties were categorized for their reaction based on average data of both the years (*Kharif* 2021 and *Kharif* 2022) of mean larval population, mean plant damage percent, and leaf damage scoring. Based on statistical categorization, it was determined that the varieties NWMH-2002 and JM 218 exhibited lower susceptibility. On the other hand, the varieties Dkc-9141, Maize 3845, PAC 740, Yashoda gold, PM 303, Maize Ranker, Maize 5402, and Maize 3046 were classified as moderately susceptible. Lastly, Sona 5101 and M-909 were identified as highly susceptible varieties.

**Conclusions:** It concluded that the varieties NWMH-2002 and JM 218 exhibited lower susceptibility against fall armyworm.

**Keywords:** Susceptibility; leaf damage; larval population; plant damage.

## 1. INTRODUCTION

Maize is referred to as the "Queen of Cereals" throughout the world due to its great genetic yield potential among cereals. Due to its wide genetic base and high level of genetic diversity, it is more adaptive and versatile in a variety of agro-climatic conditions. Maize is a member of the Poaceae family. It originated in South America and spread throughout the world. In 70 countries, including 53 developing countries, more than 100,000 ha of maize are grown [1]. According to Shah et al. [2], 100 g of the edible component of maize includes 71.88 g of carbohydrates, 8.84 g of protein, 4.57 g of fat, 2.15 g of crude fibre, 2.33 g of ashes, and different vitamins and minerals.

With an average productivity of 3199 kg/ha, maize is grown on 98.91 million ha of land in India on an annual average basis, producing 316.46 million tonnes [3]. Karnataka is the top producer of maize among the Indian states, followed by Madhya Pradesh, Maharashtra, Tamil Nadu, and West Bengal. With a productivity of 2763 kg/ha and a yield of 38.81 million tonnes, maize is grown on 14.05 million ha of land in Madhya Pradesh [3]. Some of the insect pests that target maize fields include cutworms, maize stem borer, white grub and chaffer beetles, armyworm, gram pod borer, wireworm, hairy caterpillars, etc. [4]. In India, the fall armyworm *Spodoptera frugiperda* was discovered on a maize field in August 2018 close to Shivammoga, Bangalore, Karnataka state [5]. The maize crop has been severely damaged by this new invasive species, which is posing an important threat to the nation's ability to produce maize. Plants are harmed when larvae of fall

armyworm (FAW) consume the leaf. Young larvae typically harm leaves by boring holes in them and primarily eating on the epidermis of the leaf. Small holes in the leaves can be used to identify leaf damage. Around the damaged region, the larvae are frequently simple to find. The larger larvae in the whorls eating on the maize cob or kernels can reduce the yield and quality of older plants. Due to lower yields, higher costs of inputs for pesticide applications, and the demand for more labour and resources to control pest infestations, farmers suffer financial losses. The identification of resistant or tolerant varieties of maize by pest control screening provides specific pest management techniques, less pesticide use, enhanced resilience to pest outbreaks, and long-term sustainable pest control solutions. Varieties of maize show varied degrees of tolerance or resistance to particular pests. Thus, the present study is meant to examine with the aim of screening out the susceptible varieties of maize against fall armyworm.

## 2. MATERIALS AND METHODS

This evaluation was conducted at the Entomological Research Farm of College of Agriculture, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior, M.P. during two consecutive years i.e., *Kharif*, 2021-22 and *Kharif* 2022-23. In the experiment, twelve varieties of maize were sown, maintaining 60 cm plant to plant and 20 cm row to row distance. The size of each replicated plot was maintained as 5.0m x 3.0m. After sowing, moderate irrigation was done to facilitate the initiation of germination. Fertilizers were applied at the rate of 120 kg N,

60 kg P and 30 kg K per hectare. For the observations of fall armyworm, the number of FAW larvae and damaged plants were counted from randomly selected ten plants at weekly intervals starting from germination till harvest of the crop. Also, the fall armyworm infestation on different maize varieties was noted on the basis of leaf damage under natural infestation. A numerical scale (0-9), also known as the Davis scale, was used to evaluate leaf damage [6] as described in the list 1.

### 3. RESULTS AND DISCUSSION

It was noteworthy from the results that none of the studied varieties were fully free of pest infestation. The data regarding number of larvae per plant, plant damage (%) and leaf damage rating scale are shown in the Tables 1 and 2.

#### 3.1 Average Larval Population during Kharif 2021 and Kharif 2022

Average mean least larval population was recorded in NWMH-2002 (0.87 larvae) followed by JM 218 (0.98). However highest larval population was recorded in Sona 5101 (1.89) (Table 1). For the purpose of interpreting the results, all the varieties were categorized for their reaction based on average data of both the years (*Kharif 2021 and Kharif 2022*) of mean larval population. The statistical formula  $\bar{X} \pm \sigma$  was used to categorize the varieties, with the average value being ( $\bar{X} = 1.43$ ) and the standard deviation being ( $\sigma = 0.30$ ). As a result, three separate groupings of larval population—below 1.13, between 1.13 and 1.73, and above 1.73—were identified. Based on this statistical categorization

method, it was determined that the varieties NWMH-2002 and JM 218 exhibited lower susceptibility. On the other hand, the varieties Dkc-9141, Maize 3845, PAC 740, Yashoda gold, PM 303, Maize Ranker, Maize 5402, and Maize 3046 were classified as moderately susceptible. Lastly, Sona 5101 and M-909 were identified as highly susceptible varieties.

#### 3.2 Average Plant Damage (%) during Kharif 2021 and Kharif 2022

Minimum plant damage % was found in NWMH-2002 (9.29%) which recorded as significantly less effective in comparison to rest of the varieties. However, maximum and significantly high larval population was found in Sona 5101 (38.69%) among all the varieties (Table 1). For the purpose of interpreting the results, all the varieties were categorized for their reaction based on average data of both the years (*Kharif 2021 and Kharif 2022*) of percent plant damage. The statistical formula  $\bar{X} \pm \sigma$  was used to categorize the varieties, with the average value being ( $\bar{X} = 22.62$ ) and the standard deviation being ( $\sigma = 8.31$ ). As a result, three separate groupings of average data of two year of plant damage (%) due to fall armyworm —below 14.31 (less susceptible), between 14.31 and 30.92 (moderately susceptible), and above 30.92 (highly susceptible)- were discovered. The varieties NWMH-2002 and JM 218 were found to be less susceptible. The varieties Dkc-9141, Maize 3845, PAC 740, Yashoda gold, PM 303, Maize Ranker, Maize 5402 and Maize 3046 were found as moderately susceptible and Sona 5101 and M-909 were highly susceptible.

List 1. Rating scale showing leaf damage

Rating	Explanation/definition of damage
0	No visible leaf damage;
1	Only pin-hole damage;
2	Pin-hole and small circular hole damage to leaves.
3	Pinholes, small circular lesions and a few small elongated (rectangular shaped) lesions of up to 1.3 cm in length present on whorl and furl leaves.
4	Several small to mid-sized 1.3 to 2.5 cm in length elongated lesions present on a few whorl and furl leaves
5	Several large elongated lesions greater than 2.5 cm in length present on a few whorl and furl leaves and/or a few small to mid-sized, uni-form to irregular shaped holes (basement membrane consumed) eaten from the whorl and/or furl leaves.
6	Several large elongated lesions present on several whorl and furl leaves and/or several large uniforms to irregular shaped holes eaten from furl and whorl leaves.
7	Many elongated lesions of all sizes present on several whorl and furl leaves and/or several large uniforms to irregular shaped holes eaten from the whorl and furl leaves.
8	Many elongated lesions of all sizes present on most whorl and furl leaves plus many mid to large sized uniform to irregular shaped holes eaten from the whorl and furl leaves.
9	Whorl and furl leaves almost totally destroyed.

**Table 1. Incidence of fall armyworm on different varieties of Maize during Kharif 2021 and Kharif 2022**

Varieties	Larval Population per plant			Plant damage (%)		
	2021	2022	Average	2021	2022	Average
Dkc- 9141	1.13 (1.28)*	1.23 (1.32)	1.18 (1.30)	14.52 (22.39)**	16.90 (24.28)	15.71 (23.35)
Maize 5402	1.57 (1.44)	1.70 (1.48)	1.63 (1.46)	27.38 (31.55)	27.14 (31.39)	27.26 (31.47)
NWMH-2002	0.83 (1.15)	0.90 (1.18)	0.87 (1.17)	8.57 (16.95)	10.00 (18.43)	9.29 (17.73)
PM 303	1.45 (1.40)	1.54 (1.43)	1.49 (1.41)	21.43 (27.57)	23.33 (28.87)	22.38 (28.23)
Sona 5101	1.83 (1.53)	1.95 (1.57)	1.89 (1.55)	37.14 (37.55)	40.24 (39.37)	38.69 (38.46)
PAC 740	1.34 (1.36)	1.43 (1.39)	1.38 (1.37)	17.86 (24.99)	19.76 (26.38)	18.81 (25.7)
Yashoda Gold	1.39 (1.37)	1.47 (1.40)	1.43 (1.39)	19.76 (26.39)	21.43 (27.57)	20.60 (26.98)
JM 218	0.95 (1.21)	1.01 (1.23)	0.98 (1.22)	11.67 (19.96)	13.33 (21.41)	12.50 (20.7)
Maize 3046	1.63 (1.46)	1.75 (1.50)	1.69 (1.48)	29.76 (33.05)	30.24 (33.26)	30.00 (33.21)
Maize ranker	1.54 (1.43)	1.63 (1.46)	1.58 (1.44)	24.52 (29.67)	25.48 (30.31)	25.00 (30)
M-909	1.71 (1.49)	1.82 (1.52)	1.77 (1.51)	33.10 (35.11)	33.81 (35.55)	33.45 (35.33)
Maize 3845	1.24 (1.32)	1.33 (1.35)	1.28 (1.34)	16.90 (24.26)	18.57 (25.52)	17.74 (24.9)
SEm ±	<b>0.01</b>	<b>0.01</b>	<b>0.01</b>	<b>0.85</b>	<b>0.53</b>	<b>0.59</b>
C.D. (at 5%)	<b>0.03</b>	<b>0.03</b>	<b>0.03</b>	<b>2.58</b>	<b>1.62</b>	<b>1.80</b>

\*figures in parentheses are square root transformed values; \*\*figures in parentheses are arc sine transformed values

**Table 2. Rating of leaf damage by fall armyworm on different varieties of Maize during Kharif 2021 and Kharif 2022**

Varieties	Leaf damage scoring		
	2021	2022	Average
Dkc- 9141	1.79	1.94	1.86
Maize 5402	3.69	3.90	3.79
NWMH-2002	1.23	1.43	1.33
PM 303	2.93	3.14	3.03
Sona 5101	5.03	5.18	5.10
PAC 740	2.29	2.42	2.35
Yashoda Gold	2.48	2.63	2.56
JM 218	1.37	1.60	1.49
Maize 3046	4.05	4.21	4.13
Maize ranker	3.19	3.44	3.31
M-909	4.39	4.55	4.47
Maize 3845	2.06	2.22	2.14
SEm ±	<b>0.06</b>	<b>0.02</b>	<b>0.04</b>
C.D. (at 5%)	<b>0.17</b>	<b>0.07</b>	<b>0.11</b>

### 3.3 Leaf Damage Scoring during Kharif 2021 and Kharif 2022

According to the data presented in Table 2, on different varieties, the leaf damage rating of all observations in the average data of both the year ranged from 1.33 to 5.10. NWMH-2002 had the lowest leaf damage rating (1.33), followed by JM 218 (1.49), although they were statistically different. However, Sona 5101 (5.10) has the highest leaf damage rating (Table 2). Despite the fact that the other cultivars scored in the middle of the class in regards to leaf damage. The order of leaf damage rating from lowest to highest among all the varieties was NWMH-2002 (1.33), JM 218 (1.49), Dkc-9141 (1.86), Maize 3845 (2.14), PAC 740 (2.35), Yashoda gold (2.56), PM 303 (3.03), Maize Ranker (3.31), Maize 5402 (3.79), Maize 3046 (4.13), M-909 (4.47) and Sona 5101 (5.10). For the purpose of interpreting the results, all the varieties were categorized for their reaction based on average data of both the years (*Kharif 2021* and *Kharif 2022*) of leaf damage scoring. The statistical formula  $\bar{X} \pm \sigma$  was used to categorize the varieties, with the average value being ( $\bar{X} = 2.96$ ) and the standard deviation being ( $\sigma = 1.17$ ) as shown in the Table 1. As a result, three unique groupings of leaf damage ratings were obtained: <1.80, between 1.80 - 4.13, and >4.13. According to this, the varieties NWMH-2002 and JM 218 have been found to be less susceptible. The varieties Dkc-9141, Maize 3845, PAC 740, Yashoda gold, PM 303, Maize Ranker, Maize 5402 and Maize 3046 were found as moderately susceptible and Sona 5101 and M-909 were highly susceptible.

During this investigation, twelve varieties of maize were screened against fall armyworm. The result of both the year indicated that not all varieties were found to be completely resistant to fall armyworm. For the purpose of interpreting the results, all the varieties were categorized for their reaction based on average data of both the years (*Kharif 2021* and *Kharif 2022*) of mean larval population, mean plant damage percent, and leaf damage scoring. Based on statistical categorization by using ( $\bar{X} \pm \sigma$ ), it was determined that the varieties NWMH-2002 and JM 218 exhibited lower susceptibility. On the other hand, the varieties Dkc-9141, Maize 3845, PAC 740, Yashoda gold, PM 303, Maize Ranker, Maize 5402, and Maize 3046 were classified as moderately susceptible. Lastly, Sona 5101 and M-909 were identified as highly susceptible varieties as mentioned in the Tables 1 and 2.

The current findings are consistent with Gowda et al. [7] who observed that of the twenty-two maize genotypes evaluated, CML 71, CML 67, and DMRE63 had considerably reduced leaf damage scores to fall armyworm of 3.93, 4.00, and 4.17, respectively. CML 71, CML 67, DMRE63, CML 561, AEBY-1, CML 335, CML 345, and CML 337 had mean leaf damage scores of 3.93, 4.00, 4.17, 4.36, 4.42, 4.57, 4.72, and 4.80, respectively, and were classified as moderately resistant genotypes. When Paul and Deole [8] tested maize genotypes for fall armyworm, they recorded similar findings. The genotype DKC-9190 exhibited the lowest leaf damage score of 2.36, suggesting resistance, whereas the genotype NK-30 had the highest leaf damage score of 8.21, indicating significant susceptibility. Somashekhar [9] discovered that partially resistant hybrids LG 36607, P3550, Tata Dhanya, and S 6668 had significantly higher leaf damage scores of 3.38, 3.66, 3.80, and 7.13 at V6 leaf stage when compared to leaf damage scores of LG 36607 (2.60), P3550 (3.28), Tata Dhanya (2.93), and S 6668 (4.33) at V8 stage. In line with the present result, Nelly et al. [10] who evaluated five farmer-grown varieties of maize viz. Pioneer, Pertiwi, Bisi 18, NK212, and NK7328 found that *S. frugiperda* attacked all of the varieties. Chiriboga et al. [11] evaluated six maize cultivars under natural infestation, and in contrast to the present results, they found a slight increase in the scores for plant damage at the end of the season, suggesting that these cultivars may have undergone successive colonization after the second sampling of the season. In the three last samplings of the season under natural infestation, the cultivars "WH5070," "Rachar," and "SC Duma 430" had stable plant damage ratings.

## 4. CONCLUSION

It was concluded that the varieties NWMH-2002 and JM 218 exhibited lower susceptibility. On the other hand, the varieties Dkc-9141, Maize 3845, PAC 740, Yashoda gold, PM 303, Maize Ranker, Maize 5402, and Maize 3046 were classified as moderately susceptible. Lastly, Sona 5101 and M-909 were identified as highly susceptible varieties.

## ACKNOWLEDGEMENTS

We are grateful to Head of the Department of Entomology and Dean, College of Agriculture, Gwalior for providing necessary facilities to carry out the research work.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Dowswell C, Paliwal RL and Cantrell RP. Maize in the third world. CRC Press. 2019;282.
2. Shah TR, Prasad K and Kumar P. Studies on physicochemical and functional characteristics of asparagus bean flour and maize flour. Conceptual frame work & innovations in agroecology and food sciences. 2015;103-105.
3. Anonymous. Selected State/Season- wise Area, Production and Productivity of Maize in India (2020-2021). Available:<https://www.indiastatagri.com/table/agriculture/selected-state-wise-area-production-productivity-m/1423779>
4. Arifie U, Bano P, Ahad I, Singh P, Dar ZA, Badri Z, Maqbool S, Aafreen S, Kumar R. Insect pests of maize at different altitudes of north Kashmir, J&K. Journal of Entomology and Zoology Studies. 2019;7:1123-1128.
5. Ganiger PC, Yeshwanth HM, Muralimohan K, Vinay N, Kumar ARV, Chandrashekara K. Occurrence of the new invasive pest, fall armyworm, *Spodoptera frugiperda* (JE Smith) (Lepidoptera: Noctuidae), in the maize fields of Karnataka, India. Current Science. 2018;115(4):621-623.
6. Davis FM, Ng SS, Williams WP. Visual rating scales for screening whorl-stage corn for resistance to fall armyworm. Technical bulletin (Mississippi Agricultural and Forestry Experiment Station). 1992;186:1-9.
7. Gowda MA Prajwal, Sekhar JC, Soujanya PL, Yathish KR, Rahman SJ, Mallaiah B. Screening of maize genotypes against fall armyworm, *Spodoptera frugiperda* (J.E. Smith) under artificial infestation. Biological Forum – An International Journal. 2022;14(2a):249-254.
8. Paul N, Deole S. Screening of maize genotypes against fall army worm, *Spodoptera frugiperda* (Smith) with reference to plant morphological characters at Raipur (Chhattisgarh). Journal of Entomology and Zoology Studies. 2020;8(4):580-587.
9. Somashekhar CM. Screening of different hybrids and evaluation of insecticides against fall armyworm, *Spodoptera frugiperda* (J.E. Smith) on maize. M.Sc. (Ag.) Thesis. University of Agricultural and Horticultural Sciences, Shivamogga; 2020.
10. Nelly N, Hamid H, Lina EC, Yunisman. The use of several maize varieties by farmers and the infestation of *Spodoptera frugiperda* (Noctuidae: Lepidoptera). IOP Conference Series: Earth and Environmental Science. 2021;662, 012020.
11. Chiriboga Morales X, Tamiru A, Sobhy IS, Bruce TJA, Midéga CAO, Khan Z. Evaluation of African maize cultivars for resistance to fall armyworm *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) Larvae. Plants. 2021;10:392. DOI:<https://doi.org/10.3390/plants10020392>

© 2023 Shivani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/104947>