



# Effect of Different Pre-Sowing Treatments on Germination Characteristics of Strawberry Guava (*Psidium cattleianum* Sabine)

Prashita Rao <sup>a++</sup>, K. A. Shanmugasundaram <sup>a#\*</sup>,  
I. Muthuvel <sup>at</sup>, K. Vanitha <sup>b‡</sup> and K. Gurusamy <sup>c^</sup>

<sup>a</sup> Department of Fruit Science, Tamil Nadu Agricultural University, Coimbatore, India.

<sup>b</sup> Department of Crop Physiology, Tamil Nadu Agricultural University, Coimbatore, India.

<sup>c</sup> Department of Food Process Engineering, Tamil Nadu Agricultural University, Coimbatore, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/v13i102997

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106392>

Original Research Article

Received: 05/07/2023

Accepted: 08/09/2023

Published: 15/09/2023

## ABSTRACT

**Aim:** The aim of the present study was to investigate the effect of different pre-sowing treatments on germination characteristics of strawberry guava (*Psidium cattleianum* Sabine).

**Study Design:** The experiment consisted of eleven treatments and three replications laid out in a Completely randomized design (CRD).

**Place and Duration of Study:** The study was conducted at Orchard Nursery, Department of Fruit

<sup>++</sup>M.Sc. Scholar;

<sup>#</sup>Assistant Professor (Horticulture);

<sup>†</sup>Professor (Horticulture);

<sup>‡</sup>Assistant Professor (Crop Physiology);

<sup>^</sup>Assistant Professor (Food Process Engineering);

<sup>\*</sup>Corresponding author: E-mail: shanmugasundaramka@tnau.ac.in;

Science, Horticulture College and Research Institute, Tamil Nadu Agricultural University, Coimbatore during April-June 2023.

**Methodology:** Seeds of *Psidium cattleianum* (strawberry guava) were collected from guava germplasm from an orchard and treated with different pre-sowing chemicals. After pre-sowing treatment, the seeds were sown and various germination characteristics and growth parameters were observed and recorded.

**Results:** The seeds treated with GA3 500 ppm for 30 minutes took minimum days to germinate (20), with higher germination percentage (79.33%), maximum vigour index I, and vigour index II (708.6 and 584.23). The growth parameters such as seedling height, stem girth, and number of leaves, primary root length, number of lateral roots were recorded higher in GA3 500 ppm followed by Thiourea 1%. The biomass production of seedlings was also higher in seeds treated with GA3 500 ppm.

**Conclusion:** The seeds treated with GA3 500 ppm for 30 minutes recorded higher germination characteristics, growth parameters and biomass production. So for quicker propagation of strawberry guava to be used as rootstock, seeds can be treated with 500 ppm GA3 to obtain the large number of seedlings for the use of rootstock.

**Keywords:** Rootstock; germination percentage; vigour index; presowing; biomass production.

## 1. INTRODUCTION

Guava (*Psidium guajava* L.) is one of the major fruit crop grown in India. It belongs to the family Myrtaceae and the order Myrtales. In India, it is the fourth most important fruit crop in terms of area and production after mango, banana, and citrus. Guava occupies an area of 0.3 million hectares with an annual production of 4.5 million tonnes. In North India, Uttar Pradesh is considered as "Home of Guava" which stands first in the production of guava (5983.59 thousand MT). Besides, it is cultivated in Bihar, Maharashtra, Gujarat, Madhya Pradesh, Tamil Nadu, Rajasthan, Karnataka, West Bengal, Orissa, Kerala and Punjab. In Tamil Nadu, guava is cultivated in an area of 14.44 thousand hectares with a production of 363.07 thousand MT [1].

In recent years, guava fruits had a huge demand in the national as well as international market because it has a rich basket of bioactive compounds and nutrients. Though there is a huge demand in the market, but guava production is being hindered due to various problems such as the guava nematode-wilt complex. Nematodes and Fusarium species jointly parasitize the tree's root system, causing guava decline [2]. Unlike, the persistence of chemical nematicides is usually short in the tropics, due to run-off and leaching. Besides, many of the effective nematicides were banned from the markets owing to their risks to human health and the environment. So, alternative control measures should be adopted to control the nematode. No source of resistance to the

nematode was found in *Psidium guajava*; however, sources of resistance were also identified in genotypes of *P. cattleianum* by Carneiro et al. [3] and Miranda et al. [4].

Almeida et al. [5] identified *Psidium cattleianum* as resistant to this nematode and the disease complex. Therefore, to alleviate the nematode problem, alternative low-input, cost-effective and environment friendly management strategies will be utilizing *P. cattleianum* as root stock. Allahabad Safeda scions were grafted onto *Psidium cattleianum* were the tallest and gave the greatest yields [6]. When used as rootstocks *P. cattleianum* and *P. friedrichsthalianum* were compatible with *P. guajava* cv. Paluma [3]. Vishwakarma et al. [7] screened different wild species of guava with *M. incognita* inoculation and reported that *Psidium cattleianum* var. *cattleianum* and *P. cattleianum* var. *lucidum* were immune to both *M. incognita* (0 RGI score) and *F. oxysporum* f. sp. *psidii* (0% wilting).

*Psidium cattleianum* Sabine is a slow-growing evergreen tree, 2–4 m tall, commonly called cattley guava, Chinese guava, purple guava, and strawberry guava. It is indigenous to the lowlands of eastern Brazil and parts of Uruguay, Strawberry guava fruit has a pleasant, strawberry-like flavour when ripe, hence its common name. *P. cattleianum* is resistant to the nematode *Meloidogyne incognita*. No disease pathogens have been recorded on *P. cattleianum*, and it is known to be resistant to the rust fungus *Puccinia psidii* which affects *P. guajava*. Costa et al. [8] conducted a research on identification of secondary metabolites related to *Psidium* spp. resistance against nematode

*Meloidogyne enterolobii* and data analysis revealed that the substances regulated upon infection in were primarily hydrolysable tannins, lignans and also some terpenes, which may also act as inhibitors of *M. enterolobii* inoculation.

These characteristics of strawberry guava make it the one of ideal rootstock for guava. However, the availability of this root stock is not meeting the actual demand of grafted planting material in guava. Uowolo and Denslow [9] stated that *P. cattleianum* does not form a persistent seed bank, and loses viability after 6.5 months. Based on the economic potential of the species and its ecological importance, studies are needed on conservation, and adequate storage [10], and So, we conducted research to study the effect of different pre-sowing treatments on the rootstock strawberry guava (*Psidium cattleianum* Sabine) for overcoming dormancy, thus enabling their availability and use throughout the year for grafting with guava.

## 2. MATERIALS AND METHODS

The experiment was conducted in orchard nursery, Department of Fruit science, Tamil Nadu Agricultural University, Coimbatore. The experiment was laid out in a Completely Randomized Design with eleven treatments and three replications.

Seeds were extracted from ripened fruits harvested from the mother plants maintained at guava germplasm in the Orchard which were collected and outdoors dried for one day and packed in butter paper. Then dried seeds were used for the study. Seeds were treated with different pre-sowing treatments for the germination study. The different treatments are: T1: GA<sub>3</sub> 200 ppm T2: GA<sub>3</sub> 300 ppm T3: GA<sub>3</sub> 400 ppm T4: GA<sub>3</sub> 500 ppm for 30 minutes. T5: Thiourea 0.5% T6: Thiourea 1% for 12 hours T7: KNO<sub>3</sub> 0.5% T8: KNO<sub>3</sub>: 1% for 12 hours T9: 5% Hcl for 10 minutes T10: 10% Hcl for 10 minutes T11: Control in water for 30 minutes.

For each replication, 50 seeds were sown in cocopeat media in 98 called portray, so a total of 1650 seeds were sown. Portrays were kept inside the shade net for effective germination.

### 2.1 Number of Days taken for Germination and Germination Percentage

The germinating seeds were observed daily and the day on which the first germination of the seed

was seen was considered as the number of days taken for germination for each treatment.

The germination percentage was calculated using the following formula:

$$\text{Germination Percentage (\%)} = \frac{\text{Number of seeds germinated}}{\text{Total number of seeds germinated}} \times 100$$

## 2.2 Growth Parameters

The growth parameters such as seedling height (cm), length of the primary root (cm), number of leaves and stem girth (mm) were measured from selected five seedlings of each treatment and mean was computed. The observations were recorded at 60 days from sowing.

The stem girth was measured with the help of Vernier calliper at 3 cm height from the base of the seedling.

Leaf area was calculated by using conversion factor (k) given by Nava et al. [11]

$$\text{Leaf area (cm}^2\text{)} = \text{Length of leaf} \times \text{Width of leaf} \times k$$

Where

$$k = 0.72581647$$

## 2.3 Biomass Parameters

### 2.3.1 Fresh weight of total seedling, shoot (g), and root (g)

The five selected seedlings of each department were uprooted 90 days after sowing and adhered soil to the roots was washed thoroughly seedlings were shade dried to remove excess water and the weight of the total seedling, root, and shoot was taken on electric balance was recorded and mean was computed in grams.

### 2.3.2 Dry weight of total seedling, shoot (g) and root(g)

Freshly weighed seedlings, shoots, and roots were chopped and labelled properly and were kept in brown paper bags and oven dried at 60°C weight was measured and the average value was computed.

## 2.4 Seedling Vigour Index

Vigour index defines the state of health of seedlings and also the production of biomass of

the plant. It was calculated using the formula given by Abdul-Baki and Anderson [12]:

**Seedling Vigour Index I(cm)** = Germination percentage × Length of the seedling (cm)

**Seedling Vigour Index II (cm)** = Germination percentage × Dry weight of seedling (g)

Vigour index I was calculated at 60 days after sowing and Vigour index was calculated at 90 days after sowing.

This experiment was laid out in completely randomized design (CRD). The data collected were statistically analyzed by computing ANOVA at a 5% level of significance using R studio software version 4.3.1 with suitable packages.

### 3. RESULTS AND DISCUSSION

#### 3.1 Number of Days taken for Germination and Germination Percentage

The minimum number of days taken for germination in the present investigation was seen in GA<sub>3</sub> 500 ppm (20.0) and maximum days were recorded in control (31.0). These results can be attributed to the activity of GA<sub>3</sub> on embryos causing de novo synthesis of enzymes such as amylase which is utilized for embryo growth and thereby enhancing the germination. Uravshi et al. (2020) also obtained the same

results when guava seeds were treated with GA<sub>3</sub>.

The germination percentage was recorded higher in seeds treated with GA<sub>3</sub> 500 ppm (79.33%) followed by Thiourea 1% (74.33%) and minimum germination percentage was recorded in control (34.66%) (Fig. 1). The higher germination percentage in seeds treated with GA<sub>3</sub> can be attributed to the fact that GA helps in breaking the seed dormancy and increases the activity of endogenous auxins and gibberellins like substances [13]. These results are in close conformity with Gurung et al. [14] in passion fruit and by Chandra and Govind (1990) in guava. Higher germination in thiourea-treated seeds can be due to the antagonized effect of thiourea on inhibitors present in the seed or due to the activity of increased cytokinin by thiourea which helps in overcoming the inhibiting effect of the seed coat [15].

#### 3.2 Growth Parameters of Seedlings

The growth parameters of strawberry guava seedlings showed a positive response to the pre-sowing treatments and are presented in (Table 2). One of the important characteristics of seedling growth is the height of the seedling and it was recorded highest in GA<sub>3</sub> 500 ppm (8.93 cm) and lowest in Control (3.76 cm). The maximum seedling length in GA-treated seeds can be due to the fact that it enhances nutrient uptake and increases the process of cell multiplication in internodal tissue causing increased seedling length [16].

**Table 1. Effect of pre-sowing seed treatments on germination parameters of strawberry guava seedlings**

| Treatment                  | Days taken for germination | Germination % | Seedling vigour index - I | Seedling vigour index - II |
|----------------------------|----------------------------|---------------|---------------------------|----------------------------|
| T1:GA <sub>3</sub> 200ppm  | 27.5                       | 59            | 328.66                    | 238.23                     |
| T2: GA <sub>3</sub> 300ppm | 26                         | 65            | 404.76                    | 311.86                     |
| T3: GA <sub>3</sub> 400ppm | 24                         | 70.37         | 508.86                    | 398.16                     |
| T4: GA <sub>3</sub> 500ppm | 20                         | 79.33         | 708.6                     | 584.23                     |
| T5:Thiourea 0.5%           | 24.5                       | 61.34         | 376.5                     | 261.84                     |
| T6: Thiourea 1%            | 22                         | 74.42         | 612.23                    | 529.28                     |
| T7: KNO <sub>3</sub> 0.5%  | 25                         | 69.39         | 480.26                    | 373.08                     |
| T8: KNO <sub>3</sub> : 1%  | 24                         | 73            | 523.16                    | 452.74                     |
| T9: 5% Hcl                 | 28                         | 47.64         | 215.64                    | 168.43                     |
| T10: 10% Hcl               | 29                         | 41.72         | 163.83                    | 131.39                     |
| T11: Control               | 31                         | 34.84         | 129.86                    | 82.46                      |
| Mean                       | 25.5                       | 61.42         | 404.76                    | 321.06                     |
| SE d                       | 1.46                       | 1.93          | 17.92                     | 16.61                      |
| CD(5%)                     | 3.04                       | 4             | 37.18                     | 34.44                      |

**Table 2. Effect of pre-sowing seed treatments on shoot and root parameters of strawberry guava seedlings**

| Treatment                  | Seedling height (cm) | Stem girth (mm) | Total No. of leaves | Leaf area (cm <sup>2</sup> ) | Length of primary root | No. of lateral roots |
|----------------------------|----------------------|-----------------|---------------------|------------------------------|------------------------|----------------------|
| T1:GA <sub>3</sub> 200ppm  | 5.56                 | 0.62            | 5.93                | 1.73                         | 2.14                   | 4.46                 |
| T2: GA <sub>3</sub> 300ppm | 6.23                 | 0.71            | 6.7                 | 2.05                         | 2.52                   | 5.4                  |
| T3: GA <sub>3</sub> 400ppm | 7.23                 | 0.81            | 7.46                | 2.25                         | 2.90                   | 6.43                 |
| T4: GA <sub>3</sub> 500ppm | 8.94                 | 0.99            | 8.43                | 2.6                          | 3.7                    | 7.65                 |
| T5:Thiourea 0.5%           | 6.12                 | 0.66            | 6.06                | 1.92                         | 2.28                   | 5.06                 |
| T6: Thiourea 1%            | 8.20                 | 0.91            | 8.25                | 2.4                          | 3.23                   | 8.33                 |
| T7: KNO <sub>3</sub> 0.5%  | 6.94                 | 0.74            | 7.03                | 2.17                         | 2.70                   | 5.96                 |
| T8: KNO <sub>3</sub> : 1%  | 7.16                 | 0.81            | 8.3                 | 2.35                         | 3.16                   | 6.93                 |
| T9: 5% Hcl                 | 4.56                 | 0.54            | 4.7                 | 1.70                         | 1.85                   | 4.7                  |
| T10: 10% Hcl               | 3.94                 | 0.46            | 3.9                 | 1.58                         | 1.71                   | 4.05                 |
| T11: Control               | 3.76                 | 0.36            | 3.5                 | 1.43                         | 1.36                   | 3.6                  |
| Mean                       | 6.24                 | 0.69            | 6.39                | 2.01                         | 2.50                   | 5.69                 |
| SE d                       | 0.24                 | 0.03            | 0.29                | 0.1                          | 0.06                   | 0.28                 |
| CD(5%)                     | 0.5                  | 0.07            | 0.6                 | 0.2                          | 0.12                   | 0.58                 |

**Table 3. Effect of pre-sowing seed treatments on seedling biomass parameters of strawberry guava seedlings**

| Treatment                  | Fresh weight of seedling(g) | Dry weight of seedling(g) | Fresh weight of shoot(g) | Dry weight of shoot(g) | Fresh weight of root(g) | Dry weight of root(g) |
|----------------------------|-----------------------------|---------------------------|--------------------------|------------------------|-------------------------|-----------------------|
| T1:GA <sub>3</sub> 200ppm  | 10.54                       | 4.03                      | 9.58                     | 3.65                   | 0.96                    | 0.38                  |
| T2: GA <sub>3</sub> 300ppm | 12.62                       | 4.78                      | 11.36                    | 4.24                   | 1.26                    | 0.54                  |
| T3: GA <sub>3</sub> 400ppm | 15.23                       | 5.65                      | 13.6                     | 4.94                   | 1.63                    | 0.71                  |
| T4: GA <sub>3</sub> 500ppm | 17.26                       | 7.36                      | 15.13                    | 6.13                   | 2.13                    | 1.23                  |
| T5:Thiourea 0.5%           | 11.87                       | 4.26                      | 10.77                    | 3.82                   | 1.10                    | 0.44                  |
| T6: Thiourea 1%            | 16.85                       | 7.12                      | 14.84                    | 6.06                   | 2.00                    | 1.06                  |
| T7: KNO <sub>3</sub> 0.5%  | 13.97                       | 5.38                      | 12.54                    | 4.74                   | 1.43                    | 0.63                  |
| T8: KNO <sub>3</sub> : 1%  | 16.05                       | 6.2                       | 14.13                    | 5.18                   | 1.92                    | 1.01                  |
| T9: 5% Hcl                 | 9.20                        | 3.54                      | 8.36                     | 3.21                   | 0.84                    | 0.32                  |
| T10: 10% Hcl               | 8.09                        | 3.15                      | 7.41                     | 2.91                   | 0.68                    | 0.24                  |
| T11: Control               | 6.71                        | 2.39                      | 6.08                     | 2.26                   | 0.63                    | 0.12                  |
| Mean                       | 12.58                       | 4.89                      | 11.25                    | 4.28                   | 1.32                    | 0.61                  |
| SE d                       | 0.4                         | 0.17                      | 0.36                     | 0.15                   | 0.06                    | 0.03                  |
| CD(5%)                     | 0.84                        | 0.35                      | 0.75                     | 0.31                   | 0.12                    | 0.08                  |

The maximum stem girth was observed in seeds treated with GA<sub>3</sub> 500 ppm (0.99 mm) and minimum in control (3.76) (Fig. 2). Higher stem girth can also be attributed to the same fact as in the case of seedling length as observed by Dhankhar and Singh [17] in aonla.

Leaf is the most important photosynthetic site of the plant as the plant derives its energy for its metabolic activities from leaves and its major function is carbon assimilation [18]. Regarding number of leaves and leaf area, seeds treated with GA<sub>3</sub> 500 ppm showed higher number of leaves (8.43) and higher leaf area (2.6 cm<sup>2</sup>) followed by thiourea 1% (8.26) and leaf area (2.4

cm<sup>2</sup>) (Fig. 3). The lowest values were seen in control. The favorable effect of thiourea might be due to the presence of the SH- group which has diverse biological activities such as the transport of photosynthates from source to sink [19]. Similar results were also obtained by [20].

Regarding the length of the primary root and number of lateral roots, seeds treated with GA<sub>3</sub> 500 ppm showed the highest length of root (3.7 cm), number of lateral roots (7.66), and lower values were seen in control i.e., 1.36 cm and 3.6 number of lateral roots. Higher values in GA-treated seeds might be due to the availability of a sufficient amount of gibberellic acid which in turn

gives higher cell elongation and multiplication. These might be reflected in the increase in the length of roots and the higher number of secondary roots. These results are in close conformity with Nurunga et al. (2013) in passion fruit.

### 3.3 Biomass Production of Seedlings

The biomass production of seedlings is presented in (Table 3). Significant differences

were observed between all the treatments for fresh weight and dry weight of root, shoot, and seedling. The highest fresh weight of the shoot was seen in seeds treated with GA3 500 ppm (15.13 g) as compared to the control (6.08 g). Similarly, a higher fresh weight of root was also obtained in 500 ppm GA3 (2.13 g) as compared to the control (0.63) (Fig. 4). As a result of the higher fresh weight of root and shoot, the higher total fresh weight of seedling were also obtained in seeds treated with GA3 500 ppm (17.26 g) as




Fig. 1. Effect of pre sowing treatments on germination (%) of strawberry guava seedling




Fig. 2. Effect of pre sowing treatments on stem grith (mm) of strawberry guava seedling




Fig. 3. Effect of pre sowing treatments on leaf area (cm<sup>2</sup>) of strawberry guava seedling




Fig. 4. Effect of pre sowing treatments on fresh weight of seedling, shoot and root of strawberry guava seedling

compared to control (6.71 g). These results might be due to the production of more photosynthesis and their translocation to the various parts of the plant. Similarly, because of the higher

accumulation of photosynthesis the dry weight of total seedling, shoot, and root was also seen in GA treated as compared to control (Fig. 5).

### 3.4 Seedling Vigour Index

The seedling vigor index is directly proportional to the germination percentage and overall growth of seedlings. Seedling Vigour Index I and Vigour Index II recorded from this investigation are shown in Table 1. Seed treated with GA3 500

ppm was recorded with the highest vigor index I (708.6) and Vigour index II (584.23) as compared to the control (Fig. 6). These can be linked to the fact that in the present study, GA3 500 ppm gave higher germination percentage, seedling length and dry weight of seedling.




Fig. 5. Effect of pre sowing treatments on dry weight of seedling, shoot and root of strawberry guava seedling




Fig. 6. Effect of pre sowing treatments on Vigour index I and Vigour index II of strawberry guava seedling

#### 4. CONCLUSION

Based on the results of the experiment it can be concluded that the strawberry guava seeds treated with GA<sub>3</sub> 500 ppm for 30 minutes gave the maximum germination percentage (79.33%), higher growth parameters (height of the seedling (8.93 cm); the number of leaves (8.43); higher leaf area (2.6 cm<sup>2</sup>); length of root (3.7 cm), number of lateral roots (7.66)) and higher biomass production (fresh weight of shoot (15.13 g)). So to break the dormancy of strawberry guava seed for quick mass propagation of strawberry guava seedling to be used as rootstock for commercial guava, strawberry guava seeds have to be treated with 500ppm GA<sub>3</sub> for 30 minutes.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Available: <https://www.indiastat.com.elibrarytnau.remotexs.in/table/uttar-pradesh-state/guava/selected-state-wise-area-production-productivity-g/1424487>
2. Gomes VM, Souza RM, Mussi-Dias V, Silveira SF, Dolinski C. Guava decline: A complex disease involving *Meloidogyne mayaguensis* and *Fusarium solani*. *Journal of Phytopathology*. 2011 Jan;159(1):45-50.
3. Carneiro RM, De Freitas VM, Mattos JK, Castro JM, Gomes CB, Carneiro RG. Major guava nematodes and control prospects using resistance on *Psidium spp.* and non-host crops. In III International Symposium on Guava and other Myrtaceae. 2012 Apr 23;959:41-49.
4. Miranda GB, Souza RM, Gomes VM, Ferreira TD, Almeida AM. Avaliação de acessos de *Psidium spp.* quanto à resistência a *Meloidogyne enterolobii*. *Bragantia*. 2012;71:52-8.
5. Almeida EJ, Santos JM, Martins AB. Resistência de goiabeiras e araçazeiros a *Meloidogyne mayaguensis*. *Pesquisa Agropecuária Brasileira*. 2009;44:421-3.
6. Teotia SS, Phogat KP. Effect of rootstocks on growth, yield and quality in guava (*Psidium guajava*). *Progressive Horticulture*. 1971;2(4):37-45.
7. Vishwakarma PK, Vasugi C, Umamaheswari R, Sriram S, Nandeesh P, Sankaran M, Shivashankara KS. Screening of *Psidium* species and their interspecific hybrid progenies for resistance to *Fusarium oxysporum* f. sp. *psidii* and southern root-knot nematode. *South African Journal of Botany*. 2023 Apr 1;155:249-60.
8. de Oliveira Costa SN, e Silva MV, Ribeiro JM, e Castro JM, Muzitano MF, da Costa RG, Oliveira AE, Fernandes KV. Secondary metabolites related to the resistance of *Psidium spp.* against the nematode *Meloidogyne enterolobii*. *Heliyon*. 2023 Jul 1;9(7).
9. Uowolo AL, Denslow JS. Characteristics of the *Psidium cattleianum* (Myrtaceae) Seed Bank in Hawaiian Lowland Wet Forests. *Pacific Science*. 2008 Jan;62(1):129-35.
10. Silva AD, Perez SC, Paula RC. Qualidade fisiológica de sementes de *Psidium cattleianum* Sabine acondicionadas e armazenadas em diferentes condições. *Revista Brasileira de Sementes*. 2011;33: 197-206.
11. Nava AD, Hernández VA, Jaimes MN, Castro EH, Álvarez DV, Villaseñor GD, Alberto FP, Lagunas BC. Growth Kinetics of Vegetative and Reproductive Organs of Guava (*Psidium guajava* L.) in Iguala Guerrero, Mexico this work is licensed under the Creative Commons Attribution International License (CC BY).
12. Abdul-Baki AA, Anderson JD. Vigor determination in soybean seed by multiple criteria 1. *Crop Science*. 1973 Nov;13(6): 630-3.
13. Pawshe YH, Patil BN, Patil LP. Effect of pregermination seed treatments on the germination and vigour of seedlings in custard apple (*Annona squamosa* L.). *Annals of Plant Physiology*. 1997;11:150-4.
14. Gurung N, Swamy GS, Sarkar SK, Ubale NB. Effect of chemicals and growth regulators on germination, vigour and growth of passion fruit (*Passiflora edulis* Sims.). *The Bioscan*. 2014;9(1):155-7.
15. Hore JK, Sen SK. Role of presowing seed treatment on germination, seedling growth and longevity of ber (*Zizyphus mauritiana* Lam) seeds. *Indian Journal of Agricultural Research*. 1994;28(4):285-9.
16. Barathkumar TR. Studies on influence of different seed treatments on dormancy breaking in aonla (*Phyllanthus embolic* L). *Journal of Pharmacognosy and Phytochemistry*. 2019;8(2S):131-3.
17. Dhankhar DS, Singh M. Seed germination and seedling growth in aonla (*Phyllanthus*

- emblica* Linn.) as influenced by gibberellic acid and thiourea. CROP RESEARCH-HISAR-. 1996; 12:363-6.
18. Parab AM, Mathad JC, Malshe KV. Effect of Presoaking Chemicals on Germination and Subsequent Seedling Growth of Papaya (*Carica papaya*) Cv. Solo. IJCS. 2017;5(4):1812-6.
  19. Meena SK, Jat NL, Sharma B, Meena VS. Effect of plant growth regulators and sulphur on productivity of coriander (*Coriandrum sativum* L.) in Rajasthan. J. Environ. Sci. Int. 2014;6:69-73.
  20. Balai LR, Keshwa GL. Effect of thiourea on yield and nutrient uptake of coriander (*Coriandrum sativum* L.) varieties under normal and late sown conditions. Journal of Spices and Aromatic Crops. 2012 Apr 5;20(1).

---

© 2023 Rao et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/106392>