



# Impact of Different Formulations of Biofertilizer Consortia (NPK & NPKZn) on Proximate Nutritional Composition of Hybrid Maize (Co H (M) 8) Grains

M. Akash <sup>a</sup>, M. Gnanachitra <sup>a\*</sup>, D. Balachandar <sup>a</sup>  
and P. Geetha <sup>b</sup>

<sup>a</sup> Department of Agricultural Microbiology, Tamil Nadu Agricultural University, Coimbatore-3, India.

<sup>b</sup> Centre for Post-Harvest Technology, Tamil Nadu Agricultural University, Coimbatore-3, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i234257

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110899>

Original Research Article

Received: 07/10/2023

Accepted: 19/12/2023

Published: 21/12/2023

## ABSTRACT

The study focused on the effect of liquid and powder formulations of biofertilizer consortia developed for NPK and NPKZn nutrition using *Azospirillum*, phosphobacteria, potash bacteria and zinc solubilizing bacteria on the proximate nutrient component analysis of maize grains and to evaluate their potential as nutrient-rich sources for both human and animal consumption. Cereals, particularly maize, is one of the globally utilized and significant crops, contributing substantially to human diet, as animal feed, substrate for fuel and occupying a considerable portion of agricultural land. The nutritional composition of maize, encompassing starch, carbohydrate, protein, crude fiber and ash plays an important role in dietary planning and epidemiological research. The experiment involves different treatments, each representing a specific biofertilizer consortium application

\*Corresponding author: E-mail: [mgc74@tnau.ac.in](mailto:mgc74@tnau.ac.in);

method, such as liquid and powder formulations. The liquid formulation of NPK and NPKZn biofertilizer consortia applied through seed treatment and powder formulation of NPK and NPKZn biofertilizer consortia applied through seed coating and compared with the existing recommendation in maize crop. The results indicate variations in moisture content, protein levels, starch yield, carbohydrate composition, ash content, and crude fiber among the treatments. Notably, seed coating with NPKZn biofertilizer consortia in powder form, stands out with promising attributes, including high protein content (8.08%), significant starch yield (69.88%), and balanced amylose to amylopectin ratio. The findings contribute valuable insights for selecting optimal biofertilizer consortia formulations to enhance the nutritional quality of maize, crucial for both agricultural and dietary considerations.

**Keywords:** Maize; NPK and NPKZn biofertilizer consortia; proximate nutritional composition.

## 1. INTRODUCTION

Cereals stand out as the most commonly cultivated and consumed crops globally, as they play a crucial role in supplying a significant portion of the calories in human diets and account for over half of the world's agricultural land area (approximately 45% and 54%, respectively) on a global scale [1,2]. Among cereals, maize (*Zea mays* L.), recognized as a highly adaptable and continually evolving cereal crop, thrives in diverse agroclimatic conditions and holds a significant position within the Poaceae family, serving as a dietary staple in numerous countries worldwide [3]. Indigenous farmers are credited with a remarkable achievement in plant breeding the domestication and diversification of maize. Archaeological evidence traces the initial cultivation of maize to Mexico and Central America [4]. Renowned as the "queen of cereals" globally, maize stands out due to its unparalleled genetic production potential. Besides being a crucial staple, maize finds extensive industrial use, particularly in the production of high fructose corn syrup [3]. Maize grain, constituting a significant portion of broiler feed (60%), is utilized for non-ruminants in its whole form [5]. Notably, its high starch and low fibre content contribute to a superior conversion ratio to meat and eggs compared to other grains. In tropical regions, there's a preference for yellow maize in animal feed. Maize holds pivotal importance as a raw material in the starch, feed, and food industries, providing enhanced health benefits when incorporated into food processing and feeding [6].

Additionally, maize nutritional composition plays a vital role in nutritional planning and supports epidemiological research [7]. Maintaining a safe

moisture level of 12% to 14% when wet is recommended for cereal grains, including maize [8]. Maize seeds encompass moisture (11.6-20.0%), ash (1.10-2.95%), protein (4.50-9.87%), fat (2.17-4.43%), fibre (2.10-26.70%), and carbohydrates (44.60-69.60%) [1,6]. This examination holds particular significance in the food sector to guarantee steadfast quality and safety of products for consumers. Through the measurement of macronutrient levels, scientists can evaluate factors such as energy content, amino acid composition, and fiber presence, offering valuable information about the nutritional value of plant-derived foods [9]. The versatility of maize is evident in its various products, such as whole corn, corn flour, corn starch, corn gluten, corn syrup, tortillas, tortilla chips, polenta cornmeal, corn oil, popcorn, and cornflakes. Maize processing is carried out by three primary industries: wet millers for starch, sweeteners, and maize oil; dry millers for maize flour and grits; and distillers for beverage and bioethanol production. The rising demand for maize as a carbohydrate source in the processing sector has led to the expansion of breeding programs focusing on hybrids with specialized features for distinct uses [10]. Maize encompasses a variety of micronutrients essential for maintaining optimal health. Common vitamin-related disorders such as beriberi, pellagra, anemia, and more can be addressed through the consumption of maize, given its rich content of vitamin B-complex, including thiamin (B1), niacin (B3), pantothenic acid (B5), pyridoxine (B6), and folate (B9). The enhancement of vitamins in plant-derived foods is achievable through the improvement of growth conditions, traditional plant breeding methods, or the application of transgenic techniques—a process known as biofortification [11]. Regular consumption of maize has been associated with facilitating the

removal of harmful dietary components and expediting food transit through the gut. It also plays a protective role in the digestive system, reducing stomach acidity, and enhancing gallbladder function [12,13]. Despite limited research on comparing the nutritional content of different biofertilizer methods used on maize seeds in farmers' fields, this study aims to explore the proximate composition of various biofertilizer consortium techniques applied to maize seedlings and their quality. Understanding these variations will aid in selecting optimal biofertilizer consortiums (NPK and NPKZn) in liquid or powder formulations for both human and animal consumption.

## 2. MATERIALS AND METHODS

### 2.1 Plant Materials

Maize seeds COH(M)8 cultivar, sourced from the Department of Millets at the Center for Plant Breeding and Genetics, Tamil Nadu Agricultural University in Coimbatore, were utilized for the study.

### 2.2 Development of Biofertilizer Consortia for Maize

Two distinct biofertilizer consortia (NPK, NPKZn) were developed in two different formulations (liquid and powder) with the aim of enhancing the efficiency of applied fertilizers and augmenting the availability of applied essential nutrients (nitrogen, phosphorus, potassium, and zinc) to the soil in the form of urea, single super phosphate, muriate of potash and zinc sulphate for maize cultivation. The liquid formulations of biofertilizer consortia were developed using the standard biofertilizers viz., *Azospirillum brasilense* (Sp7), *Bacillus megaterium* (Pb1), *Paenibacillus mucilaginosus* (KRB9), and *Pseudomonas chlororaphis* (ZSB15) normally recommended for maize cultivation with respect to NPK and NPKZn nutrition. Using these liquid formulations of biofertilizer consortia, powder formulation was developed with carrier. The liquid formulation of NPK &NPKZn consortia was used for seed treatment of maize; whereas the powder formulation is used for seed coating of maize by seed coating mixture [14]. These biofertilizer consortia (NPK &NPKZn) coated maize seed is a kind of ready to use input. A field trial was taken in maize with five different

treatments. The liquid biofertilizer consortia formulation was used for seed treatment of maize (as per the recommended dosage); whereas powder formulation of consortia used for seed coating of maize (the powder formulation of the consortia was used for development of coated maize seed as a ready to use input (@ 5 g/ 100 g ). These two consortia were compared with the existing recommended practice and farmer's practice. This study is mainly to investigate and compare the effect of two different formulations of NPK and NPKZn consortia on enhancing the nutritional parameters of maize grain. Because, in general biofertilizers enhance the quality of the farm produce.

### 2.3 Sample Preparation for Proximate Analysis

After harvesting the maize cobs, the grains were removed and used for the entire analysis. Initially, the maize grains were dried up to 12% moisture content and the dried maize grain samples further powdered and passed through a 60-mesh sieve. The powdered samples were packed in a high-density polyethylene (HDPE) cover stored at room temperature and used for the entire study.

#### 2.3.1 Estimation of moisture content

Moisture Content was determined by the method described previously [15,16]. Moisture is a critical component of food. The moisture content of any food is determined not only to assess the chemical makeup of the food item on a moisture-free basis, but also to determine the shelf life of the product. A dry and weighted petri plate held a 10 g sample. At regular intervals, the petri dish containing the material was weighed until a steady weight was attained. The percentage of moisture was calculated using the following formula:

$$\text{Moisture}(\%) = \frac{W_1 - W_2}{W} \times 100$$

where,

$W_1$  = weight of sample with crucible (g),  
 $W_2$  = weight of dried sample with crucible (g),  
and  
 $W$  = weight of wet sample (g).

### 2.3.2 Estimation of carbohydrate

The carbohydrate was estimated by anthrone method [17]. In a boiling tube, 100 mg of maize grain sample was taken and added with 5 ml of 2.5 N HCl. This mixture was then hydrolyzed in a boiling water bath with for 3 hours and cooled to room temperature. In order to remove the effervescence, the hydrolyzed mixture was then neutralized with solid sodium carbonate and made up the volume to 100ml for centrifugation. After centrifugation, from the supernatant 0.5 and 1mL of sample aliquots were taken for assessment and compared with the standards taken at 0 (Blank), 0.2, 0.4, 0.6, 0.8, and 1ml. The sample reagents taken at different volume were made up to 1 ml with fresh water and added with 4mL of anthrone reagent. This reaction mixture was then kept in a boiling water bath for eight minutes and cooled quickly. The resultant colour developed after reaction from green to dark green was read at 630nm and compared the absorbance against the standard concentration (glucose) taken on X axis. From the standard graph, the quantity of carbohydrate was arrived using the following formula.

$$\text{carbohydrate (g/100g)} = \frac{\text{mg of glucose}}{\text{Volume of test sample}} \times 100$$

### 2.3.3 Estimation of starch

To estimate starch, first the sugars in the maize grain sample has to be eliminated [17]. To eliminate the sugars, 0.5 g of maize grain sample was taken, homogenized in 80% ethanol and the residue was obtained by centrifugation. The residue was rinsed with 80% ethanol until it becomes colourless with addition of anthrone reagent. Once the residue becomes colorless, then it was carefully dried in a water bath. To the dried residual substance, 5.0 mL of distilled water and 6.5 mL of 52% perchloric acid was added and performed the extraction at 0°C for 20 minutes. Now the sample contents were centrifuged, and repeated the extraction with fresh perchloric acid. After extraction, all the supernatants were pooled to yield a total volume of 100 ml. From the supernatant 0.2 ml was taken and diluted to 1 ml with water. Simultaneously, standards were also taken at 0.2, 0.4, 0.6, 0.8, and 1 mL of water in each tube

and added with 4 mL of anthrone reagent for reaction and kept in a boiling water bath for eight minutes. After the reaction, the sample was cooled rapidly and measured the intensity of color changed from green to dark green at 630 nm and the amount of glucose was calculated as per the following equation,

$$\text{Starch (\%)} = \text{glucose (\%)} \times 0.9$$

### 2.3.4 Estimation of amylose

The powdered sample of 100 mg was taken and added with 1 ml of distilled ethanol added, followed by 10 ml of 1 N NaOH, and allowed it to stand overnight. The extract volume was then raised to 100 ml. From that, 2.5 ml of the extract was taken and to that 20 ml of distilled water, and few drops of phenolphthalein were added. To this reaction mixture, 0.1 N HCl was added drop by drop until the pink color completely disappears and the volume was adjusted to 50 ml with 1 ml of iodine reagent, and read at 590 nm. Standard amylase solutions of 0.2 ml, 0.4 ml, 0.6 ml, 0.8 ml, and 1 ml were also taken, and allowed till it develop the color as in the example. Using a standard graph, the quantity of amylose in the sample was calculated. For a blank, 1ml of iodine reagent diluted in 50ml of distilled water was used.

Absorbance corresponds to 2.5ml of test solution = 'x' mg amylose

100 ml contains = ('x' ÷ 2.5) × 100mg amylose = % amylose

Subtracting the amylose content from the starch content provides a quantity of amylopectin.

### 2.3.5 Estimation of protein content

The AOAC standard technique was used to determine the crude protein content [18]. Protein content was calculated using three stages: digestion, distillation, and titration.

i) N<sub>2</sub> content calculation:

$$\% \text{ of N}_2 = \text{Burette reading} \times \text{Normality of H}_2\text{SO}_4 \times \text{mL equivalent of N}_2$$

In this case, the Normality of H<sub>2</sub>SO<sub>4</sub> is 0.2 and the ml equivalent of N<sub>2</sub> is 1.4.

ii) Protein content calculation:

$$\% \text{ Protein} = \% \text{ of N}_2 \times \text{Protein factor}$$

In this case, Protein factor = 6.25.

### 2.3.6 Estimation of crude fibre

The crude fibre content was determined using the wende approach through the acid-alkali digestion technique [19]. Fibre, consisting of insoluble vegetable matter such as cellulose, hemi-cellulose and lignin, is resistant to digestion by proteolytic or diastatic enzymes and can only be utilized through microbial fermentation. In this method, three grams of a moisture and fat-free maize grain samples from different treatments were placed in a 500 ml beaker, boiled for 30 minutes with 200 ml of 1.25 percent sulphuric acid sodium hydroxide solution. The samples were boiled, then filtered through muslin cloth. The residue was then rinsed with hot distilled water until the alkali was no more visible, and then 50 ml of alcohol and ether were used to wash it. The residue was then placed in the crucible ( $W_1$ ) and dried in an oven at 130°C for 2 - 3 hours before being cooled and weighed ( $W_2$ ). In a muffle furnace at 600 °C for 2-3 hours, a crucible containing dry residue was cooled, and weighed again ( $W_3$ ). Crude fibre was calculated using the method below;

$$\text{Crude fibre}(\%) = \frac{(W_2 - W_1) - (W_3 - W_1)}{\text{Weight of sample}} \times 100$$

Where,

- $W_1$ = Weight of empty crucible
- $W_2$ = Weight of crucible with dry residue
- $W_3$ = Weight of crucible with heated residue

### 2.3.7 Estimation of ash content

The ash content of a maize sample was determined using the AOAC standard procedure [18]. Five gram of powdered maize sample was taken in a pre-weighed crucible and placed in a muffle furnace at 550°C to 600°C for 5-6 hours. After complete ashing, the crucible was cooled and dried in a desiccator before being weighed. From the weights recorded the present ash content was calculated by the following equation.

$$\text{Ash}(\%) = \frac{W_1 - W_2}{W} \times 100$$

Where,

- $W_1$  = weight of ash with crucible (g)
- $W_2$  = weight of empty crucible (g)
- $W$  = weight of sample (g).

## 2.4 Statistical Analysis

A statistical analysis was conducted to determine the nutritional composition of maize in a Completely Randomized Design (CRD) with five replications and analyzed using statistical software (SPSS windows version 20). ANOVA was performed on triplicate values. Mean comparisons were performed using Duncan's multiple range tests for significant effect at least of  $p < 0.05$ .

## 3. RESULTS AND DISCUSSION

The proximate analysis of maize grains evaluated under five different treatments ( $T_1$  to  $T_5$ ), revealed variations in key nutritional parameters.

### 3.1 Moisture Content

The moisture content in food holds significant economic importance for both producers and consumers, as it exhibits an inverse relationship with the dry matter present. This factor is also pivotal for ensuring the stability and overall quality of food products. The results from the present study revealed that the moisture content recorded ranges from 3.6 % to 4.0 % (Table 1; Fig. 1). The significantly maximum moisture content of 4.0 % was observed in maize seed treated with NPK liquid biofertilizer consortia ( $T_2$ ). In contrast to Oluwalana's findings, who recorded a moisture content ranging from 9.85% to 11.35% in maize, the current study observed a decreased level of moisture content irrespective of all the treatments (3.6 to 4.0 %) [20]. The moisture content measured at 4.0% was lower than the reported values of 8.0% by Hamisu et al. [21], 6.00% by Atiku et al. [22], 11.74% by Danish et al. [23], and 4.6% by Danje [24]. The quantity of moisture present in a food item commonly influences its weight, texture, taste, and durability on the shelf. Even a slight deviation from the specified standard can have

adverse effects on the physical characteristics of a food substance. For example, excessively dry grains could compromise the consistency of the end product. Additionally, higher levels of total water content can accelerate microbial growth, potentially leading to spoilage and necessitating disposal of the affected grains [25]. This itself showed a positive sign about the quality of maize grain to be used as food or feed. Because, higher moisture levels tend to decrease the shelf life of the maize; and the observed lower moisture content -a favourable characteristic for prolonged storage. Moreover, grains with elevated moisture levels are susceptible to quick deterioration due to issues like mold growth and insect damage, underscoring the need for careful consideration of moisture content in food processing and consumption [26,27].

### 3.2 Carbohydrate

Maize is recognized for its high carbohydrate content, providing energy, facilitating the utilization of body fats through metabolic processes, and supporting the proper functioning of the intestinal tract. The analysed maize grain samples showed carbohydrate content ranged from 67.76 to 74.40 %. This is in accordance with the results previously reported [28 29]. The study also revealed a significant difference in carbohydrate content among the five treatments (Table 1; Fig 2), with carbohydrate levels ranging from 69.25% to 71.13%. The maximum carbohydrate content of 71.13% was noted in the maize seed coated with NPKZn (T<sub>5</sub>) biofertilizer consortia, compared to the carbohydrates content in maize seed coated with NPK biofertilizer consortia (T<sub>4</sub>), maize seed treated with NPKZn liquid biofertilizer consortia (T<sub>3</sub>), and uninoculated (T<sub>1</sub>) recorded 70.89%,70.89% and 69.28%, respectively. In accordance with the present study findings, Ullah and his co-workers also recorded a carbohydrate percentage ranging from 69.659% to 74.549% in maize [6]. This aligns with existing literature, as Wilson (1987) indicated that the primary chemical component of maize grain is carbohydrates, constituting approximately 72-73% of the grain [30]. Ekanem and his co-workers [25] documented findings ranged between 67.07% to

70.70%, exhibiting a close resemblance to the carbohydrate content values observed in the present study. Similarly, yellow corn exhibited the highest total carbohydrate content at 76.57%, while White corn recorded 76.00%. Additionally, sweet corn and popcorn displayed carbohydrate contents of 69.25% and 74.12%, respectively [31].

### 3.3 Estimation of Starch


Starch is the main carbohydrate reserve in plants and an important part of our nutrition. Starch undergoes a variety of physical, enzymatic, or chemical modifications to yield different products. Among these products are maltodextrin, dextrin, dextrose monohydrate, sorbitol, liquid glucose, high maltose syrup, dextrose syrup, and anhydrous dextrose. These derivatives find applications in the production of beverages, bakery items, pastries, meat, soups, sauces, baby food, textiles, dextrans, paper, and pharmaceutical products [32,33]. starch (60-70%) being the main product, the by-products include steep liquor/solubles, maize germ, maize fibre and maize gluten [33]. The mean starch yield of the maize treatments ranged from 67.24% to 69.88% (Table 1; Fig 2). The starch obtained from maize seed coated with NPKZn biofertilizer consortia (T<sub>5</sub>) was significantly ( $P < 0.05$ ) higher (69.88 %) than maize seed coated with NPK biofertilizer consortia (T<sub>4</sub>) recorded 68.36 which is on par with the other two liquid formulation of biofertilizer consortia treatments (T<sub>3</sub> & T<sub>2</sub>). The starch yield is less than the values obtained for potato and corn starch, which were 93.4% and 96.5% respectively [34]. Deepak and Jayadeep, also reported in their results the starch content ranged from 67.0% to 73.0%, demonstrating close consistency with the starch content values observed in the present study [33]. While the wide range of starch content among the cultivars is interesting, further information regarding specific starch components should be more helpful in formulating an effective strategy for their further utilization. The amylose percentage ranges from 20.22% to 21.98%, while amylopectin percentage varies from 46.47% to 47.90%, indicating different starch structures.

**Table 1. Effect of biofertilizer consortia (NPK and NPKZn) on the proximate nutritional composition of maize grain**

S. No.	Treatments	Moisture (%)	Carbohydrate content (%)	Starch content (%)	Amylose (%)	Amylopectin (%)	Crude protein (%)	Crude fibre (%)	Ash content (%)
1	T <sub>1</sub>	3.70 (±0.05) <sup>b</sup>	69.28 (±0.31) <sup>a</sup>	67.76 (±0.58) <sup>b</sup>	20.22(±0.04) <sup>b</sup>	47.54 (±0.08) <sup>a</sup>	7.51 (±0.04) <sup>b</sup>	1.82 (±0.03) <sup>b</sup>	1.98 (±0.03) <sup>c</sup>
2	T <sub>2</sub>	4.00 (±0.05) <sup>a</sup>	69.25 (±0.92) <sup>a</sup>	67.24 (±0.09) <sup>b</sup>	20.76 ±0.23) <sup>b</sup>	46.47(±0.73) <sup>a</sup>	7.70 (±0.03) <sup>b</sup>	1.87(±0.01) <sup>ab</sup>	2.03 (±0.03) <sup>bc</sup>
3	T <sub>3</sub>	3.80 (±0.06) <sup>b</sup>	70.89 (±0.29) <sup>a</sup>	68.22 (±0.12) <sup>b</sup>	20.45(±0.05) <sup>b</sup>	47.77 (±0.75) <sup>a</sup>	8.02 (±0.02) <sup>a</sup>	1.90 (±0.00) <sup>a</sup>	2.11 (±0.02) <sup>ab</sup>
4	T <sub>4</sub>	3.72 (±0.01) <sup>b</sup>	70.89 (±0.36) <sup>a</sup>	68.36 (±0.30) <sup>b</sup>	21.84 ±0.27) <sup>a</sup>	46.53 (±0.06) <sup>a</sup>	8.03 (±0.11) <sup>a</sup>	1.92 (±0.01) <sup>a</sup>	2.07(±0.01) <sup>abc</sup>
5	T <sub>5</sub>	3.60 (±0.05) <sup>b</sup>	71.13(±0.02) <sup>a</sup>	69.88 (±0.40) <sup>a</sup>	21.98 ±0.12) <sup>a</sup>	47.90 (±0.13) <sup>a</sup>	8.08 (±0.07) <sup>a</sup>	1.94 (±0.02) <sup>a</sup>	2.15 (±0.03) <sup>a</sup>
	Mean	3.76	70.29	68.29	21.05	47.24	7.87	1.89	2.07


Mean with same alphabet in the same column are not significantly different from each other at  $p < 0.05$

T<sub>1</sub> –Uninoculated maize seeds (control), T<sub>2</sub>–Seed treatment with liquid formulation of NPK biofertilizer consortium, T<sub>3</sub>–Seed treatment with liquid formulation of NPKZn biofertilizer consortium, T<sub>4</sub>–Seed coating with powder formulation of NPK biofertilizer consortium, T<sub>5</sub>– Seed coating with powder formulation of NPKZn biofertilizer consortium.



**Fig. 1. Effect of biofertilizer consortia (NPK and NPKZn) on the moisture, crude protein and crude fibre content of maize grain**

*T<sub>1</sub>–Uninoculated maize seeds (control), T<sub>2</sub>–Seed treatment with liquid formulation of NPK biofertilizer consortium, T<sub>3</sub>–Seed treatment with liquid formulation of NPKZn biofertilizer consortium, T<sub>4</sub>–Seed coating with powder formulation of NPK biofertilizer consortium, T<sub>5</sub>– Seed coating with powder formulation of NPKZn biofertilizer consortium*



**Fig. 2. Effect of biofertilizer consortia (NPK and NPKZn) on the carbohydrate, starch, amylose and amylopectin content of maize grain**

*T<sub>1</sub>–Uninoculated maize seeds (control), T<sub>2</sub>–Seed treatment with liquid formulation of NPK biofertilizer consortium, T<sub>3</sub>–Seed treatment with liquid formulation of NPKZn biofertilizer consortium, T<sub>4</sub>–Seed coating with powder formulation of NPK biofertilizer consortium, T<sub>5</sub>– Seed coating with powder formulation of NPKZn biofertilizer consortium*

### 3.4 Estimation of Protein Content

Protein is the most important biochemical character; since it contributes to the synthesis of nitrogen-containing compounds, including antibodies and enzymes, which are essential for

normal physiological functions. In the present study, compared to uninoculated control all the biofertilizer consortia formulation treatments recorded an increase in crude protein content (7.70 to 8.08%) (Table 1; Fig 1). The significantly maximum value of protein content 8.08% was



observed in maize seeds coated with NPKZn biofertilizer consortia (T<sub>5</sub>) maize grains, followed by seed coating with NPK biofertilizer consortia (T<sub>4</sub>) recorded high protein. Uninoculated maize seeds (T<sub>1</sub>) registered a minimum protein content of 7.51%. Similar to this study, the effect of *Azospirillum* and *Azotobacter* was found to increase the protein content in maize as 6.34 & 7.00 % respectively [35]. Various studies have documented protein levels in maize hybrids ranging from 7.77% to 13.84% [36] and 8.40% to 11.84% [25]. As per FAO (1983) enhancing the nutritional value of maize protein is possible through improved maize breeding, storage, cooking practices, and fortification methods [6,30]. Maize protein, typically ranging from 6.0 to 12 percent, is deemed subpar due to its low lysine and tryptophan levels. Nonetheless, a considerable number of the assessed maize cultivars exhibited elevated protein content, rendering them suitable for applications in baby food formulations and other dietary requirements.

### 3.5 Estimation of Crude Fibre

Crude fibre, primarily composed of cellulose and hemicellulose, offers favourable effects in humans by enhancing water retention capacity as food moves through the digestive tract. A diet abundant in crude fibre is considered healthful [37]. as it contributes to the production of larger and softer faeces. It may be beneficial for conditions like diabetes and high blood cholesterol [38]. The results of crude fibre analysis in various treatments ranged from 1.82% to 1.94% (Table 1; Fig 1). The maximum crude fibre content, 1.94%, was observed in maize seeds coated with NPKZn biofertilizer consortia (T<sub>5</sub>), followed by maize seeds coated with NPK biofertilizer consortia (T<sub>4</sub>) at 1.92%, while the minimum crude fibre content, 1.82%, was noted in uninoculated maize seeds (T<sub>1</sub>). The crude fibre percent within the range of 0.80% to 2.32% was reported closely aligning with the findings of the present study [6]. Ekanem and co-workers also reported almost the same range of 1.62% to 4.36%, demonstrating close consistency with the crude fibre content values observed in the present study [25].

### 3.6 Estimation of Ash Content

The analysis of foodstuffs for ash content percentage serves to identify the non-organic

matter component within the dry matter, representing the residue after the organic matter undergoes processes like oven drying, ignition, or complete oxidation. This ash content provides a rough estimate of the overall mineral content present in the food. In the case of maize seeds under different treatments, the percentage ash content ranged from 1.98% to 2.15% (Table 1). Comparable findings of 0.70% to 2.50% in various maize hybrids were previously reported [39-42]. Maziya-Dixon and co-workers also reported results within the range of 1.4% to 3.3%, demonstrating close consistency with the ash content values observed in the present investigation [43]. which is in agree with the results obtained by Anyaegbu et al. [44] with a range of 1.25% to 1.52%. Ekanem and co-workers [25] documented ash content of 1.21% to 1.46%, exhibiting a close resemblance to the ash content values observed in the present study. Whereas, it (2.15%) was observed to be lower than the reported values by Hamisu et al. [21] as 4.0%.

## 4. CONCLUSION

This research investigated the impact of liquid and powder formulation of NPK and NPKZn biofertilizer consortia, on the proximate nutritional composition of maize grain under different treatments. In general, bio-organic fertilization improves soluble sugars, starch, carbohydrates, protein, and amino acid contents in maize seeds. In the present study also revealed notable variations in moisture content, carbohydrate composition, starch content, amylose and amylopectin ratios, crude protein, crude fibre, and ash content among the treatments. Maize seeds coated with NPKZn biofertilizer consortia exhibited superior nutritional characteristics, including higher protein content and starch yield. The study also emphasized the importance of moisture content, as lower levels contribute to prolonged shelf life. The results contribute valuable insights into the nutritional enhancement of maize through biofertilizer application, offering potential benefits for both human and animal consumption. The study provides a comprehensive understanding of the proximate nutritional changes induced by different biofertilizer treatments, offering practical guidance for optimizing maize cultivation strategies. Additionally, the study underscores the versatility of maize in various industrial applications, from food processing to bioethanol

production, further highlighting its significance in global agriculture and nutrition.

## ACKNOWLEDGEMENT

Indian Council of Agricultural Research - All India Network Project on Soil Biodiversity and Biofertilizers (ICAR – AINP – SBB) and Department of Agricultural Microbiology, Tamil Nadu Agricultural University is acknowledged for the technical and financial support.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Enyisi IS, Umoh V, Whong C, Alabi O, Abdullahi I. Chemical and nutritional values of maize and maize products obtained from selected markets in Kaduna. *Journal of Pharmaceutical and Allied Sciences*. 2014;11(2):2106-2113.
2. FAOSTAT, FAOSTAT data - food balance. 2017: Rome, Italy.
3. Rouf Shah T, Prasad K, Kumar P. Maize— A potential source of human nutrition and health: A review. *Cogent Food & Agriculture*. 2016;2(1):1166995.
4. Singh M, Kumar S, Broadening the genetic base of grain cereals, Springer; 2016.
5. Dahlke F, Ribeiro A, Kessler A, Lima A. Corn particle size and physical form of the diet and their effects on performance and carcass yield of broilers. *Brazilian Journal of Poultry Science*. 2001;3:241-248.
6. Ullah I, Ali M, Farooqi A. Chemical nutritional properties of some maize (*Zea mays* L.) varieties grown in NWFP, Pakistan. *Pakistan Journal of Nutrition*. 2010;9(11):1113-1117.
7. Ali M, Ali W, Ahmed S, Ullah I. Mineral composition, quality and physico-chemical parameters of the local tallow of Pakistan. *Pakistan Journal of Nutrition*. 2008;7(5): 717-720.
8. Bala BK. Drying and storage of cereal grains. John Wiley & Sons; 2016.
9. Rizzo NS, Jaceldo-Siegl K, Sabate J, Fraser GE, JotAoN, and Dietetics, Nutrient profiles of vegetarian and nonvegetarian dietary patterns. 2013;113(12):1610-1619.
10. Pajić Z, Radosavljević M, Filipović M, Todorović G, Srdić J, Pavlov M. Breeding of speciality maize for industrial purposes. *Genetika*. 2010;42(1):57-66.
11. Muhammad II, Ullah SK, Kumar N, Bala HA, Mathur AJJoBS. Efficiency of vitamin B6 (Pyridoxine HCl) on production of Endogenous IAA for growth promotion in *Zea mays* varieties (Azam & Jalal). 2021; 6(4):486-492.
12. Elsgaard L, Børgesen C, Olesen J, Siebert S, Ewert F, Peltonen-Sainio P, Rötter R, Skjelvåg A. Shifts in comparative advantages for maize, oat and wheat cropping under climate change in Europe. *Food additives & Contaminants: Part A*. 2012;29(10):1514-1526.
13. Li L, Sun J, Zhang F, Li X, Yang S, Rengel Z, Wheat/maize or wheat/soybean strip intercropping: I. Yield advantage and interspecific interactions on nutrients. *Field Crops Research*. 2001;71(2):123-137.
14. Priyadharshini C, Gnanachitra M, Balachandar D, Jayanthi D. Assessment of shelf-life and efficacy of the seed-coating delivery system of biofertilizers in maize. *International Journal of Plant and Soil Science*. 2022;34(22):548-558.
15. Lee H, Htoon A, Paterson J. Alkaline extraction of starch from Australian lentil cultivars Matilda and Digger optimised for starch yield and starch and protein quality. *Food Chemistry*. 2007;102(3):551-559.
16. Ndukwe OK, Edeoga H, and Omosun G, Varietal differences in some nutritional composition of ten maize (*Zea mays* L.) varieties grown in Nigeria. *International Journal of Academic Research Reflection*. 2015;3(5):1-11.
17. Hodge J, Hofreiter B. Determination of reducing sugars and carbohydrates, in *Methods in carbohydrate chemistry*, R. Whistler and M. WOLFROM, Editors. Academic Press Inc., New York. 1962; 380.
18. Chemists A, Association of Official Analytical Chemists (AOAC) Official methods of analysis. AOAC; 2004.
19. James CS, Analytical chemistry of foods. Springer Science & Business Media; 2013.
20. Oluwalana IB, Comparative effects of sprouting on proximate, mineral composition and functional properties of white and yellow sweet maize (*Zea mays* var *saccharata*). *Journal of Emerging*

- Trends in Engineering Applied Sciences. 2014;5(7):111-115.
21. Hamisu A, Muhammad II, Ullah SK, Hani DG, Comparative study on the nutritional value of local maize variety (Hakorin Hajiya) and improved varieties of maize (sammaz14 and golden strawberry).
  22. Atiku Y, Abdulsalam S, Mohammed J, Ahmed SJJAS, Management E, Physicochemical and Microbial Characterization of Maize Cob for Bioethanol Production. 2023;27(6):1193-1201.
  23. Danish M, Naqvi M, Farooq U, Naqvi SJEP. Characterization of South Asian agricultural residues for potential utilization in future 'energy mix'. 2015;75:2974-2980.
  24. Danje S. Fast pyrolysis of corn residues for energy production. Stellenbosch: Stellenbosch University; 2011.
  25. Ekanem S, Audu A. Lieberman MJJoCSO, proximate analysis of grain samples from strategic grain reserve facilities in Nigeria. 2022;47(3).
  26. Suleiman R, Rosentrater KA, Bern C, Effects of deterioration parameters on storage of maize: A review. Journal of Natural Sciences Research. 2013;3(9): 147-165.
  27. Sweets L, Stored grain fungi. Agricultural Electronic Bulletin Board University of Missouri Extension-CAFNR; 2018.
  28. Ape D, Nwogu N, Uwakwe E, Ikedinobi C, Comparative proximate analysis of maize and sorghum bought from Ogbete main market of Enugu State, Nigeria. Greener Journal of Agricultural Sciences. 2016. 6(9):272-275.
  29. Okonkwo C, Agharandu U. Proximate and vitamin composition of selected cereals commonly used for weaning babies' food preparation in South-Eastern Nigeria. Journal of Biology, Agriculture and Healthcare. 2017;7:22.
  30. Iken J, Amusa N, Obatolu VJJOTiA. Nutrient Composition and Weight Evaluation of some Newly Developed Maize Varieties in Nigeria. 2002;7(1):27-29.
  31. Nweke FNJAJoB. Rate of water absorption and proximate analysis of different varieties of maize cultivated in Ikwo Local Government Area of Ebonyi State, Nigeria. 2010;9(52):;8913-8917.
  32. Wronkowska MJJoFP and Preservation, Wet-milling of cereals. 2016;40(3):572-580.
  33. Deepak TS, Jayadeep PAJFT. Biotechnology, Prospects of maize (corn) wet milling by-products as a source of functional food ingredients and nutraceuticals. 2022;60(1):109-120.
  34. Nadiha MN, Fazilah A, Bhat R, Karim AA, Comparative susceptibilities of sago, potato and corn starches to alkali treatment. Food Chemistry. 2010;121(4): 1053-1059.
  35. Fadlalla HA, Abukhlaif HA, Mohamed SS. Effects of chemical and bio-fertilizers on yield, yield components and grain quality of maize (*Zea mays* L.). African Journal of Agricultural Research. 2016;11(45):4654-4660.
  36. Jiang H, Zhu Y, Wei L, Dai J, Song T, Yan Y, Chen S. Analysis of protein, starch and oil content of single intact kernels by near infrared reflectance spectroscopy (NIRS) in maize (*Zea mays* L.). Plant Breeding. 2007;126(5):492-497.
  37. Capuano E. The behavior of dietary fiber in the gastrointestinal tract determines its physiological effect. Critical reviews in food science and nutrition. 2017;57(16):3543-3564.
  38. Sadkiewicz JJEiT. International association for cereal science and technology ICC. 2005;13(5):215-216.
  39. Saleem M, Ahsan M, Aslam M, Majeed A. Comparative evaluation and correlation estimates for grain yield and quality attributes in maize. Pakistan Journal of Botany. 2008;40(6):2361-2367.
  40. Liu K. Effects of particle size distribution, compositional and color properties of ground corn on quality of distillers dried grains with solubles (DDGS). Bioresource Technology. 2009;100(19): 4433-4440.
  41. Egesel C, Kahrman F. Determination of quality parameters in maize grain by NIR reflectance spectroscopy. Journal of Agricultural Sciences. 2012;18(1):31-42.
  42. Ntuli V, Mekbib SB, Asita A, Molebatsi N, Makotoko M, and Chatanga P, Microbial and physicochemical characterization of

- maize and wheat flour from a milling company, Lesotho; 2013.
43. Maziya-Dixon B, Kling J, Okoruwa A. Physical, chemical and water absorption characteristics of tropical maize hybrids. African Crop Science Journal. 2000;8(4): 419-428.
44. Anyaegbu C, Oledibe O, Amadi JJEJoB. Proximate and phytochemical analysis of healthy and infected maize grains in Anambra State. 2020;5(2):15-40.

---

© 2023 Akash et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/110899>