



Development of Ready-to-eat (Spirulina Pancake Premix) Nutritional Supplement for Anemic Adolescent Girls: Nutritional and Microbial Analysis

Janhavi Srivastava^{a++}, Neetu Singh^{a#*} and Alka Nanda^{a†}

^a Department of Food and Nutrition, Food Science & Technology School for Home Science, Babasaheb Bhimrao Ambedkar University, Lucknow, 226025, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2023/v42i484328

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111223>

Original Research Article

Received: 15/10/2023

Accepted: 22/12/2023

Published: 24/12/2023

ABSTRACT

This study aimed to create an iron-rich dietary supplement utilizing spirulina and other nutritionally dense ingredients—quinoa, soybean, and amaranth—in the form of a pancake premix. The pancake premix was formulated, prepared, and analysed for its nutritional composition. The premix was meticulously developed by combining ingredients from various food groups, incorporating vanilla powder for flavour. The preparation involved simple steps to produce the final pancake product. Following the preparation, a thorough nutritional analysis was performed, covering moisture, ash, fat, protein, iron, crude fibre, and total carbohydrate content, adhering to established protocols such as AOAC methods.

⁺⁺ PG Scholar;

[#] Associate Professor;

[†] Research Scholar;

^{*}Corresponding author: E-mail: neetubbau@gmail.com;

The results revealed that the spirulina pancake premix contained 4.5% moisture, 12% ash, 4.6% fat, 11.7% protein, 67.8% carbohydrates, 5.7% iron, and 0.83% crude fibre. Furthermore, based on the Food Composition Table, the premix provided 460 kcal of energy, 16g of protein, 87g of carbohydrates, 3g of fat, and 8.5g of iron per serving, meeting substantial nutritional requirements. Microbial analysis indicated limited microbial growth, attributed to the airtight sealing of the premix in silver plastic bags, which effectively prevented contamination and extended the product's shelf life. The pancake premix demonstrated promising nutritional content and microbial control, suggesting its potential as an iron-rich dietary supplement for addressing nutritional deficiencies, particularly among anemic adolescent girls.

Keywords: Anemia; spirulina; premix; nutritional analysis; microbial analysis.

1. INTRODUCTION

Foods which are nutritionally dense and sustainable are high in demand for the food industries. It gets more serious to provide low-cost functional food when it comes to cope up with the deficiencies like anemia so to provide the needfuls to all the groups of the society. One of the low costs and readily available food is spirulina. Spirulina is a cyanobacterium that occurs in the form of either blue-green bacteria or blue-green algae which is available only in lakes with high alkalinity [1,2]. There is extensive documentation on indigenous populations across various regions using *S. platensis* as a food source [3]. The concentrated nutrition of spirulina makes it an excellent food supplement for people of all ages and lifestyles [4]. Spirulina is one of the natural sources containing the highest amount of protein, essential and nonessential amino acids, beta-carotene, a precursor of vitamin A, vitamin B12, and essential fatty acid γ -linolenic acid [5]. In addition to this spirulina also contains a host of other beneficial nutrients including; carotenoids, vitamin E, copper, manganese, magnesium, iron, selenium, and zinc [6]. Numerous research investigations demonstrate the wide array of health advantages offered by Spirulina, encompassing antioxidant, immune-modulating, anti-inflammatory, anticancer, antiviral, and antibacterial properties. Additionally, it exhibits positive effects against conditions such as hyperlipidemia, malnutrition, obesity, diabetes, toxicity induced by heavy metals and chemicals, as well as anemia [7,8,9,10]. Additionally, growing spirulina is better for the environment than making animal protein or plant alternatives. It's grown in controlled settings with low pollution and produces a lot in a small space [11].

Ready-to-eat foods require no additional preparation and are typically kept refrigerated or at room temperature and offers

the benefits of convenience, health and variety [12,13]. Hence, spirulina can be used preparing pancake premix by incorporating with other super flours such as quinoa, soy flour and amaranth. This would require no expert skills and is less time consuming.

Quinoa (*Chenopodium quinoa* Willd.) is a plant species in the Chenopodiaceae family that was first appeared in the Andes and can adapt regardless of soil and environmental circumstances [14]. It is a gluten-free pseudo cereal accompanied by great nutritional advantages because of its high protein, fat, fibre, essential fatty acids, vitamin, and mineral content [15,16]. Quinoa has a significant impact on the rheological, technical, and characteristics perceived by the senses in baked goods and could be used to manufacture bread products [17].

Soybean has been able to provide high-quality proteins at low prices [18]. Soy proteins are distinct among plant proteins due to their high biological value and the inclusion of necessary lysine, which is a limiting amino acid in most cereals [19,20]. Defatted soy flour (DSF) is a less expensive, more convenient, traditional, and high-protein source for the world's rapidly growing population [21].

Amaranth on the other hand, is a saviour in terms of nutrition for the malnourished world. Amaranth is a plant that exists in various variations, typically characterized by its wide leaves and purplish-green color, capable of growing up to a height of eight feet [22]. In India, amaranth is used widely in fasting rituals as it is considered as a "pseudocereal". It holds a wealth of nutrients like protein, lysine, carotenoids, dietary fibre, iron, vitamin C, vitamin A, riboflavin, thiamine, folic acid, calcium, among others. Moreover, it boasts a significant quantity of bioactive elements such as tocopherol (α , β , and γ), DPPH (2,2-diphenyl-1-picrylhydrazyl),

anthocyanins, lutein, and various phenolic compounds [23]. Apart from nutritional richness, it is also high in satiety value.

In this study, the aim was to prepare an iron rich dietary supplement for anemic adolescent girls and do nutritional analysis.

2. MATERIALS AND METHODS

2.1 Materials

The research was conducted at Babasaheb Bhimrao Ambedkar University's Food Science and Technology Laboratory (FSTL). The ingredients were procured from the local vendors of Lucknow and online retailer (Amazon & blink it) after designing a realistic meal that precisely reflects the desired proportion and nutrition.

2.2 Preparation of Premix Powder

The pancake premix powder was prepared by choosing at least one ingredient from each food groups given by Indian Council of Medical Research which are listed Table 1.

Apart from this, flavouring was achieved by adding vanilla powder into the mixture. The combined weight of all the components used to prepare the pancake premix was 155.5g per serving.

2.3 Preparation of Pancakes

The premix was prepared by combining all the ingredient in one bowl. After that, it was used to prepare the final product.

- **Prepare the Batter:** In a mixing bowl, pour the pancake premix according to the requirement. Add the specified amount of water and mix until the batter is smooth. Keep at rest for 10-15 minutes to set. Do not over mix the batter.
- **Preheat the Pan:** Heat a non-stick frying pan or griddle over medium heat. Optionally, add a small amount of butter or oil to avoid adherence.
- **Pour the Batter:** Once the pan is heated, pour a small amount of batter onto the cooking surface. Use a ladle or measuring cup to maintain consistent pancake sizes. Leave space between each pancake to allow for spreading.
- **Cook the Pancakes:** Cook the pancakes until bubbles form on the surface. This

usually takes around 2-3 minutes. Check the edges; they should start to look set.

- **Flip the Pancakes:** Once bubbles have formed and the edges look set, gently flip the pancakes using a spatula. Continue cooking for another 1-2 minutes on the opposite side until it turns a golden-brown color.
- **Repeat:** Continue making pancakes with the remaining batter, adding more oil or butter to the pan as needed to prevent sticking.
- **Serve:** Once all pancakes are cooked, stack them on a plate. Add your favourite toppings, such as syrup, honey, fresh fruit, nuts, or whipped cream.

2.4 Nutritional Analysis

Moisture: As per AOAC 2020, a petri plate was dried in an oven at 105 degrees Celsius for 20 minutes to remove the any moisture content. The petri dish was weighed without and then with the sample. 5 grammes of sample was maintained at 135 degrees for 2 to 3 hours. The sample was cooled, weighed, and the results were recorded. Finally, the calculations were completed by putting all of the values into the formula, which is:

$$\text{Moisture \%} = \frac{W_s - (W_2 - W_1)}{W_s} \times 100$$

Where, W_1 , weight of dish
 W_2 , weight of dish after drying
 W_s , weight of sample

Ash: Ash estimation was done on the basis of AOAC 2020 where a clean porcelain crucible was taken and dried properly to remove the moisture. The crucible was placed in a hot air for drying at 105 degrees Celsius for 20 minutes. The crucible was placed in the desiccator for cooling and was weighed empty. Then, a marginalized portion of the sample was kept in the pre weighed crucible and was weighed. At 700 degrees Celsius, the sample was maintained for two to three hours in the muffle furnace. Then, the sample was taken out and was kept in the desiccator for cooling. At last, the sample was weighed and calculation was done by putting the values into the formula:

$$\text{Ash\%} = \frac{W_2 - W_1}{W_s} \times 100$$

Where, W_1 , weight of crucible
 W_2 , weight of crucible with ash
 W_s , weight of sample

Table 1. Ingredients of spirulina pancake premix

| Food Groups | 1. Milk and Meat Products | 2. Pulses and Legumes | 3. Cereals, Grains and Products | 4. Fruits and Vegetables | 5. Fats and Sugar |
|-------------|---------------------------|-----------------------|---|--------------------------|-------------------------|
| Ingredients | Milk 30 g | Soya Flour 20g | Refined wheat flour 25 g Quinoa 20g Amaranth 5g | Date Powder 30g | Butter 15g Sugar 18g |




Fig. 1. Flow diagram of pancake preparation

Fat Estimation: As per AOAC 2020, the Soxhlet method was used to calculate the fat content of the product. With a filter paper, a thimble was made and weight on the prepare thimble was noted. 3 grams of the spirulina premix sample was taken into the thimble. Then after putting a cotton inside, thimble was folded to enclose the sample. the Soxhlet extraction unit was set by placing the sample in it. About 200ml of di-ethyl ether was added into the Soxhlet flask for the process of fat extraction. The sample was kept for 6 hours of continuous cycle for the completion of the process.

After cooling, the thimble was removed followed by collecting the remaining diethyl ether with extracted fat into a pre weighed beaker. The beaker was then put on a hot plate to evaporate the excess solution until the extracted fat was obtained in dried form.

The beaker with extracted fat was finally weighed and the crude fat was calculated by putting the obtained values in the formula

$$\text{Crude Fibre \%} = \frac{W_2 - W_1}{W_s} \times 100$$

Where, W_1 , weight of flask
 W_2 , weight of flask with fat
 W_s , weight of sample

Protein Estimation: The protein content of samples was estimated by the Kjeldahl method. 5gm of samples was introduced in digestion flask and to that 10 ml of concentrated H_2SO_4 and 5gm of digestion mixture of K_2SO_4 : $CuSO_4$: Na_2SO_4 (equal ratio) was added. The flask was stirred to fully combine the ingredients before being placed on a heater to begin digestion until the mixture became transparent (blue green in colour). The entire procedure took three hours to complete.

The resulting solution was brought to room temperature and put to a volumetric flask of 100 ml. Distilled water was used to top up the volume. Thereafter, ten millilitres of digest were gently added to the distillation tube, along with 10 ml of 0.5 N NaOH.

Due to the presence of NH_4OH , a yellowish tint emerged during distillation. Next, the distilled liquid was subjected to titration with a 0.25mol/L standard HCL solution until a pink colour was obtained. In order to determine how much titrant was utilised, the initial and final readings were recorded at this phase and marked as Vs. The nitrogen content of acetanilide or tryptophan after addition of 1 g of saccharose was determined at the titration stage for the blank, and the volume

of titrant utilised was indicated as V_b . The %N in samples was calculated via the given formula followed by the calculation of % P by multiplying the %N with the protein factor (PF) that is 6.38 [24].

$$\text{Protein \%} = \frac{V_s - V_b \times F \times C \times f \times M(N)}{m} \times 1000 \times 100$$

Where,

V_s - volume of titrant used for sample
 V_b - volume of titrant used for blank
 F- molar reaction factor of titrant (HCl-1 and H₂SO₄- 2)
 C- concentration of titrant (mol/L) = 0.25mol/L
 f- Factor of titrant = 1
 M(N)- Molecular weight of Nitrogen = 14.007g/mol
 m- sample weight
 1000- conversion factor (ml into L)
 %N- % weight of N

Iron Estimation: The concentration of metals in the samples was determined using Inductively Coupled Plasma-Mass Spectrometry (ICP-MS) with a Perkin Elmer instrument. Before the analysis, 100 μ L of the extracted samples was combined with 100 μ L of an internal standard (IS). This mixture was then adjusted to a total volume of 5 mL by adding a diluent comprising a solution of 15 mL methanol + 0.005% v/v Triton X + 10 mL nitric acid, topped up to 1000 mL with double distilled water. The metal concentrations were assessed by comparing them against a standard through a linearity curve [25,26].

Total Carbohydrate: Measurement of total carbohydrate content in a sample is calculated based on calculations.

$$\text{Total Carbohydrate} = 100\% - (\text{protein} + \text{fat} + \text{ash} + \text{water}).$$

Microbial Analysis: The pour plate method was employed to analyze the microbial content in Spirulina pancake premix powder. The process began with the preparation of nutrient agar, where the agar powder was mixed with distilled water, sterilized, and cooled to maintain its liquid state. Subsequently, a series of dilutions were made for the premix powder to facilitate the enumeration of viable microbial colonies. These dilutions were then aseptically transferred onto nutrient agar plates and evenly distributed. Around 15-20 ml of the cooled nutrient agar was carefully poured onto each plate to cover the

surface uniformly. Following this, the prepared plates were incubated at an appropriate temperature for microbial growth, typically ranging from 30-37°C. After 24-48 hours of incubation, colonies that developed on the plates were manually counted to estimate the microbial load in the original Spirulina pancake premix powder.

$$\text{CFU} = \frac{\text{Number of Colonies Counted} \times \text{Dilution Factor}}{\text{Volume Plated}}$$

Where,

Number of Colonies Counted: The count of visible colonies on the agar plate.

Dilution Factor: The factor used for dilution of the original sample.

Volume Plated: The volume of sample plated onto the agar plate.

3. PACKAGING OF PRODUCT

The premix powder was packed with no preservatives in silver disposable bags which were bought from a nearby market and were sealed to air tight to prevent spoilage from microbial growth.

Package labelling was done in this study as per Food Safety and Standard (Packaging & Labelling) regulation, 2011. The label was prepared with all the necessary details on Microsoft word and the printout was then pasted on the packets as finishing.

4. RESULTS AND DISCUSSION

The obtained results of the proximate analyse reveals that spirula pancake premix contains 4.5% of the total moisture content, 12% ash, 4.6 % fat, 11.7 % protein, 67.8% carbohydrates, 5.7 % iron and 0.83% crude fibre.

According to the Food Composition Table, the premix contains 460kcal of energy, 16g of protein, 87g of carbohydrates, 3g of fat, and 8.5 g of iron which is good enough to fulfil the requirements of nutrition in one meal.

It was found that there was less microbial growth which was 110 cfu/ml. Air tight sealing in silver plastic bags provided the same atmosphere to the food, hence prevent microbial contamination and increase the shelf life of the product.

Also, the study did not include all the labelling parameters as the product was produced on experimental basis and not for commercial purpose [27].




Fig. 2. Diagram showing proximate values of spirulina pancake premix




Fig. 3. Bar graph showing proximate values of spirulina pancake premix




Fig. 4. Label for packaging (Front and Back)

5. CONCLUSION

The development of an iron-rich dietary supplement in the form of a spirulina-based pancake premix, incorporating quinoa, soybean, and amaranth, presents promising prospects for combating nutritional deficiencies, especially anemia among adolescent girls. The comprehensive nutritional analysis revealed substantial protein, iron, and other vital nutrients within the premix, showcasing its potential to fulfill significant dietary requirements in a single serving. The limited microbial growth observed, owing to the effective airtight packaging, suggests an extended shelf life and enhanced safety of the product. These findings underscore the feasibility and convenience of utilizing this premix as a convenient and nutritionally dense supplement, highlighting its potential significance in addressing specific dietary deficiencies, particularly in vulnerable demographic groups. Further studies focusing on broader applicability, taste optimization, and clinical trials would reinforce its viability as an accessible solution for nutritional support.

DISCLAIMER

The goods employed in this study are widely and mostly used in our research area and country. There is no conflict of interest between the writers and manufacturers of the items because we want to use these products for the advancement of knowledge rather than for litigation. Furthermore, the research was not supported by the production firm, but rather by the authors' own personal efforts.

ACKNOWLEDGEMENT

The facility provided by the Department of Food and Nutrition, Babasaheb Bhimrao Ambedkar University, Lucknow for conducting this study in their laboratories is sincerely appreciated and the authors would like to express their gratitude for this.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumari DJ, Babitha B, Jaffar S, Prasad MG, Ibrahim MD, Khan MS. Potential

- health benefits of *Spirulina platensis*. *Int J Adv Pharm Sci*. 2011;2:417-22.
2. AlFadhly NKZ, Alhelfi N, Altemimi AB, Verma DK, Cacciola F, Narayanankutty A. Trends and technological advancements in the possible food applications of *Spirulina* and their health benefits: A review. *Molecules*. 2022;27(17):5584.
3. Bensehaila S, Doumandji A, Boutekrabt L, Manafikhi H, Peluso I, Bensehaila K, et al. The nutritional quality of *Spirulina platensis* of Tamenrasset, Algeria. *Afr J Biotechnol*. 2015;14(19):1649-54.
4. Sharoba AM. Nutritional value of *Spirulina* and its use in the preparation of some complementary baby food formulas. *J Food Dairy Sci*. 2014;5(8):517-38.
5. Mohan A, Misra N, Srivastav D, Umapathy D, Kumar S. *Spirulina*, the nature's wonder: A review. *Lipids*. 2014;5:7-10.
6. Dillon JC, Phuc AP, Dubacq JP. Nutritional value of the alga *Spirulina*. *World Rev Nutr Diet*. 1995;77:32-46.
7. Lee JB, Hayashi T, Hayashi K, Sankawa U, Maeda M, Nemoto T, et al. Further purification and structural analysis of calcium spirulan from *Spirulina platensis*. *J Nat Prod*. 1998;61(9):1101-4.
8. Lorenz RT. A review of *Spirulina* and *Haematococcus* algae meal as a carotenoid and vitamin supplement for poultry. *Spirulina pacifica* technical bulletin. 1999;53:1-14.
9. Hosseini SM, Khosravi-Darani K, Mozafari MR. Nutritional and medical applications of *Spirulina* microalgae. *Mini Rev Med Chem*. 2013;13(8):1231-7.
10. Kulshreshtha A, Zacharia AJ, Jarouliya U, Bhadauriya P, Prasad GB, Bisen PS. *Spirulina* in health care management. *Curr Pharm Biotechnol*. 2008;9(5):400-5.
11. Fantechi T, Contini C, Casini L. Pasta goes green: consumer preferences for *Spirulina*-enriched pasta in Italy. *Algal Res*. 2023;75:103275.
12. Kour DP, Sharma S, Gandotra A, Gupta N. Evaluation and development of healthy pancake premix from pseudocereals amaranth and buckwheat; 2022.
13. Muktawat P, Muktawat N, Varma N. Impact of ready to eat food taken by single living male and female. *Int J Sci Res Publ*. 2013;3(11):1-3.
14. El-Said ET, Soliman AS, Abbas MS, Aly SE. Treatment of anaemia and malnutrition by Shamy bread fortified with *Spirulina*,

- quinoa and chickpea flour. Egypt J Chem. 2021;64(5):2253-68.
15. Navruz-Varli S, Sanlier N. Nutritional and health benefits of quinoa (*Chenopodium quinoa* Willd.). J Cereal Sci. 2016;69:371-6.
 16. Stikic R, Glamoclija D, Demin M, Vucelic-Radovic B, Jovanovic Z, Milojkovic-Opsenica D, et al. Agronomical and nutritional evaluation of quinoa seeds (*Chenopodium quinoa* Willd.) as an ingredient in bread formulations. J Cereal Sci. 2012;55(2):132-8.
 17. Atef A, Abou-Zaid A, Wafaa SY, Emam H. Use of quinoa meal to produce bakery products to celiac and autism stuffs. Int J Sci Res. 2014;3(9):1344-54.
 18. Kulthe AA, Pawar VD, Kotecha PM, Chavan UD, Bansode VV. Development of high protein and low-calorie cookies. J Food Sci Technol. 2014;51(1):153-7.
 19. Riaz MN. Soybeans as functional foods. Cereal Foods World. 1999;44(2): 88-92.
 20. Kaur A, Ahluwalia P, Singh B. Studies on use of isabgol (*Psyllium mucilloid*) husk in atta for chapati making. J Food Sci Technol. 2005;38:75-8.
 21. Tripathi AK, Mishra AK. Soybean-a consummate functional food-a review. J Food Sci Technol. 2005;42:111-9.
 22. Tucker JB. Amaranth: The once and future crop. BioScience. 1986;36(1):9-13.
 23. Bhattarai G. Amaranth: A golden crop for future. Himalayan J Sci Technol. 2018;2:108-16.
 24. Kirk PL. Kjeldahl method for total nitrogen. Anal Chem. 1950;22(2):354-8.
 25. Şahan Y, Basoglu F, Gücer S. ICP-MS analysis of a series of metals (Namely: Mg, Cr, Co, Ni, Fe, Cu, Zn, Sn, Cd and Pb) in black and green olive samples from Bursa, Turkey. Food Chem. 2007;105(1):395-9.
 26. Chan KC, Yip YC, Chu HS, Sham WC. High-throughput determination of seven trace elements in food samples by inductively coupled plasma mass spectrometry. J AOAC Int. 2006;89(2):469-79.
 27. Longvah T, Ananthan R, Bhaskarachary K, Venkaiah K. Indian food composition tables; 2017.

© 2023 Srivastava et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/111223>