



Flexor Tendon Sheath Ganglion of the A1 Pulley Causing Trigger Thumb: A Case Report

**Abhishek Sugumar K. ^{a++}, Vidya Sagar ^{a++}, Raviraj A. ^{a++}
and Ashish Anand ^{b,c#*}**

^a Apollo Hospitals, Bannerghata Road, Bangalore-560076, India.

^b VAMC Jackson, University of Mississippi Med Ctr, Jackson, MS, USA.

^c William Carey School of Osteopathic Medicine, Hattiesburg, MS, USA.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJMPCR/2024/v17i1363

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112571>

Case Report

Received: 02/12/2023

Accepted: 07/02/2024

Published: 13/02/2024

ABSTRACT

Ganglion as cause of triggering of fingers have been described previously but they are not commonly described for the Thumb. We present the case of a 34-year-old lady who presented with triggering of the right thumb of few months duration. Patient chose to decline further imaging work up or injections. She elected to proceed with surgery. Surgical exploration revealed a 4x2mm ganglion in the A-1 pulley which was excised and resulted in complete resolution of the symptoms. To the best of our knowledge this is only the second described case report in English Literature.

Keywords: Ganglion; thumb; A1 pulley; trigger thumb; hand tumors.

⁺⁺ Consultant Orthopedic Surgeon;

[#] Staff Orthopedic Surgeon, Assoc. Prof. (Orthopedics) & Prof.;

^{*}Corresponding author: E-mail: ashishanandortho@yahoo.com;

1. INTRODUCTION

Stenosing tenosynovitis is a common cause of hand disability [1]. Bimodal distribution is observed with peaks observed in the 1st decade and in the 5th and 6th decades of life. It has a prevalence of 2-3% in the adult population [2-3]. Higher prevalence is seen in patients with diabetes mellitus, other endocrine disorders and inflammatory arthropathies [4].

Critical review of literature suggest that ganglion cysts flexor synovial sheath is uncommon cause of triggering. Ganglion cysts around the hand and wrist comprise about 60% of all hand tumors of which 5-16% occur along the flexor tendon sheaths [5-6]. Ganglion cyst of A1 pulley resulting in triggering of thumb is highly uncommon and this is only the second reported case in English literature.

2. CASE REPORT

A 34-year-old female-stay at home mother presented to the out-patient department with complaints of pain at palmar aspect of base of right thumb and triggering for 3 months.

Triggering was also elicited on clinical examination. A small firm non-tender swelling was palpable at the level of thumb flexion crease. It was not moving with tendon excursion. Patient was advised an ultrasound/Magnetic resonance Imaging but was declined by the patient. Patient had declined aspiration as well. She denied a traumatic event or similar swellings on contralateral side. Patient was agreeable for surgical excision. Following routine pre-anesthetic check and investigations patient was taken up for surgery.

At the thumb flexion crease local anesthetic infiltration was given and under loupe magnification transverse incision at the thumb flexion crease was used. A 1x1cm swelling was noted at the A1 pulley of the thumb and dissected out with care being taken to preserve the digital neurovascular bundle. Swelling was excised with surrounding portion of A1 pulley and sent for histopathological examination. Patient was relieved of triggering symptom. Patient was placed in a well-padded bulky dressing. Sutures were removed at 3 weeks and patient was seen at 3 months and 1 year following surgery. Patient had no recurrence noted at 1 year follow-up.



Fig. 1. Transverse incision at the thumb flexion crease. Neurovascular bundles dissected and retracted. Ganglion noted over the A1 pulley of thumb flexor sheath




Fig. 2. The 4x2mm Cyst, filled with gelatinous fluid

3. DISCUSSION

Ganglions are common hand tumors and account for about 50-70% of all upper limb swellings and a leading cause for the significant loss of normal hand function. The etiology of flexor sheath ganglions is still an enigma [6]. A higher prevalence is noted in females [7]. There are several hypotheses regarding the formation of these cysts including myxoid degeneration, increased production of hyaluronic acid and nonspecific proliferation of mesenchymal cells [7]. Doyle proposed possibility of tear in the synovial membrane and subsequent healing with an extracapsular segment of synovium that secretes synovial fluid [5]. Nelson et al proposed a post traumatic etiology [8-9]. In our patient no specific etiology could be attributed and presumably it was degenerative in origin.

A variety of treatment modalities exist for treatment of flexor sheath ganglion from manual compression [10] to ultrasound guided aspiration with or without steroid injection [10]. The disadvantage of steroid injection is high risk of recurrence and digital nerve injury. Surgical excision is the preferred mode of treatment as it avoids the pitfalls of aspiration and injection. Surgical excision of the cyst is combined with excision of a small portion of the pulley to prevent recurrence and enhance patient satisfaction.

To the best of our knowledge this is the second described case report in English Literature. Atlaf et al. [11] had previously reported Ganglion in A1 pulley as cause of triggering. The patient in their case report was a software professional and was using his hands on the computer. The authors reported possible over use as a cause of ganglion. Patient had relevant image work up done and had failed conservative treatment and subsequently underwent surgery with resolution of symptoms. In our case the patient was not indulging in excessive use of the computer and had declined advanced imaging work up or conservative treatment. Our patient underwent surgical excision and the histopathology observed was small Ganglion in A1 pulley which was excised. The Histopathology was similar to the case reported by Atlaf et al. [11].

4. CONCLUSION

Trigger finger and ganglion cysts of the flexor sheath are both common conditions, but their simultaneous occurrence is infrequent. Clinicians must devote careful attention to the diagnosis and open surgical excision of the A1 pulley ganglion under magnification to reduce the risk of neurovascular injury and recurrence.

CONSENT

As per international standards or university standards, patient written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Gil JA, Hresko AM, Weiss AC. Current concepts in the management of trigger finger in adults. J Am Acad Orthop Surg. 2020, Aug 1;28(15):e642-e650. DOI: 10.5435/JAAOS-D-19-00614. PMID: 32732655.
2. David M, Rangaraju M, Raine A. Acquired triggering of the fingers and thumb in adults. BMJ. 2017;359:j5285.
3. Lunsford D, Valdes K, Hengy S. Conservative management of trigger finger: A systematic review. J Hand Ther. 2019;32:212-221.
4. Kuczmarski AS, Harris AP, Gil JA, Weiss APC. Management of diabetic trigger finger. J Hand Surg Am. 2019;44:152-153.
5. Gunaseelan P, Jeremy P, Chua CK, Rashdeen F. Ganglion of the flexor tendon sheath at the A2 pulley - Case report. Malays Orthop J. 2015 Mar;9(1):28-29. DOI: 10.5704/MOJ.1503.002. PMID: 28435592; PMCID: PMC5349344.
6. Abe Y, Watson HK, Renaud S. Flexor tendon sheath ganglion: Analysis of 128 cases. Hand Surg. 2004, Jul;9(1):1-4. DOI: 10.1142/s0218810404001930. PMID: 15368618.
7. Jung, Kyu, Choi, Hwanjun, Kim Jun. Ganglion of the flexor tendon sheath between A1 pulley and A2 pulley. Archives of Hand and Microsurgery. 2014;23:29-32. DOI:10.15596/ARMS.2014.23.1.29.
8. Nelson CL, Sawmiller S, Phalen GS. Ganglions of the wrist and hand. J Bone Joint Surg Am. 1972;54:1459-64.
9. Al-Khawashki H, Hooper G. The distribution of fibrous flexor sheath ganglions. J Hand Surg Br. 1997;22:226-7.
10. Hvid-Hanson O. On the treatment of ganglia. Acta Chir Scand. 1970;136:471-6.
11. Altaf W, Attarde D, Sancheti P, Shyam A. Triggering of thumb by a ganglion cyst of the flexor tendon sheath at A1 pulley: A case report. Journal of Orthopaedic Case Reports. 2021, March;11(3):10-12.

© 2024 Sugumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/112571>