

## **Role of Intercropping in Sustainable Insect-Pest Management: A Review**

**Mohd Salim Mir <sup>a\*</sup>, Amal Saxena <sup>a</sup>, Raihana Habib Kanth <sup>a</sup>, Waseem Raja <sup>b</sup>,  
Khursheed Ahmad Dar <sup>a</sup>, S. S. Mahdi <sup>a</sup>, Tauseef Ahmad Bhat <sup>a</sup>,  
Nasir Bashir Naikoo <sup>c</sup>, Aijaz Nazir <sup>d</sup>, Zakir Amin <sup>e</sup>, Tajamul Mansoor <sup>c</sup>,  
May Zar Myint <sup>a</sup>, Mohammad Rafiq Khan <sup>a</sup>, Idrees Mohammad <sup>a</sup>  
and Suhail Ahmad Mir <sup>a</sup>**

<sup>a</sup> Division of Agronomy, Faculty of Agriculture, SKUAST-Kashmir, India.

<sup>b</sup> Mountain Research Centre for Field Crops, Khudwani, SKUAST-Kashmir, India.

<sup>c</sup> Division of Soil Science and Agricultural Chemistry, Faculty of Agriculture, SKUAST-Kashmir, India.

<sup>d</sup> Agromet Field Unit, Faculty of Horticulture, SKUAST-Kashmir, India.

<sup>e</sup> Division of Plant Pathology, Faculty of Agriculture, SKUAST-Kashmir, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJECC/2022/v12i111390

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92762>

**Review Article**

**Received 15 August 2022**  
**Accepted 20 October 2022**  
**Published 27 October 2022**

### **ABSTRACT**

Reduced soil fertility and rising pest and disease pressures are contributing to the already serious problem of global food insecurity. Monoculture is the most labour and resource-intensive form of crop production around the globe. Unfortunately, monocultures are more vulnerable to pests, diseases, and weeds, so the expansion of this system is accompanied by a host of biological issues. Negative effects on the environment, human health, and ecosystem stability are all associated with monocropping because it relies so heavily on the use of chemical plant protection products of all generations of pesticides. Although crop production strategies are important for overall enhancement in production, the intercropping can help farmers in attaining raised economic returns by taking multiple crops in a single season. Intercropping is an alternative strategy for improved resource use efficiency, environmental safety, and sustainable pest management without the use of chemical pesticides that can help mitigate these risks. Intercropping (two or more crop species coexisting) is a cultural practice in pest management that reduces insect pests by increasing ecosystem diversity. Intercropping and planting crops that kill or repel pests, attract

natural enemies, or have antibacterial effects can reduce disease and pest damage and pesticide use. Intercropping, where crops grow between main crops, reduces the likelihood of pest infestation. Intercropping is a potential pest management practice because it diversifies crops in an agro-ecosystem to reduce insect populations and attacks. Intercropping relies on a deep understanding of insect ecology and crop traits. Intercropping can be used alone or in combination with host-plant resistance and biological control. Intercropping ensures crop yield stability, protects against crop failure, improves soil fertility, increases soil conservation, and reduces pesticide use, minimizing agriculture's environmental impact. The aim is to define the role and importance of intercropping as a strategy in crop pest management and as a boost for crop production vis-à-vis soil fertility.

*Keywords: Environmental impacts; intercropping; pesticide; sustainable pest management; insect ecology; environmental safety; pest infestation.*

## 1. INTRODUCTION

A doubling (100-110%) of crop production relative to its 2005 level is required to feed the 9.6-12.3% increase in the global population by 2100 [1,2]. Losses of 10-16% of crop production due to pests [3,4] pose a real threat to entire world regions and significantly reduce agricultural yield [5,6,7,8]. Furthermore, there is growing concern that climate change will cause an increase in plant damage from pests in the coming decades [9,10,11]. Global yield losses of rice, maize, and wheat grains are projected to increase between 10 and 25% per degree of global mean surface warming. The average increases in yield losses due solely to pest pressure amount to 59, 92, and 62 metric megatons per year for wheat, rice, and maize, respectively, in a projected scenario of a 2°C warmer climate [12].

Alternative control methods within the framework of integrated pest management (IPM) have gained popularity as a result of the negative effects of synthetic pesticides [13] on the environment and human health [14], as well as reduced efficacy due to resistance within pest populations [15]. The "push-pull" strategy and the introduction of biological control agents are two common IPM systems that are widely used to achieve long-term control [16,17]. One important aspect of push-pull strategies is the use of volatile plant compounds to influence insect behavior [18,19,20]. Volatile compounds produced by plants are used by insects to identify and locate potential food and breeding plants [21,22]. So, non-host plants (e.g., aromatic plants) can be used to create insect repellents, anti-feedants, or insecticides because they release volatiles with repellent or deterrent properties in response to an attack [23,24]. However, the 'appropriate/inappropriate landings

theory' suggests that the presence of non-host plants may interfere with host-plant finding and host-plant acceptance behavior by giving insects a choice of green surfaces on which to land (host and non-host plant leaves) [25,26]. Attracting natural enemies to the area through intercropping, particularly with aromatic plants [27,28], providing food resources [29,30,31], or providing shelter and oviposition sites can all increase the efficacy of biological control. It is important to evaluate the effects of intercropped plants on both pests and natural enemies in order to optimize pest control in intercropping systems [32]. In this regard, an attempt was made to scrutinize the available literature and collect the information on the research carried out on intercropping worldwide over a period of time and look for the possible insights for future research in this domain.

Improving crop yield and income through intercropping is a time-honored agronomic practice [33,34,35,36,37,38]. To reduce the prevalence of pests and diseases and to encourage the growth of natural predators, it is an extremely useful strategy [39]. Reducing the damage caused by diseases and pests and lowering the need for pesticides can be accomplished through the strategic use of intercropping and the planting of crops that can kill or repel pests [40,41], attract natural enemies [42,43,44], or possess antibacterial effects in between the rows of economic crops [45,46,47,48]. Intercropping, which promotes plant diversity, can boost crop productivity and ensure greater and more consistent yields, bringing additional economic benefits [49]. As a result, it provides more financial security than monoculture and is thus well-suited to small farms that require a lot of human labor. As an added bonus, intercropping reduces the need for fertilizer and pesticides [50], lessening the negative effects that agriculture has on the

environment and allowing for more effective biological insect pest management.

## 2. OBJECTIVES OF INTERCROPPING

The major objectives of intercropping include

1. Intercropping is aimed at production maximization under limited land availability
2. It can serve as an alternate for generation of additional income through crop diversification.
3. It can serve as a soil fertility booster by improving the soil microflora and ecological diversification in soil.
4. It can help in minimizing the dependency on chemical fertilizers especially under legume intercropping system through improvement in soil fertility status and nutrient utilization capacity.
5. It can serve as a potential alternative for enhancing land use efficiency by occupation of limited arable land under diverse crops.

## 3. PROBLEMS ASSOCIATED WITH MONOCROPPING

Monocropping is the utilization of same land for single crop year after year. The concept is ideal for crop production but needs favorable conditions and may require higher production costs. However, the bottlenecks still remain as it does not provide any insurance under a complete crop failure owing to weather adversities or incidence of pests and diseases. The problems associated with monocropping include:

### 3.1 Higher Pest Infestation

Due to the lack of genetic diversity of plants, pests thrive on farms where only one type of crop is grown year after year [51].

### 3.2 Higher Pesticide Use

Reduced biodiversity makes monoculture crops more vulnerable to pests and diseases, prompting a higher reliance on pesticides to protect the crop, which in turn increases the likelihood of pest resistance [52,53].

### 3.3 Soil Degradation and Fertility Loss

The delicate equilibrium of soils is disrupted by monoculture farming. Growing the same crop

year after year depletes soil nutrients and leads to a decline in the diversity of bacteria and microorganisms [54].

### 3.4 Higher Use of Fertilizers

Growing only one type of crop on a given plot of land eventually depletes and exhausts the soil because it has no living organisms to replenish its natural diversity. Farmers often use chemical fertilizers, which disrupt the delicate balance of the ecosystem by altering the soil's natural composition, to artificially increase the fertility of their degraded fields [55].

### 3.5 Higher Water Use

If only one type of crop is grown on a piece of farmland, the soil will deteriorate and water will be wasted because the root systems aren't strong enough to hold the soil together. This is why such farmlands have an uneven distribution of water after rainfall. It is necessary for farmers to increase their water consumption to compensate for this loss.

### 3.6 Decrease in Biodiversity

As a result of the overuse of chemical pesticides in monoculture, biodiversity declines, and the ecosystem as a whole suffers.

### 3.7 Negative Impact on Pollinators

As pesticide use increases in monoculture, pollinating insects become sicker and eventually die off, and their absence from the ecosystem is a direct result. The use of chemicals has a serious impact on pollinizers and therefore it is often recommended to avoid chemical sprays during bloom.

### 3.8 Economic Risks

In the event of a catastrophic failure in crop development, such as an unusually long drought, unusually heavy rainfall, or an infestation of a particularly virulent strain of a pest, for example, a monoculture farm could lose its entire harvest all at once [56].

### 3.9 Environmental Degradation

Chemicals are overused in monoculture, leading to environmental degradation. These chemicals

not only deteriorate the soil health but also lead to deterioration of soil micro biota especially the beneficial microbes [57].

#### 4. PESTICIDE CONSUMPTION IN WORLD AND INDIA


Pesticides are the chemical or mixture of chemical substances aimed to prevent or destroy the losses due to biotic species likewise weeds, diseases, insects, etc., by mitigating the target pest. A pesticide is typically a chemical or biological agent, such as a virus, bacterium, antibiotic, or disinfectant, that discourages, renders ineffective, or kills pests. It is frequently applied to get rid of or manage a wide range of agricultural pests that can harm livestock and crops and lower farm output. Global production and use of pesticides are on the rise (Fig. 1) production of BHC began in India in 1952 at a plant in Calcutta, and today India is the largest producer of pesticides in Asia after China and the twelfth in the world, according to the Food and Agriculture Organization. Technical grade pesticide production in India has increased steadily from 5,000 metric tonnes in 1958 to 102,240 metric tonnes in 1998. In 1996–97, it was estimated that worldwide demand for pesticides would reach Rs. 22 billion (USD 0.5 billion), representing roughly 2% of the total market [58]. The graph shows that in the last seven decades, pesticide use in India has increased hundreds of times, from 154 MT in 1953-54 to 57,000 MT in 2016-17. India used 80,000 MT of pesticides in just one year between 1994 and 1995 [59].

The use of pesticides was highest in Maharashtra in 2016–17, followed by Uttar Pradesh, Punjab, and Haryana. Pesticide use was highest in Punjab (0.74 kg per acre), then Haryana (0.62 kg), and finally Maharashtra (0.40 kg) (0.57 kg). Fig. 2 shows that the two states of Maharashtra and Uttar Pradesh are responsible for 41% of India's total consumption of pesticides. Collectively, the six most populous states in India use more than 70 percent of the country's crop protection chemicals. Due to rising health and environmental awareness, many people today opt for non-chemical, all-natural alternatives to traditional, synthetic ones. Biopesticide is becoming increasingly popular as a result of its many advantageous properties, such as low toxicity to non-target organisms, high efficiency, rapid biodegradation, and suitability


for use in integrated pest management (IPM) initiatives. It's no secret that biopesticides could be used without negatively impacting the ecosystem [60].

##### 4.1 Adverse Effects of Pesticides

- Pesticides can have a major impact on the production workers, formulators, sprayers, mixers, loaders, and agricultural farm workers and all those who are directly or indirectly exposed. This can cause serious health hazards in humans and may damage the ecological diversity as well. Besides being possible human carcinogens, mutagens, and acetylcholinesterase inhibitors, most of these pesticides cause chronic toxicities like reproductive toxicity, genotoxicity, endocrine disruption, kidney damage, metabolic alterations, liver and bladder toxicity, gastrointestinal problems, etc.[61,62,63].
- The effects of pesticides on food commodities as a result of poisoning and increased residue retention on crops is a serious problem especially to the final consumers and the presence of residues in the food products makes the produce unacceptable in the global markets following the strict guidelines from international food standard committees like CODEX Alimentarius Commission, FSSAI, etc., [64,65].
- Pollution of water sources and other forms of plant life are just two examples of the environmental effects. Pesticides aren't just bad for the insects and weeds they're meant to kill; they can also be harmful to birds, fish, beneficial insects, and non-target plants.
- Surface and subsurface water contamination from untreated plant and soil runoff in addition to detrimental effects on soil health, such as those caused by persistent pesticides.
- Reducing the population of pollinators and beneficial soil microorganisms can result in a loss of biodiversity, which has an effect on soil fertility.
- Sprays of pesticides can contaminate nearby air, soil, and non-target plants in several ways: by landing directly on the plants, by drifting from the treated area, or by evaporating into the air.



**Fig. 1. Pesticide usage and consumption in World**  
Source: [56]



**Fig. 2. State wise pesticide consumption in India**  
Source: [56]

### 5. INTERCROPPING AS AN ALTERNATIVE TO CHEMICAL PESTICIDES FOR SUSTAINABLE INSECT PEST MANAGEMENT

❖ One of the key cultural practices in pest management, intercropping (where two or more crop species are grown together and coexist for a time) is predicated on the idea that increasing biodiversity in a given area will lead to a decrease in the prevalence of insect pests [66]. Therefore, reducing the harm inflicted by pests and diseases and lowering the need for pesticides can be

accomplished through the use of appropriate crop pairing for intercropping and the planting of crops that can kill or repel infestations, attract natural enemies or acquire antimicrobial action in between the rows of economic crops. Pests are less likely to infest intercropped crops than main crops. By increasing crop diversity within a given agro-ecosystem, intercropping has a potential to serve as a cultural practice based strategy for pest management by lowering the insect population and the frequency with which it attacks individual plants (Fig. 3).




Fig. 3. Illustration of benefits of intercropping

## 6. TYPES OF INTERCROPPING

### 6.1 Row-Intercropping

Planting two or more crops at once, with one or more crops planted in straight rows and the other(s) planted either next to or at random among them.

### 6.2 Mixed- Intercropping

Concurrently cultivating two or more crops in a field without using a definite row arrangement. This variety has the potential to thrive in pastures where grass and legumes are grown in symbiosis. When you plant a variety of crops together, you fortify your main crop against storms, freezes, and droughts.

### 6.3 Strip-Intercropping

To practice multi-cropping, two or more crop varieties are grown on the same plot of land in narrow strips that are wide enough to accommodate individual plantings but narrow enough to prevent physical interactions between the plants.

### 6.4 Relay- Intercropping

Multiple crop cultivation during a single growing season. Planting a second crop occurs when the first has completed its reproductive cycle but is still a few months away from being harvested.

### 6.5 Alley Cropping

Planting rows of food between hedges, trees, or bushes. The strong root systems of taller plants protect the soil from erosion and provide shade for the roots of shorter plants.

### 6.6 Trap Cropping

The intercropping method is useful for preventing damage to the primary crop by trapping pests. The basic idea is to lure pests or disease-causing organisms to the secondary crops instead of the primary cash crop.

### 6.7 Repellent Intercropping

The use of plants that naturally deter pests is a sustainable method of pest control that can be implemented in this cropping. Certain species are used for their natural repellent properties, which keep predators away from the cash crop.

## 7. MECHANISMS OF REDUCTION IN PEST INCIDENCE BY INTERCROPPING

Reducing the population growth rate of the attacking organism is one of three ways in which associated plants in an intercropping system can mitigate pest or disease attacks.

- ❖ When an associate crop is present, plants of the attacked component suffer (e.g. the host component is less vigorous and smaller in the intercrop than when sole-cropped).
- ❖ By diluting the host plant, the associate crop makes it more difficult for the attacking organism to do its damage-inducing work.
- ❖ The host's environment is altered by the associate crop in a way that benefits the organism's natural predators.

## 8. PUSH-PULL AGRICULTURAL PEST MANAGEMENT

- Intercropping with repellent "push" plants and trap "pull" plants is known as "push-pull technology," and it is used to reduce the number of agricultural pests. Stem borers, for instance, are a common pest of cereal crops like maize and sorghum (Fig. 4). Planting grasses around the crop's perimeter attracts and traps the pests, while planting plants like *Desmodium* in the spaces between the maize rows deters pests and helps keep *Striga* under control.

The ICIPE in Kenya and Rothamsted Research in the United Kingdom created the push-pull system.

### 8.1 Principles of Push-Pull Strategy of Insect Pest Management

The push-pull strategy seeks to maximize control efficacy, efficiency, sustainability, and output while minimizing unintended consequences to the environment. Each part of the strategy is unlikely to reduce pest populations as effectively as a single application of a broad-spectrum insecticide. Yet, when both push and pull elements are used together, performance improves [67]. Pests can be contained more effectively and their population reduced more rapidly by moving them to one central location. The use of broad-spectrum, synthetic insecticides is discouraged in favor of biological control methods or highly selective botanicals for population reduction. It is recommended that semiochemicals be made from renewable sources like plants, and this is increasingly possible [68]. If you want to get the most out of your farm without spending too much money, you should use harvestable intercrops or trap crops instead of sacrificial crops [69]. Understanding the biology of the pest and the behavioral/chemical ecology of its interactions with its hosts, conspecifics, and natural enemies is crucial to the development of reliable, robust, and sustainable push-pull strategies. Each tactic

**Table 1. Benefits and uncertainties of intercropping systems**

Benefits	Uncertainties
Efficient use of available land	Inadequate possibilities for production mechanization
Possibility of multiple harvests per year	Harvesting produce more challenging
Increasing crop diversity to ensure market supply	Demand for management has increased
Potential crop failure risks may be mitigated	There is no major harvest of staple or cash crops
Farmers may be able to handle price fluctuations	Disparately matched intercrops reduce yields of primary crops
Greater productivity and less wasteful use of materials	It's possible that increasing the soil's nitrogen levels through intercropping won't have much of an effect
Increase the amount of nitrogen in the soil gradually over time, especially if legumes are planted	The use of herbicides might be limited
Different plant species have different root systems, and this may lead to better soil structure	Possibility of allelopathy
Improving rotational control of soil erosion	Decline in crop specific pesticide use
Reduction in pests, diseases and weeds	May increase drudgery
Reducing eutrophication and emissions by decreasing the use of energy-intensive farming inputs	Potentially challenging intercultural operations



**Table 2. Recommended crops that impact the occurrence of pests under intercropping conditions**

SL. No.	Intercrop	Major Taxa	Main pest	Reference
1.	Alfalfa, carrot, maize, mungbean, wheat, sunflower, sorghum	Aphids	<i>Aphis gossypii</i>	[70]
2.	Cowpea, safflower, sunflower	Plant bugs	<i>Lygus Hesperus</i>	[71]
3.	Apricot	Thrips	<i>Thrips tabaci</i>	[72]
4.	Wheat	Predatory bugs	<i>Nabis sinoferus</i>	[73]
5.	Maize	Green Lacewings	<i>Chrysopa sinica</i>	[74]
6.	Soybean	Spider mite	<i>Tetranychus cinnabarinus</i>	[75]

**Table 3. Examples of crop pests for which changes in the behavior or development of the population have been observed due to intercropping**

Name of pest	Host plant	Type of intercropping	Changes in the pest behavior and pest population	Reference
<i>Acalymma vittata</i>	Cucumber	Inter-row cultivation, cucumber and corn or broccoli in separate rows	❖ Three times fewer beetles than in pure cucumber crop. ❖ Reduction in the reproductive rate and decrease in the period of foraging	[76]
<i>Phyllotreta Cruciferae</i>	Broccoli	Inter-row cultivation, broccoli in rows and vetch and bean between rows	❖ Decrease in the foraging period ❖ Decreasing of the population	[77]
<i>Aphis craccivora</i>	Groundnut	Row-crop mixture, groundnut and common beans in separate rows	❖ Common bean's sticky tendrils kept aphids away ❖ A reduction in aphids by decreasing foraging time	[78]
<i>Oulema spp</i>	Oat, barley	Row-crop mixture of both cereals	❖ Mixed cultivation of each species reduces the degree of damage to oat leaves by 48% and barley by 51% compared with pure stands	[79]
<i>Sitobion avenae</i>	Barley	Row-crop mixture of barley with yellow lupine and pea	❖ The number of aphids on barley heads was 3–6 times lower in crops with legumes	[80]

**Table 4. Examples of some intercrop used as trap crop in farming practices**

SL. No.	Main crop	Trap crop	Pest controlled
1.	Bengal gram	Marigold	<i>Heliothis sp.</i>
2.	Cowpea	Cotton	<i>Heliothis sp.</i>
3.	Soybean	Corn	<i>Heliothis sp.</i>
4.	Beans	Soybean	Mexican bean beetle
5.	Sunflower	Cotton and Marigold	<i>Heliothis sp.</i>
6.	Groundnut	Cowpea	Leaf folder
7.	Mustard	Cabbage	Cabbage head borer
8.	Cotton	Marigold	<i>Heliothis sp.</i>
9.	Tomato	Cabbage	Diamond back moth
10.	Brinjal	Coriander	Shoot and Fruit borer
11.	Cabbage	Radish	Flea beetle
13.	Radish	Cabbage	Flea beetle
14.	Potato	Marigold	Nematodes


Source: [81]



**Table 5. Intercropping for pest reduction – reported successful scientific trials**

SL. No.	Crop	Intercrop	Pest reduced/controlled	Mechanisms
1.	Apple	<i>Phacelia</i> sp.	San Jose scale, aphid	Parasitic wasps
2.	Barley	Alfalfa, red clover	Aphid	Predators
3.	Bean	Goosegrass	Leafhopper	Chemical repellent
4.	Cabbage	Tomato	Diamondback moth	Uncertain
5.	Carrots	Onion	Carrot fly	Chemical repellent
6.	Cauliflower	White or red clover	Cabbage aphid	Predators
7.	Collards	Tomato	Flea beetle	Chemical repellent
8.	Corn	Beans	Leafhoppers, beetle, fall armyworm	Physical interference, Predators
9.	Cow pea	Sorghum	Leaf beetle	Chemical repellent
10.	Crucifers	Wild mustard	Cabbageworm	Parasitic wasps
11.	Fruit trees	Wheat, sorghum	European red mite	Predators
12.	Walnut	Weedy ground cover	Walnut aphid	Parasitic wasps

Source: [82]



**Fig. 4. Showing push-pull pest management in maize crop**

uses a unique set of components that is tailored to the pest at hand (based on factors like its specificity, sensory abilities, and mobility) and the resource that is being guarded.

### 9. ADVANTAGES AND DISADVANTAGES OF INTERCROPPING SYSTEMS

Table 1 outlines the positives and negatives of intercropping systems in agriculture. Table 2 lists crops that are recommended for their ability to reduce pest occurrence when grown in between other crops. In Table 3, several examples of crop pests whose population dynamics have been

observed to shift as a result of intercropping are given. In Table 4, some examples of intercrops used as trap crops in farming practices are presented, and in Table 5, the results of scientific trials are given, that have used intercropping as a tool for reducing pest populations.

### 10. CONCLUSION

Intercropping is an additional workable strategy for warding off insect pests in crops. Diversifying crops in a given agro-ecosystem can help lower the insect population and, in turn, the risk of attack, making it a promising cultural practice for

pest management. There are a few different ways in which intercropping protects the primary crop. The choice of companion crops and their additional valuation after harvest, as well as the farmers' knowledge and the mechanization practiced, play a role in the success of intercropping for pest management. Planting multiple crop types in a single field is known as intercropping, a cultural practice used in the IPM system to reduce the need for pesticides. Concerns about potential negative impacts of pesticide on human health and the environment, pesticide resistance, the resurgence of insect pests, and general considerations of agricultural production led to the development of intercropping as an IPM tool. Since not all possible crop combinations have the desired effect, figuring out which ones will reduce pest abundance is a major challenge when trying to select the best intercrop combination for pest suppression. Therefore, intercropping can either stand on its own as a pest management strategy or be used in tandem with others, such as host-plant resistance, supplementary biological control, and chemical control. To effectively use intercropping for insect pest management in today's modern agriculture, one must have an in-depth understanding of how different crop characteristics and combinations will affect the behaviour of pests.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Gerland P. World population stabilization unlikely this century. *Science* 2014;346:234-237.
- Tilman D, Balzer C, Hill J, Befort BL. Global food demand and the sustainable intensification of agriculture. *Proc. Natl Acad. Sci. USA*. 2011;108:220-260.
- Flood J. The importance of plant health to food security. *Food Secur.* 2010;2:215-231.
- Chakraborty S, Newton AC. Climate change, plant diseases and food security: An overview. *Plant Pathol.* 2011;60:2-14.
- Oerke EC. Crop losses to pests. *J. Agric. Sci.* 2006;144:31-43.
- Strange RN, Scott PR. Plant disease: A threat to global food security. *Annu. Rev. Phytopathol.* 2005;43:83-116.
- Fisher MC, Henk DA, Briggs CJ, Brownstein JS, Madoff LC, McCraw SL, Gurr SJ. Emerging fungal threats to animal. *Plant and Ecosystem Health.* *Nature.* 2012;484:186-194.
- Khan ZR, Pickett JA, van den Berg J, Wadhams LJ, Woodcock CM. Exploiting chemical ecology and species diversity: Stem borer and striga control for maize and sorghum in Africa. *Pest Manage. Sci.* 2000;56:957-962.
- Thomson LJ, Macfadyen S, Hoffmann AA. Predicting the effects of climate change on natural enemies of agricultural pests. *Biol. Control.* 2010;52:296-306.
- Howden SM, Soussana JF, Tubiello FN, Chhetri N, Dunlop M, Meinke H. Adapting agriculture to climate change. *Proc. Natl Acad. Sci. USA.* 2007;104:19691-19696.
- Gu S, Han P, Ye Z, Perkins LE, Li J, Wang H, Zalucki M, Lu Z. Climate change favours a destructive agricultural pest in temperate regions: Late spring cold matters. *J. Pest Sci.* 2018;91: 1191-1198.
- Deutsch CA, Tewksbury JJ, Tigchelaar M, Battisti DS, Merrill SC, Huey RB, Naylor RL. Increase in crop losses to insect pests in a warming climate. *Science.* 2018;361: 916-919.
- Lai RQ, You MS, Jiang LC, Lai BT, Chen SH, Zeng WL, Jiang DB. Evaluation of garlic intercropping for enhancing the biological control of *Ralstonia solanacearum* in flue-cured tobacco fields. *Biocontrol Science and Technology.* 2011; 21:755-764.
- Degri MM, Mailafiya DM, Mshelia JS. Effect of intercropping pattern on stem borer infestation in Pearl millet (*Pennisetum glaucum* L.) grown in the Nigerian Sudan Savannah. *Advances in Entomology.* 2014;2:81-86.
- Brewer MJ, Goodell PB. Approaches and incentives to implement integrated pest management that addresses regional and environmental issues. *Annu. Rev. Entomol.* 2011;57:41-59.
- Brodeur J, Abram PK, Heimpel GE, Messing RH. Trends in biological control: Public interest, international networking and research direction. *Biocontrol.* 2018; 63:11-26.
- Hassanali A, Herren H, Khan ZR, Pickett JA, Woodcock CM. Integrated pest management: The push-pull approach for

- controlling insect pests and weeds of cereals, and its potential for other agricultural systems including animal husbandry. *Philos. Trans. R. Soc. B Biol. Sci.* 2008;363:611-621.
18. Dudareva N, Negre F, Nagegowda DA, Orlova I. Plant volatiles: Recent advances and future perspectives. *Crit. Rev. Plant Sci.* 2006;25:417-440.
  19. Shrivastava G, Rogers M, Wszelaki A, Panthee DR, Chen F. Plant volatiles-based insect pest management in organic farming. *Crit. Rev. Plant Sci.* 2010;29:123-133.
  20. Beck JJ, Torto B, Vannette RL. Eavesdropping on plant-insect-microbe chemical communications in agricultural ecology: A virtual issue on semiochemicals. *J. Agric. Food Chem.* 2017;65:5101-5103.
  21. Kuhnle A, Muller C. Relevance of visual and olfactory cues for host location in the mustard leaf beetle *Phaedon cochleariae*. *Physiol. Entomol.* 2011;36:68-76.
  22. Wynde FJH, Port GR. The use of olfactory and visual cues in host choice by the capsid bug *Lygus rugulipennis* Poppius and *Liocoris tripustulatus* Fabricius. *PLoS ONE* 2012;7:e46448.
  23. War AR, Paulraj MG, Ahmad T, Buhroo AA, Hussain B, Ignacimuthu S, Sharma HC. Mechanisms of plant defence against insect herbivores. *Plant Signal. Behav.* 2012;7:1306-1320.
  24. Bakkali F, Averbeck S, Averbeck D, Idaomar M. Biological effects of essential oils-A review. *Food Chem. Toxicol.* 2008; 46:446-475.
  25. Finch S, Collier RH. The influence of host and non-host companion plants on the behaviour of pest insects in field crops. *Entomol. Exp. Appl.* 2011;142:87-96.
  26. Finch S, Billiald H, Collier RH. Companion planting-Do aromatic plants disrupt host-plant finding by the cabbage root fly and the onion fly more effectively than non-aromatic plants? *Entomol. Exp. Appl.* 2003;109:183-195.
  27. Batista MC, Fonseca MCM, Teodoro AV, Martins EF, Pallini A, Venzon M. Basil (*Ocimum basilicum* L.) attracts and benefits the green lacewing *Ceraeochrysa cubana* Hagen. *Biol. Control.* 2017;110:98-106.
  28. Togni PHB, Venzon M, Muniz CA, Martins EF, Pallini A, Sujii ER. Mechanisms underlying the innate attraction of an aphidophagous coccinellid to coriander plants: Implications for conservation biological control. *Biol. Control* 2016;92:77-84.
  29. Tang GB, Song BZ, Zhao LL, Sang XS, Wan HH, Zhang J, Yao YC. Repellent and attractive effects of herbs on insects in pear orchards intercropped with aromatic plants. *Agrofor. Syst.* 2013;87: 273-285.
  30. Song B, Zhang J, Wiggins NL, Yao Y, Tang G, Sang X. Intercropping with aromatic plants decreases herbivore abundance, species richness, and shifts arthropod community trophic structure. *Environ. Entomol.* 2012;41:872-879.
  31. Song BZ, Wu HY, Kong Y, Zhang J, Du YL, Hu JH, Yao YC. Effects of intercropping with aromatic plants on the diversity and structure of an arthropod community in a pear orchard. *Biocontrol.* 2010;55:741-751.
  32. Raseduzzaman M, Jensen ES. Does intercropping enhance yield stability in arable crop production? A meta-analysis. *European Journal of Agronomy.* 2017;91: 25-33.
  33. Rao MR, Mathuva MN. Legumes for improving maize yields and income in semi-arid Kenya. *Agriculture, Ecosystems & Environment.* 2000;78:123-137.
  34. Singh RJ, Ahlawat IPS. Productivity, competition indices and nutrients dynamics of Bt cotton (*Gossypium hirsutum* L.)-groundnut (*Arachis hypogaea* L.) intercropping system using different fertility levels. *Indian Journal of Agricultural Sciences.* 2011;81:606-611.
  35. Lopes T, Hatt S, Xu Q, Chen J, Liu Y, Francis F. Wheat (*Triticum aestivum* L.)-based intercropping systems for biological pest control. *Pest Management Science.* 2016;72:2193-2202.
  36. Himmelstein J, Ares A, Gallagher D, Myers J. A meta-analysis of intercropping in Africa: Impacts on crop yield, farmer income, and integrated pest management effects. *International Journal of Agricultural Sustainability.* 2017;15:1-10.
  37. Kumar R, Turkhede AB, Nagar RK, Kumar R. Effect of American cotton based intercropping system on yield, quality and

- economics. Research in Environment and Life Sciences. 2017;10: 75-77.
38. Singh A, Weisser WW, Hanna R, Houmgny R, Zytynska SE. Reduce pests, enhance production: Benefits of intercropping at high densities for okra farmers in Cameroon. Pest Management Science. 2017;73:2017-2027.
  39. Redlich S, Martin EA, Steffan-Dewenter I. Landscape level crop diversity benefits biological pest control. Journal of Applied Ecology. 2018;55:2419-2428.
  40. Schader C, Zaller JG, Köpke U. Cotton–basil intercropping: Effects on pests, yields and economical parameters in an organic field in Fayoum, Egypt. Biological Agriculture & Horticulture. 2005;23:59-72.
  41. Wang Y. Scientific intercropping can reduce pests and diseases. Pesticide Market News. 2015;10:47-68.
  42. Xue L. Preliminary analysis on ecological control of pests and diseases. Modern Agriculture. 2015;7:40.
  43. Ning C, Qu J, He L, Yang R, Chen Q, Luo S, Cai K. Improvement of yield, pest control and Si nutrition of rice by rice–water spinach intercropping. Field Crops Research. 2017;208:34-43.
  44. Baidoo PK, Mochiah MB, Apusiga K. Onion as a pest control intercrop in organic cabbage (*Brassica oleracea*) production system in Ghana. Sustainable Agriculture Research. 2012;1:50-59.
  45. Seni A. Role of intercropping practices in farming system for insect pest management. Acta Scientific Agriculture. 2018;2:8-11.
  46. Gibbons D, Morrissey C, Mineau P. A review of the direct and indirect effects of neonicotinoids and fipronil on vertebrate wildlife. Environmental Science and Pollution Research. 2015;22:103-118.
  47. Zhao ZH, Hui C, Hardev S, Ouyang F, Dong Z, Ge F. Responses of cereal aphids and their parasitic wasps to landscape complexity. Journal of Economic Entomology. 2014;107:630-637.
  48. Jeyasankar A. Antifeedant, Insecticidal and growth inhibitory activities of selected plant oils on black cutworm, *Agrotis ipsilon* (Lepidoptera: Noctuidae). Asian Pacific Journal of Tropical Disease. 2012;2:347-351.
  49. Khan MR, Raja W, Bhat TA, Mir MS, Naikoo NB, Amin Z, Nazir A, Mir SA, Mohammad I, Wani AA, Patyal D. Zero budget natural farming: A way forward towards sustainable agriculture. Current Journal of Applied Science and Technology. 2022;41(13):31-43.
  50. Mir MS, Naikoo NB, Amin Z, Bhat TA, Nazir A, Kanth RH, Singh P, Raja W, Singh L, Fayaz S, Ahngar TA, Palmo T, Rehman U. Integrated farming system: A tool for doubling farmer's income. Journal of Experimental Agriculture International. 2022;44(3):47-56.
  51. Naikoo NB, Chesti MH, Bhat MA, Mir AH, Bashir O, Bhat TA, Mir MS, Gull A, ZA, Ayoub L. Biostimulants towards Soil Health Improvement: A Review. Agricultural Reviews. 2022;R-2526:1-8.
  52. Mir MS, Naikoo NB, Kanth RH, Bahar FA, Bhat MA, Nazir A, Mahdi SS, Amin Z, Singh L, Raja W, Saad AA, Bhat TA, Palmo T, Ahngar TA. Vertical farming: The future of Agriculture: A review. The Pharma Innovation Journal. 2022;11(2): 1175-1195.
  53. Kumar S. Use of pesticides in agriculture and livestock animals and its impact on environment of India. Asian Journal of Environmental Science. 2013;8(1) 51-57.
  54. Agnihotri, N. Pesticide Consumption in Agriculture in India - An Update. Pestic. Res. Journal. 2000;150-155.
  55. Kumar S, Singh A. Biopesticides: Present status and the future prospects. J. Fertil. Pestic. 2015;6.
  56. FAO; 2020. Available:<http://www.fao.org/faostat>
  57. Rahimi V, Madadi H, Nazari BS. Effect of additive series intercropping kidney bean (*Phaseolus vulgaris* L.) with some aromatic plants on *Tetranychus urticae* Koch. (Tetranychidae) population. In: Proceedings of the International Electronic Conference on Entomology; 2021
  58. Duraimurugan P, Regupathy. Push-pull strategy with trap crops, neem and nuclear polyhedrosis virus for insecticide resistance management in *Helicoverpa armigera* (Hubner) in cotton. Am. J. Appl. Sci. 2005;2:1042-48.
  59. Nalyanya G, Moore CB, Schal C. Integration of repellents, attractants, and insecticides in a “push-pull” strategy for managing German cockroach (Dictyoptera: Blattellidae) populations. J. Med. Entomol. 2000;37:427-34.

60. Birkett MA, Pickett JA. Aphid sex pheromones: from discovery to commercial production. *Phytochemistry*. 2003;62:651-56.
61. Huang Y, Shi T, Luo X, Xiong H, Min F, Chen Y, Nie S, Xie M. Determination of multi-pesticide residues in green tea with a modified QuEChERS protocol coupled to HPLC-MS/MS. *Food chemistry*. 2019;275: 255-64.
62. Ramakrishnan B, Venkateswarlu K, Sethunathan N, Megharaj M. Local applications but global implications: Can pesticides drive microorganisms to develop antimicrobial resistance? *Science of the Total Environment*. 2019;654:177-89.
63. Zhang M, Zeiss MR, Geng S. Agricultural pesticide use and food safety: California's model. *Journal of Integrative Agriculture*. 2015;14(11):2340-57.
64. Joint FAO/WHO Codex Alimentarius Commission. *Codex alimentarius*. Food & Agriculture Org.; 1992.
65. FSSAI. Food Safety and Standards. Version-III (31.08.2018) (1) (2) (3). 2018. Available: [https://agriexchange.apeda.gov.in/food\\_standards/ind\\_document/Compendium\\_Contaminants\\_Regulations\\_07\\_09\\_2018.pdf](https://agriexchange.apeda.gov.in/food_standards/ind_document/Compendium_Contaminants_Regulations_07_09_2018.pdf)
66. Bruce TJA, Birkett MA, Blande J, Hooper AM, Martin JL, et al. Response of economically important aphids to components of *Hemizygia petiolata* essential oil. *Pest Manag. Sci*. 2005;61: 1115-21.
67. Hooper AM, Farcet JB, Mulholland NP, Pickett JA. Synthesis of 9-methylgermacrene B, racemate of the sex pheromone of *Lutzomyia longipalpis* (Lapinha), from the renewable resource, *Geranium macrorrhizum* essential oil. *Green Chem*. 2006;8:513-15.
68. Khan ZR, Pickett JA. The 'push-pull' strategy for stemborer management: A case study in exploiting biodiversity and chemical ecology. In *Ecological Engineering for Pest Management: Advances in Habitat Manipulation for Arthropods*, ed. GMGurr, SDWratten, MA Altieri, Wallington, Oxon, UK: CABI. 2004; 155-64.
69. Lu XY, Li SY, Zhu JB, Cheng FR, Liu FZ, Jiang BL, Yu C. Effect of spring corn and cotton intercropping on the population dynamics of *Aphis gossypii* and *Rhopalosiphum maidis*. *China Cotton*. 2017;44:24-27.
70. Yan S, Yu J, Han M, Michaud JP, Guo LL, Li Z, Zeng B, Zhang QW, Liu XX. Intercrops can mitigate pollen-mediated gene flow from transgenic cotton while simultaneously reducing pest densities. *Science of the Total Environment*. 2020; 711:134855.
71. Zhang LJ, Gao XK, Luo JY, Zhang S, Wang CY, Zhu XZ, Wang L, Cui JJ. Strategy for resistance risk management of Bt transgenic cotton in Australia. *China Cotton*. 2018;45:3-6.
72. Wang W, Yao J, Zhang Y, Liu H. Effect of apricot trees on insect pests and their natural enemies in nearby cotton fields in southern Xinjiang. *Chinese Journal of Applied Entomology*. 2012;49: 951-956.
73. Chen M, Luo JC, Li GQ. Evaluating alfalfa cutting as a potential measure to enhance predator abundance of *Aphis gossypii* (Homoptera: Aphididae) in cotton-alfalfa intercropping system. *Acta Agrestia Sinica*. 2011;19:922-926.
74. Luo J, Zhang S, Wang C, Lyu L, Li C, Cui J. Ecological effects of different trap crop to sucking pests and natural enemies in cotton fields. *China Cotton*. 2014;41:14-16.
75. Zhang XM, Liu XJ, Yang YM, Duan DY. Occurrence of *Lygus lucorum* in the Bt cotton and jujube field. *Entomological Journal of East China*. 2005;14:28-32.
76. Andrews DJ, Kassam A.H. The importance of multiple cropping in increasing world food supplies. *Am. Soc. Agron*. 1976;27: 32.
77. Garcia L, Celette F, Gary C, Ripoche A, Valdes-Gomez H, Metay A. Management of service crops for the provision of ecosystem services in vineyards: A review. *Agric. Ecosyst. Environ*. 2018;251:158-170.
78. Farrell JAK. Effects of intersowing with beans on the spread groundnut rosette virus by *Aphis craccivora* Koch (Homoptera: Aphididae) in Malawi. *Bull. Entomol. Res*. 1976;66:331-333.
79. Wenda-Piesik A, Piesik, D. The spring cereals food preferences of *Oulema* spp. in pure and mixed crops. *Elec. J. Pol. Agric. Univ. Agron*. 1998;1:1-12.

80. Wenda-Piesik, A. Health status of spring barley grown in monocrop and in mixtures with cereals or leguminous plants. J. Plant Prot. Res. 2001;41:388-394.
81. Dewangan SR, GD Sahu. "Trap cropping- A valuable pest management technique". Popular Kheti. 2014;131-134.
82. Miguel A. Altieri, Clara I. Nicholls with Marlene A. Fritz. In: Manage insects in your farm: A guide to ecological strategies. Recent Advances in Ecological Pest Management. Sustainable Agriculture Research and Education (SARE). 2014; 107-109.

---

© 2022 Mir et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/92762>