



The Effect of Demographic Factors on Work-related Injuries in the Oil and Gas Industry in the Niger Delta Region in Nigeria

Oseibhi Oyamienlen ^a, Ify L. Nwaogazie ^{a*}
and Chinemerem Patricks ^a

^a Centre for Occupational Health, Safety and Environment (COHSE), University of Port Harcourt, Choba, Port Harcourt, Rivers State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ACRI/2023/v23i2554

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96301>

Original Research Article

Received: 02/12/2022

Accepted: 06/02/2023

Published: 13/02/2023

ABSTRACT

This study aimed to find out the effect of Demographic factors on work-related Injuries in the Oil and Gas Industry in the Niger Delta Region in Nigeria. The research method that was used for this study was a correlation design to test the demographic variables and how related they were to work-related injuries. The instrument used for this study was a well-structured questionnaire on Analysis of Work-related Accidents which was sent to e-mails of oil and gas workers in Akwa Ibom state and the Rivers state of Nigeria for the collection of data to provide answers to the questions. The reliability of the instrument was assessed using Cronbach's alpha which was 0.767, Different statistical test methods (Mann-Whitney, Kruskal Wallis, z-test, ANOVA and Dunn's test) were used. The findings revealed that gender does not play any significant role in work-related injuries with p-

*Corresponding author: Email: ifynwaogazie@yahoo.com;

values greater than 0.05, Education plays a role in certain types of injuries i.e. (contusion and MSDs with P-values of 0.013 and 0.015 respectively, and musculoskeletal disorders were the common types of injuries experienced by staff with a p-value of 0.05. Fractures and Amputations were experienced by workers who have spent above 16 years working offshore with p-values of 0.031, 0.035 and 0.020, 0 & 022 respectively. In Summary, certain demographic factors have effect on work-related Injuries.

Keywords: Demographic factors; work-related injury; oil and gas industries.

1. INTRODUCTIONS

Occupational or work-related injuries became more common after the industrial revolution. Over time, in response to the great demand for better working conditions, the first occupational laws were enacted in the 19th century in England. Occupational injuries are associated with lots of suffering and loss at individual, community, societal and organizational levels [1]. Occupational accidents and ill health can significantly hurt the profitability of the business and, it can lead a production platform out of business because WRAs and diseases cost money. When an accident occurs there will be direct costs that arise directly from the incident itself and indirect costs which arise indirectly as a consequence of the more significant incidents [2]. Occupational accidents and work-related diseases result in more than 2.3 million fatalities globally every year, of which over 350,000 results from occupational accidents and around 2 million results from work-related diseases. Statistics show that every 15 seconds, one worker dies from a work-related incident, and 153 workers have a work accident [3]. Additionally, 6,300 people pass away daily as a result of work-related incidents or work-related ill-health [3].

The majority of the oil and gas resources are domiciled in offshore fields. Past studies [4] indicate that there are over 7270 offshore oil and gas installations in place globally with distribution in more than fifty-three countries. Consequently, Oil and gas are the cornerstones of growth and production across the nations because it is omnipresent in all sectors, which makes them indispensable [5,6]. Offshore oil and gas operations are highly associated with life-threatening risk factors for rig workers. Every year, hundreds of people die and thousands of them are injured while performing their tasks in oil and gas drilling and production industries [7]. Leigh et al. [8] Mekkodathil, et al. [1] ranks oil and gas industries as one of the industries that spend high costs on occupational injuries. Even

in high-income economies, occupational injury figures remain remarkably high [9]. This argument is hinged on the nature of these projects which are characterized by having extreme locations, harsh environmental conditions, and a large number of participants from diverse orientations, etc [10,11]. Furthermore, the offshore oil and gas industry also presents a typical example of a safety-critical sector because of the interplay of inherent technological, environmental, and human challenges According to the International Labor Organization [12], due to the unpredictable and hazardous nature of the operation, several chemical, safety, environmental and ergonomic hazards have been reported for decades in all over the world in the sector [13,14].

Work-related accidents (WRAs) constitute a social phenomenon and one of the major problems all over the world [3]. Despite all the implemented safety control measures in any organization and the improvements in work-related safety which have been known in the last decades, things can still go wrong, and accidents and incidents still do happen. Ignoring the health and safety of workers cannot be the stepping stone to a good corporate sustainability strategy. The incident rates have increased alongside the industrial revolution and the rapid globalization of the world. Research on Occupational and non-occupational factors associated with work-related injuries among construction workers in the USA have also found disparities in injury risk by race/ethnicity, gender, age, geographic region, and education [15]. This study reviews the literature on the association between demographic variables (e.g., gender, Education, work experience, and types of staff) with work-related injuries.

2. METHODS

The type study was a correlation design to test the demographic variables and how related they were to work-related injuries. A stratified sampling technique was used for this study

which involves dividing the entire population into smaller groups or strata (according to their Department). The researcher stratified the population before proportionally selecting a sample at random in 2021. The instrument used for this study was a Consistency reliability which was estimated using Cronbach's alpha coefficient formula [16]. The validity of the instrument was tested by first administering the questionnaires to panel of experts including the study supervisors and health and safety experts in the offshore oil and gas industry who will

evaluate and rate the structure, contents, clarity, consistency and relevancy of the instrument to the research. This process enabled the researcher to identify ambiguities and inconsistencies in the questionnaire constructs and thus, corrected to address the main focus in the research study. A well-structured questionnaire was sent to e-mails of oil and gas workers in Akwa Ibom state and the Rivers state of Nigeria for the collection of data to provide answers to the questions.




Fig. 1. Map of Nigeria showing Niger Delta Area, the Oil & Oil servicing companies in the region

Source: <https://www.sec.gov/Archives/>

The questionnaire contains six sections consisting of questions on demographics, hazards likely to occur in an offshore environment, injury types experienced by offshore workers, injured body parts prevalent among offshore workers, types of accidents in the offshore environment and consequences of accidents in the offshore environment were used in the data collection phase of the study. The study was performed using a population of personnel that work offshores in selected oil and gas industries in Akwa Ibom and Rivers state respectively. The population of this study covers Maintenance personnel, Marine Engineers, Mooring masters, Operations personnel, Engineers, Asset owners, Emergency Responders, and Medics in selected oil and gas offshore facilities of the company operating in Rivers and Akwa Ibom state irrespective of the level of education, sex, job specialty, ethnicity, etc.

2.1 Instruments

The research instrument adopted was the questionnaire method. A variety of four-point Likert scales ranging from 1 (strongly disagree) to 4 (strongly agree) was used. The constructs for the work-related injuries and some demographic criteria were obtained from well-Structured Health and Safety questionnaires [17] after it was vetted by health and safety expert and Cronbach's alpha was used for the reliability test. The questionnaire was adopted to fit with the practices in the offshore installations.

Hazards Likely to Occur in Offshore Environment: this measures their perception of factors that can cause hazards in the offshore environment. 15 questions were asked to elucidate their response as adopted from the offshore industry safety questionnaire [17]. An example of such a question was; how likely is wind/strong current to occur in an offshore environment?

Injury Types Experienced by Offshore Workers: This measures their opinion on the common injuries experienced in an offshore environment. 10 investigative variables were used to find out their response as embraced from the offshore safety questionnaire [17]. A sample of the question was; if laceration was probable to appear offshores. "Others" were used in the questionnaire to represent Musculoskeletal Disorders.

Injured Body parts prevalent among offshore Workers: This measures their Opinion on the most affected Body parts when an Injury Occurs Offshores. 15 body parts were identified from the offshore safety questionnaire [17]. Questions on if the hand was a prevalent body part to get injured offshores were asked.

Types of Accidents in Offshore Environment: Respondents were asked to carefully choose accidents prevalent in an offshore environment. 8 investigative variables were used to find out their response as embraced from the offshore safety questionnaire [17]. Industrial chemical accident was asked to know their perception of the likelihood of such accident to occur.

Consequences of Accidents in Offshore Environment; Interviewees were asked about their perception on the effect of accidents offshores. . 8 investigative variables were used to find out their response as embraced from the offshore safety questionnaire [17]. Effect on the personnel was one of the questions asked by the researcher to find out the response on the likelihood of the consequence to happen.

2.2 Data Analysis and Procedures

The responses from the participants were analyzed using SPSS version 26 and the coding was done according to the Likert scale used for the questionnaire. Composite scores were computed for each of the constructs and reliability was carried out using Cronbach alpha. Descriptive statistics were done on the composite score to understand the general view of the participants on each of the work-related injuries. Transformations were carried out to meet assumptions for Parametric Statistics and compare values between Parametric and non-parametric tests to see if there would be any significant difference. Mann-Whitney Test and z test were carried out to establish any significant difference in work-related injuries for different genders. Kruskal Wallis test and ANOVA test were carried out to establish any significant difference in work-related injuries and educational qualification. Mann-Whitney and z-Test were carried out to establish any Significant differences in the work-related injuries and type of staff. Kruskal Wallis test and ANOVA test were carried out to establish any significant difference in work-related injuries and years of working experience.

3. RESULTS

A total of 450 questionnaires were distributed among workers that work in the oil and gas sector in the Niger Delta region using an online questionnaire survey method. From the 450 questionnaires distributed, a total of 70 questionnaires were not returned which accounted for about 15.55% of the total questionnaires. A total of 54 questionnaires were not properly filled which accounted for about 12% of the total questionnaires distributed and 325 questionnaires were deemed ok to be used which accounted for about 72.44% of the total questionnaires.

Table 1 shows the distribution of demographic variables for Gender, Educational Qualification, work experience and types of Staff.

Table 2 shows the Test for Significant differences in work-related injuries for gender using the

Mann-Whitney test and Z test after transformation. The significance test on injuries using the Mann-Whitney test for Significance and z-test had values greater than p –value greater than 0.05.p Which shows that there is no significance relationship between work- related injuries and Gender.

Kruskal Wallis test and ANOVA test for Significant differences in the work-related injuries for educational qualification using Kruskal- Wallis non-Parametric Statistics and transformation was done so that Parametric Statistics can be used for analysis as shown in Table 3. Contusion was significant using both Kruskal-Wallis and ANOVA and “Others” which represents musculoskeletal disorder was significant using nonparametric statistics and not Significant using Parametric Statistics. Which implies that Education has a role to play in musculoskeletal injuries using the Kruskal-Wallis test of Significance.

Table 1. Distribution of demographic variables

Demographic criteria	Gender	Frequency	Percentage (%)	Cumulative percentage (%)
Gender	Male	225	69.2	69.2
	Female	100	30.8	100
Educational Qualification	Primary	36	11.1	11.1
	Secondary	39	12.0	23.1
	Tertiary	250	76.9	100
Marital Status	Single	156	48.0	48.0
	Married	162	49.8	97.8
	Divorced	7	2.2	100
Work Experience (Years)	0 - 5	101	31.1	31.1
	6 - 10	79	24.3	55.4
	11 - 15	86	26.5	81.8
	16 - 20	50	15.4	97.2
	21 and Above	9	2.8	100.0
Type of Staff	Contract	98	30.2	30.2
	Permanent	227	69.8	100

Table 2. Mann-Whitney and Z-Test for Significant difference in the work-related injuries and different gender

Variable\Test	Mann-Whitney	Z-test
Amputation	0.338	0.331
Contusion	0.323	0.262
Burn	0.463	0.411
Laceration	0.369	0.457
Fracture	0.777	0.990
Dislocation	0.750	0.755
Strains and Sprains	0.472	0.447
Bruises	0.787	0.863
Cut	0.938	0.906
Others	0.185	0.206

Table 3. Kruskal Wallis and ANOVA tests for Significant difference in the work-related injuries and educational qualification

Variable\Test	Kruskal-Wallis	ANOVA
Amputation	0.435	0.401
Contusion	0.013	0.015
Burn	0.130	0.118
Laceration	0.176	0.144
Fracture	0.838	0.783
Dislocation	0.439	0.315
Strains and Sprains	0.365	0.324
Bruises	0.761	0.654
Cuts	0.125	0.078
Others	0.046	0.068

Table 4. Dunn multiple comparison test for contusion injuries

Sample	Frequency	Mean of ranks	Groups
I2 Tertiary	250	155.010	A
I2 Primary	36	185.583	A B
I2 Secondary	39	193.372	B

Category	LS means	Groups
Secondary	2.846	A
Primary	2.778	A B
Tertiary	2.448	B

Table 5. Dunn multiple comparison test for “other” injuries

Sample	Frequency	Mean of ranks	Groups
I10 Tertiary	250	157.180	A
I10 Secondary	39	170.654	A B
I10 Primary	36	195.125	B

Table 6. Mann-Whitney and z-test for Significant difference in the work-related injuries and type of staff

Variable\Test	Mann-Whitney	z-test
Amputation	0.080	0.070
Contusion	0.276	0.265
Burn	0.947	0.891
Laceration	0.798	0.842
Fracture	0.568	0.518
Dislocation	0.059	0.057
Strains and Sprains	0.435	0.482
Bruises	0.243	0.268
Cut	0.338	0.364
Others	0.005	0.006

From Table 4, Dunn’s test is used for testing two or more experimental groups. Dunn’s multiple comparison tests for contusion injuries using the mean of ranks showed that Tertiary degrees had a mean ranking of 155.010 which shows workers with Tertiary Education experience fewer injuries than Primary and secondary school holders

which had a mean ranking of 185.583 and 193.372 respectively.

From Table 5: Dunn multiple comparison tests for “other” injuries. (“Others” refers to musculoskeletal Diseases. Musculoskeletal disorders (MSDs) are conditions that can affect

your muscles, bones, and joints. MSDs include tendinitis, carpal tunnel syndrome, osteoarthritis, rheumatoid arthritis (RA), and fibromyalgia). Tertiary degrees had a mean ranking of 157.108 which shows workers with Tertiary Education experience fewer injuries than Secondary and primary school holders which had a mean ranking of 170.654 and 195.125 respectively.

Table 6 shows Mann-Whitney and Z-Test for Significant differences in the work-related injuries for the type of staff. Mann-Whitney, non-Parametric Statistics and transformation were done so that Z-test can be used for analysis. The significance test on work-related injuries for the type of Staff using the Mann-Whitney test for Significance has a p –the value of 0.05 which dhowes that Musculoskeletal Disorders was a significant type of injury experienced by staff.

Table 7 shows Kruskal Wallis and ANOVA test for Significant differences in work-related injuries for years of working experience. Amputation and Fracture were statistically significant having a p-value of less than 0.05.

Table 8 shows the Dunn multiple comparison tests for amputation injuries using the mean of ranks. Greater than 20-year work experience had a mean rank of 100.444, and 11 to 15 years of work experience had a mean rank of 151.988.personnel with 6 to 10 years of working experience had a mean rank of 167.178. Personnel with 0 to 5 years of working experience had a mean ranking of 168.178 and personnel with 16-20years of work experience had the highest mean rank of 176.300.

Table 9 shows the Dunn multiple comparison test for Fracture injuries using the mean of ranks. Greater than 20-year work experience had a mean rank of 196.944, and personnel with 11 to 15 years of work experience had a mean rank of 196.944. Personnel with 6 to 10 years of working experience had a mean rank of 151.234. Personnel with 0 to 5 years of work experience had a mean ranking of 158.886 and personnel with 16-20 years of work experience had a mean rank of 1144.200.

Table 7. Kruskal-Wallis and ANOVA tests for Significant difference in the work-related injuries and year of working experience

Variable\Test	Kruskal-Wallis	ANOVA
Amputation	0.031	0.035
Contusion	0.125	0.128
Burn	0.433	0.437
Laceration	0.696	0.606
Fracture	0.020	0.022
Dislocation	0.663	0.495
Strains and Sprains	0.803	0.806
Bruises	0.237	0.160
Cut	0.416	0.265
Others	0.659	0.738

Table 8. Dunn multiple comparison test for amputation injuries

Sample	Frequency	Mean of ranks	Groups
I2 >20 years	9	100.444	A
I2 11-15 years	86	151.988	A B
I2 6-10 years	79	167.076	B
I2 0-5 years	101	168.178	B
I2 16-20 years	50	176.300	B

Table 9. Dunn multiple comparison test for fracture injuries

Sample	Frequency	Mean of ranks	Groups
I5 16-20 years	50	144.700	A
I5 6-10 years	79	151.234	A
I5 0-5 years	101	158.886	A
I5 11-15 years	86	185.727	B
I5 >20 years	9	196.944	B

4. DISCUSSION

The result shows that certain demographic criteria influence work-related accidents and certain kinds of intervention should be introduced to ensure that all working on offshore platforms are safe. For this study, most Parametric statistics were as significant as non-parametric statistic. There is no significant relationship between work-related injuries and Gender. From our findings, gender does not play any significant role in work-related injury experienced offshores. This is so as they have both been exposed to same training on causes of accident and how to prevent accidents in the work place [18]. This is in tandem with the findings of [19] who agreed on occupational injuries among students were not gender-based.

There is no significant relationship between work-related injuries and Education, from the study, education has a significant role to play in contusion and Musculoskeletal Disorders and work-related injuries. Dunn's multiple comparison tests have shown that Musculoskeletal Disorders' work-related injuries decrease with an increase in Educational qualification findings [20,21] also agreed with these findings that Education has a role to play in the cause of injuries. However, Contusion Injuries were more on workers with Secondary school education, this is so because these were the class of personnel that interface with the machinery.

There is no significant relationship between work-related injuries and types of Staff. Musculoskeletal Disorders were a significant type of injury experienced by staff. This happens as a result of awkward positions that workers have to attain while carrying out their job. According to [22], workers from the oil and gas industry are largely uncovered to ergonomic-related injuries and injury risks such as repeated bending, lifting heavy items, pushing and pulling weight loads, reaching overhead, performing the same or similar tasks repetitively, and working in awkward body postures. As a result, managers in the oil and gas business have frequently received worker complaints such as chronic fatigues, back pains, headaches, upper body pains, sleep disorders, and stresses [23]. These complaints are indications of ergonomic deficiencies in the work system from the industry.

There is no significant relationship between work-related injuries and years of Experience. From the work, years of Experience were

significant for Amputation and Fracture [24]. Believes that years of experience have a role to play, however, their study attributes work-related injury to lower levels of years of experience between 0-5 years. Dunn's Test of Multiple comparisons has shown that Amputation was common amongst workers that have worked for 16-20 years and this could largely be associated with complacency. While workers who have spent more than 20 years on the job experienced more fractures which are closely associated with the age of the personnel.

5. CONCLUSION

In Summary, work-related Injuries about demographic factors were examined. Gender does not play any significant role in work-related injuries, but Education plays a role in certain types of injuries, musculoskeletal disorders were the common types of injuries experienced by staff, and fracture and Amputations were experienced by workers who have spent above 16 years working offshore. More study on this research should cover more offshore locations including the newly discovered oil wells in Northern Nigeria. One of the main limitation of this study was that we were not granted access to physically visit those offshore location as a result of Covid-19 restrictions.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mekkodathil A, El-Menyar A, Al-Thani H. Occupational injuries in workers from different ethnicities. *International Journal of Critical Illness and Injury Science*. 2016;6(1):25–32. DOI:<https://doi.org/10.4103/2229-5151.177365>
2. National General Certificate Unit. NEBOSH Certificate Courses - Sample Material, national examination board in occupational safety and health. national examination board in occupational safety and health, London, United Kingdom; 2015.
3. El Bouti MY, Allouch M. Analysis of 801 work-related incidents in the oil and gas industry that occurred between 2014 and

- 2016 in 6 Regions. Energy and Environment Research. 2018;8(1): 32-47.
4. Ferreira DF, Suslick SB. Identifying potential impacts of bonding instruments on offshore oil projects. Resources Policy. 2001;27(1):43-52.
 5. Department of Homeland Security. Energy Sector-Specific Plan; 2015.
 6. Higgins BKL, Vernadsky V. Economic growth and sustainability – are they mutually exclusive?; 2013.
 7. DOSH. Annual Safety and Health Report; 2014.
Available:<http://www.dosh.gov.my/index.php/en/archive-statistics/2014>
 8. Leigh JP, Waehrer G, Miller TR, Keenan C. Costs of occupational injury and illness across industries. Scandinavian Journal of Work, Environment & Health. 2004;30(3): 199–205.
Available:<http://www.jstor.org/stable/40968777>
 9. Rommel A, Varnaccia G, Lahmann N, Kottner J, Kroll LE. Occupational injuries in Germany: population-wide national survey data emphasize the importance of work-related factors. PLoS ONE. 2016;11(2): e0148798.
DOI:10.1371/journal.pone.0148798
 10. Asad MM, Hassan RB, Sherwani F, Ibrahim NH, Soomro QM. Level of satisfaction for occupational safety and health training activities: a broad spectrum industrial survey. In Journal of Physics: Conference Series. 2018b;1049(1): 012021.
 11. HSE. Key Programme 3 – Asset Integrity Programme. Health and Safety Executive, United; 2008.
 12. Ilo S. Health at work: a vision for sustainable prevention: XX World Congress on Safety and Health at Work 2014: Global Forum for Prevention, 24–27 August 2014, Frankfurt, Germany. International Labour Office; 2014.
 13. Nolan DP. Handbook of fire and explosion protection engineering principles: for oil, gas, chemical and related facilities. William Andrew; 2014.
 14. Asad MM, Hassan RB, Ibrahim NH, Sherwani F, Soomro QM. Indication of decision making through accident prevention resources among drilling crew at oil and gas industries: a quantitative survey. In Journal of Physics: Conference Series. 2018a;1049(1):012021.
 15. Xiuwen Sue Dong, Xuanwen Wang, Julie A. Largay. Occupational and non-occupational factors associated with work-related injuries among construction workers in the USA, International Journal of Occupational and Environmental Health. 2015;21:2:142-150.
DOI: 10.1179/2049396714Y.0000000107
 16. Bhardwaj U, Teixeira AP, Guedes Soares C. Analysis of FPSO Accident and Incident data; 2017.
Available:<https://www.researchgate.net/publication/317074849>
 17. HSE. HSE and workplace transport; 2005.
Available:<http://www.hse.gov.uk/workplace-transport/hsewpt.htm>.
 18. Zakaria NH, Mansor N, Abdullah Z. Workplace accident in Malaysia: most common causes and solutions. Business and Management Review. 2012;2(5):75-88.
 19. Shipp EM. et al. Substance use and occupational injuries among high school students in South Texas, UTMB Health Research Expert Profiles. Informa Healthcare; 1970.
Available:<https://researchexperts.utmb.edu/en/publications/substance-use-and-occupational-injuries-among-high-school-student>
(Accessed: January 25, 2023).
 20. Lombardi S, et al. (no date) Independent effects of sleep duration and body mass index on the risk of a work-related injury: Evidence from the US National Health Interview Survey (2004-2010), Chronobiology international. U.S. National Library of Medicine.
Available:<https://pubmed.ncbi.nlm.nih.gov/22621351/>
(Accessed: January 25, 2023).
 21. Alkhalidi M, Pathirage C, Kulatunga U. The role of human error in accidents within oil and gas industry in Bahrain. In 13th International Postgraduate Research Conference (IPGRC): conference proceedings. University of Salford. 2017;822-834.
 22. Occupational Safety and Health Administration. Safety Hazards Associated with Oil and Gas Extraction Activities. United States Department of Labor, Washington, DC 2010; 2016.
 23. Waage S, Moen BE, Pallesen S, Eriksen HR, Ursin H, Åkerstedt T, Bjorvatn B. Shift

- work disorder among oil rig workers in the North Sea. Sleep. 2009;32(4):558-565.
24. Bena A, et al. (no date) Job tenure and work injuries: A multivariate analysis of the relation with previous experience and differences by age, BMC public health. U.S. National Library of Medicine. Available:<https://pubmed.ncbi.nlm.nih.gov/24053157/> (Accessed: January 25, 2023).

© 2023 Oyamienlen et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96301>