

Dysmenorrhea and Associated Risk Factors among Adolescent Girls in Junior High School of Upper East Region, Ghana

**Florence Assibi Ziba¹, Yussif Adams^{2*}, Peter Paul Mwinsanga Dapare²,
Emmanuel Yinkela-Onni Nanao¹, Mavis Agaanamah Ayalbire¹
and Elizabeth Memuna Baba¹**

¹Department of Nursing, School of Allied Health Sciences, University for Development Studies,
Ghana.

²Department of Biomedical Laboratory Sciences, School of Allied Health Sciences, University for
Development Studies, Ghana.

Authors' contributions

This work was carried out in collaboration among all authors. Authors FAZ, YA, PPMD, EYN, MAA and EMB designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors FAZ, YA and PPMD managed the analysis of the study. Authors EYN, MAA and EMB managed the literature searches. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Abdelmonem Awad M. Hegazy, Professor, Department of Anatomy and Embryology, Faculty of Medicine, Zagazig University, Egypt.

Reviewers:

- (1) Ronald Bartzatt, University of Nebraska, USA.
(2) Shigeki Matsubara Jichi, Tokyo Medical University, Japan.
(3) Alka Ahuja, Post Graduate Institute of Medical, Education and Research, India.
Complete Peer review History: <http://www.sdiarticle3.com/review-history/49911>

Original Research Article

**Received 21 April 2019
Accepted 25 June 2019
Published 02 July 2019**

ABSTRACT

Aim: The aim of the study was to determine the prevalence of dysmenorrhea, associated risk factors, and how it is managed among adolescent girls in junior high schools of the Upper East region in Ghana.

Study Design: A descriptive cross-sectional study design was adopted.

Place and Duration of Study: The study was carried out in five (5) districts; Bolgatanga, Bongo,

Builsa, Talensi and Nabdam in the Upper East region of Ghana, between February to June 2018.

Methodology: Simple random sampling was used to collect data from 400 participants in 14 junior high schools of the five districts using a self-administered questionnaire. Likert Scale (LS) was used to rate the severity of dysmenorrhea. Data were analyzed and p-value < 0.05 considered statistically significant.

Results: The prevalence of dysmenorrhea was 85%. About 77.8% of respondents with dysmenorrhea had their first menstruation between the ages of 9-11 years. Among those who had dysmenorrhea, 46.8% experienced mild menstrual pain while 11.0% experienced severe pain. Factors that were identified as predictors of risk factors for dysmenorrhea were; menarche between the age 9-11years (aOR = 1.92, 95% CI = 1.053-3.495), short (<21 days) menstrual cyclic length (aOR = 1.98, 95% CI = 1.021-3.578) and short (<2 days) menstruation days (aOR = 2.55, 95% CI = 1.385-4.617).

Conclusion: Dysmenorrhea prevalence is high and the risk factors are; girls who had first menstruation between the ages of 9-11 years, those who experience short (<21 days) menstrual cyclic length and short (<2 days) menstrual days. The study recommends that; reproductive health should be included in school health education programs early enough and education extended to parents in order to address the reproductive health needs of females. Further work, however, is encouraged to validate the reliability of these risk factors of dysmenorrhea.

Keywords: Dysmenorrhea; adolescent; gynaecology; junior high school; Ghana.

ABBREVIATIONS

Dysmenorrhea was defined as painful menstruation experienced by an adolescent within the past 6 months.

Likert Scale (LS) was adopted to rate the severity of dysmenorrhea using the Cronbach's α coefficient ($\alpha = 0.94$) (Mcdowell, 1996). A zero (0) score on the scale represented no pain at all while ten (10) represented severe/unbearable pain. Participants were required to rate the degree of pain by ticking the number. The following were the classification of score received; mild dysmenorrhea: 1-3, moderate dysmenorrhea: 4-7 and severe dysmenorrhea: 8-10.

Regular physical activity was measured as individual involving in outdoor physical activity for 30 minutes or longer each day for at least 4 times a week.

1. INTRODUCTION

Menstruation is a common physiological change experienced by adolescent girls of which the first of its kind is called menarche. This monthly period is often associated with problems of irregular menstrual flow, profuse bleeding and dysmenorrhea. Dysmenorrhea is a painful or cramping feeling experienced in the lower abdomen during menstruation with headache, nausea, dizziness, vomiting, diarrhoea, breast tenderness, backache and leg pain being the signs and symptoms [1-3].

Dysmenorrhea is the most common gynecologic complaint among adolescent girls [4,5] and it can be primary or secondary. Primary dysmenorrhea is pain experienced during menstruation with normal pelvic anatomy while secondary dysmenorrhea is associated with pelvic pathology with common causes reported being a pelvic inflammatory disease, endometriosis and ovarian cysts [6].

High dysmenorrhea prevalence has been reported in some studies as 83.6% in Tamale [7] and 74.4% in Accra [8]. In other studies outside Ghana, a prevalence of 66.2% in Nigeria [9], 78.35% in Benin [10], 80.1% in Egypt [11], 76% in Manipur [12], 79.7% in Gwalior [13] and 84% in India [14] were found. However, 25% to 50% dysmenorrhea prevalence was reported by Harlow and Campbell [15]. Risk factors associated with dysmenorrhea reported by several studies include; excessive bleeding, short menstrual cyclic length, family history of dysmenorrhea, depression and smoking [10,12, 16-18].

Most females with dysmenorrhea experienced some level of pain and distress and this affects their daily activities including work schedules. In the United States, it was reported that dysmenorrhea is the leading cause of periodic short-term school absenteeism among adolescent girls. About 70% of adolescent girls reported regular school absenteeism in

Tbilisi (Georgia) due to the severity of menstrual pain [19]. Girls who experience painful menstruation had poor interpersonal relationships with family and friends, reduced daily physical activity and poor examination grades due to decreased class concentration [12, 20,21].

In spite of the effects of dysmenorrhea on adolescent girls, few seek medical treatment [10, 20,22]. Most girls and their parents, especially mothers, accept dysmenorrhea as a normal feature of the menstrual cycle due to lack of adequate information and the cultural myth (silence) that is associated with it. The pain and trauma related to menstruation go unexpressed and the adolescent girls suffer as a result. Previous studies conducted on dysmenorrhea among adolescent girls were carried out in Accra, the southern part of Ghana. The study, therefore, aimed at finding the prevalence of dysmenorrhea and associated risk factors, information on awareness and how it is managed among adolescent girls in Junior High Schools of the Upper East region, the northern part of Ghana.

2. MATERIALS AND METHODS

2.1 Study Area

The study was carried out in five (5) districts; Bolgatanga (number of participants = 152), Bongo (number of participants = 100), Builsa (number of participants = 49), Talensi (number of participants = 71) and Nabdam (number of participants = 28) in the Upper East region of Ghana, between February to June 2018. The Upper East region is one of the 16 regions of Ghana located in the northern part with its regional capital being Bolgatanga. This region houses diverse ethnic populations who speak different dialects and practice the three (3) main religions: Christianity, Traditional, and Islam.

2.2 Study Design

The study was a descriptive cross-sectional study among adolescent girls in Junior High School in the Upper East region of Ghana. The schools involved were; Fountain Gate International School, Kumbosco, Golden Step, Nadengzieg in Bolgatanga; Beo Roman Catholic (R/C), Sapporo, St. Luke in Bongo; Ayieta, Success International School in Builsa; Sakorit, Kpatia, Kolpellega in Talensi; Dasabligo and Kongo in Nabdam district.

2.3 Study Population

The target populations were girls between the ages of 11 - 19 years. Girls who had not seen their menstrual period for the last 6 months, refused to consent/take part in the study and those who were absent during our visit were all excluded from the study.

2.4 Sample Size Calculation

The sample size was calculated using the Cochran [23] formula. The sample size was, therefore, calculated from the expression:

$$n = \frac{Z^2(1-p)p}{e^2}$$

Where

n = Minimum sample size.

Z = Abscissa off the normal curve that cuts off an area α at the tails ($1 - \alpha$ equals the desired confidence level, ie 95%).

e = Margin of error at 5% (standard value of 0.05).

p = Prevalence of dysmenorrhea among adolescent in Accra, Ghana is 74.4% [8].

Using the information in the formula above, the sample size for this study was 293 persons. Given a targeted response rate of 90%, the sample size was recalculated as $293/0.90$. Using the above formula, the calculated sample size was approximately 326. However, 400 girls were targeted and recruited to participate in the study.

2.5 Sampling Technique

A simple random sampling technique of balloting was used to select fourteen (14) schools out of 43 junior high schools in the region. In each school, questionnaires were randomly distributed to participants who met the sampling criteria.

2.6 Data Collection

2.6.1 Pre-testing of questionnaires

In order to clear all uncertainties and any ambiguity, 10 questionnaires were pre-tested among adolescent girls at Bolgatanga Roman Catholic (R/C) junior high school in order to change the instructions that were not clear to the participants.

2.6.2 Administering of questionnaires

Data was collected using a self-administered questionnaire. The questionnaires were

developed to cover socio-demographic characteristics, history of menstruation, the severity of dysmenorrhea, physical activity levels, treatment-seeking practice and quality of life-related to dysmenorrhea such as mood, emotion, concentration and school attendance.

Dysmenorrhea was defined as painful menstruation experienced by an adolescent within the past 6 months. Likert Scale (LS) was adopted to rate the severity of dysmenorrhea using Cronbach's α coefficient ($\alpha = 0.94$) (Mcdowell, 1996). A zero (0) score on the scale represented no pain at all while ten (10) represented severe / unbearable pain. Participants were required to rate the degree of pain by ticking the number. The following were the classification of score received; mild dysmenorrhea: 1-3, moderate dysmenorrhea: 4-7 and severe dysmenorrhea: 8-10.

Regular physical activity was measured as an individual involving in outdoor physical activity for 30 minutes or longer each day for at least 4 times a week.

2.7 Statistical Analysis

Data were analyzed using Microsoft Excel 2013 and GraphPad Prism version 6.0 (www.graphpad.com). Data were presented as frequency, percentages and charts. Categorical variables were compared using the chi-square test. Bivariate and multivariate logistic regression analysis was used to assess risk factors associated with dysmenorrhea and in all cases a p-value < 0.05 was considered statistically significant.

3. RESULTS

3.1 Socio-demographic Characteristics of Study Participants

A total of 400 adolescent girls participated in the study. Majority of the respondents (63.0%) were in 14-16years age category compared with other age categories, while the age category 17-19 years had the least number of respondents (12.5%). The majority (79.5%) of respondents was Christians, in JHS 1 (35.5%) and about 38.0% reside in Bolgatanga in the Upper East Region of Ghana (Table 1).

3.2 Logistic Regression Analysis of Some Selected Variables and Dysmenorrhea Outcome

Table 2 shows univariate and multivariate logistic regression analysis of some selected variables and dysmenorrhea outcome. Menarche at age of 9-11years (OR = 1.93, 95% CI = 1.061-3.506), short (<21 days) menstrual cycle length (OR = 1.88, 95% CI = 1.017-3.490) and short (< 2 days) menstruation days (OR = 2.51, 95% CI = 1.380-4.576) were found to be risk factors for associated with dysmenorrhea among JHS girls in Upper East region.

After adjusting for confounding variables, girls (77.8%) who had menarche at age 9-11years were 2 times (aOR = 1.92, 95% CI = 1.053-3.495) more likely to experience dysmenorrhea than those who had menarche between the age 12-19years. Respondents who experienced short (<21 days) menstrual cyclic length were about 2 times (aOR = 1.98, 95% CI = 1.021-3.578) more likely to experience dysmenorrhea than those who experienced normal (22-28 days) and long (>28 days) menstrual cycle length. Furthermore, respondents whose menstruation came for 2 days or less (short menstruation days) were 3 times (aOR = 2.55, 95% CI = 1.385-4.617) more likely to experience dysmenorrhea than those with normal (2-6 days) and long (> 6 days) menstrual days (Table 2).

3.3 Distribution of Dysmenorrheic by the Severity of Dysmenorrheal

The prevalence of dysmenorrhea was 85%. Among those who had dysmenorrhea, 46.8% experienced mild menstrual pain while 11.0% experienced severe pain (Fig.1).

3.4 Percentage Distribution of Dysmenorrhea Severity and Quality of Life

Majority of the study participants with dysmenorrhea responded that interpersonal relationship with family and friends (75.9%) were affected due to painful menstruation. A total of 58.5% were depressed, 28.8% emotionally unstable and 14.7% absenting themselves from school during dysmenorrhea state. As the degree of painful menstruation increases from mild to severe, these variables (relationship with family and friends, daily physical activity, concentration during class hours, depressed mood and emotional instability) increase.

Table 1. Socio-demographic characteristics of study participants

Variable	Frequency (n = 400)	Per cent (%)
Age (years)		
11-13	98	24.5
14-16	252	63.0
17-19	50	12.5
Educational level		
JHS/Form 1	142	35.5
JHS/Form 2	135	33.7
JHS/Form 3	123	30.8
Religion		
Christian	318	79.5
Islam	77	19.3
Traditional	5	1.2
District		
Bolga	152	38.0
Bongo	100	25.0
Builsa	49	12.3
Nabdam	28	7.0
Talensi	71	17.7

Data presented as frequency and percent

About one-seventh (14.7%) of girls with dysmenorrhea had been absent from school for at least 1 day due to painful menstruation in the past 6 months and absenteeism increases with increase in severity of dysmenorrhea (Table 3).

3.5 Information on Awareness

The prevalence of dysmenorrhea awareness was 92.0%. Among those who were aware of dysmenorrhea, 85.6% had experienced dysmenorrhea at least once in their lifetime while 14.4% had not experienced dysmenorrhea (Fig. 2).

3.6 Information on Awareness Stratified by Dysmenorrhea Status

Out of the 400 participants studied, 45.8% had heard of dysmenorrhea years after menarche, of which those who had experienced dysmenorrhea were the majority (82.0%). Among those who heard of dysmenorrhea before menarche, 87.5% of the girls had experienced dysmenorrhea while 12.5% had not.

Majority of this information was from the family members (39.5%), teachers at school (20.5%), and friends (18.8%) with the least from the media (3.3%) (Table 4).

About 75.5% of the participants felt treatment was necessary for painful menstruation of which those who had experienced dysmenorrhea seek to relax (100%), take herbs (100%) or over counter drugs (100%) as a form of medication, or lower fat and sugar intake (100%) as a remedy.

Remedies were prescribed either by self (30.8%), mother (3.8%) or a doctor (10.5%) with 70.5% of the total participants studied of the opinion that, these remedies were effective in the management of painful menstruation (Table 4).

4. DISCUSSION

The study found a prevalence of dysmenorrhea to be 85%. This high prevalence is consistent with several other studies which found a dysmenorrhea prevalence of 83.6% among female students in Tamale, Ghana [7], 74.4% in adolescent girls in Accra, Ghana [8], 78.35% in the school of Parakou in Benin [10], 76% among high secondary schoolgirls of Imphal West district, Manipur [12], 79.7% among higher secondary schoolgirls in Gwalior [13] and 84% in college going girls in India [14]. However, a prevalence of 13.7% by Ogunfowokan and Babatunde [24] in Nigeria and 25% to 50% in a Cochrane systematic review of studies in developing countries by Harlow and Campbell [15] indicate much lower prevalence. These variations of prevalence may be due to sociocultural and ethnic factors since this study was carried out in different ecological zones. Another reason for variations of prevalence may be due to different diagnostic techniques/tools, the absence of a widely accepted definition of dysmenorrhea and a system for grading dysmenorrhea severity [25]. In this study, the majority (84.1%) within the 14-16years age category had experienced dysmenorrhea. It was also revealed that 46.8%, 27.3% and 11.0% of girls had experienced mild, moderate and severe menstrual pain respectively. Similar studies by Kural, Noor, Pandit et al., [14] found 29.2%, 36.6% and 34.2% and Kumar, Konjengbam and Devi [12] reported 32.8%, 46.0% and 21.2% for mild, moderate and severe pain respectively. In other studies, severe pain was reported to be 6.3% [26], 20.1% [27], 28.8% [10] and 42% [28] of adolescent girls experiencing dysmenorrhea. The observed differences may be due to a technique in grading the severity of pain. Likert scale (LS) was adopted in this study to rate the severity of dysmenorrhea using the Cronbach's α coefficient ($\alpha = 0.94$) [29] but other methods for assessing menstrual pain may include visual-

Table 2. Univariate and multivariate logistic regression analysis of some selected variables and dysmenorrhea outcome

Variable	Univariate analysis				Multivariate analysis	
	Dysmenorrhea, n(%)	No Dysmenorrhea, n(%)	OR (95% CI)	p-value	aOR (95% CI)	p-value
Age (years)						
11-13	83 (84.7)	15 (15.3)	1.03 (0.547-1.947)	.92	NA	NA
14-16	212 (84.1)	40 (15.9)	1.0			
17-19	45 (90.0)	5 (10.0)	0.60 (0.226-1.569)	.29		
Age of menarche (years)						
9-11	70 (77.8)	20 (22.2)	1.93 (1.061-3.506)	.03	1.92 (1.053-3.495)	.03
12-14	243 (86.2)	39 (13.8)	1.0		1.0	
15-17	27 (96.4)	1 (3.6)	0.20 (0.026-1.475)	.08	0.19 (0.031-1.528)	.09
Regular physical activity						
Yes	42 (93.3)	3 (6.7)	0.37 (0.112-1.247)	.10	0.37 (0.108-1.239)	.09
No	298 (83.9)	57 (16.1)	1.0		1.0	
Menstrual cycle length						
< 21 days (short)	63 (77.8)	18 (22.2)	1.88 (1.017-3.490)	.04	1.98 (1.021-3.578)	.04
22-28 days (normal)	246 (86.9)	37 (13.1)	1.0		1.0	
>28 days (long)	31 (86.1)	5 (13.9)	0.91 (0.338-2.432)	.85	1.05 (0.432-2.497)	.84
Menstruation days						
< 2 days (short)	60 (74.1)	21 (25.9)	2.51 (1.380-4.576)	.002	2.55 (1.385-4.617)	.002
2-6 days (normal)	239 (87.2)	35 (12.8)	1.0		1.0	
> 6 days (long)	41 (91.1)	4 (8.9)	0.52 (0.179-1.512)	.22	0.57 (0.175-1.524)	.23
Educational level						
JHS 1/Form 1	125 (88.0)	17 (12.0)	0.68 (0.372-1.243)	.21	0.68 (0.373-1.241)	.22
JHS 2/Form 2	116 (85.9)	19 (14.1)	0.90 (0.497-1.612)	.71	0.89 (0.485-1.608)	.71
JHS 3/Form 3	99 (80.5)	24 (19.5)	1.0		1.0	
Family history						
Yes	36 (90.0)	4 (10.0)	0.60 (0.207-1.762)	.35	0.60 (0.213-1.768)	.35
No	63 (85.1)	11 (14.9)	0.99 (0.486-2.006)	.97	0.99 (0.487-2.015)	.97
Don't know	241 (84.3)	45 (15.7)	1.0		1.0	

Data presented as number and percentages. OR: odd ratio, aOR: adjusted odd ratio, 95% CI: 95% confidence interval and p-value <0.05 considered statistically significant




Fig. 1. Distribution of dysmenorrheic by the severity of dysmenorrheal

Table 3. Percentage distribution of dysmenorrhea severity and quality of life

Variable	Total, n(%)	Mild, n(%)	Moderate, n(%)	Severe, n(%)
Interpersonal relationship				
Affected	258 (75.9)	137 (73.3)	85 (78.0)	36 (81.8)
Not affected	82 (24.1)	50 (26.7)	24 (22.0)	8 (18.2)
Depressed mode				
Affected	199 (58.5)	109 (58.3)	64 (58.7)	26 (59.1)
Not affected	141 (41.5)	78 (41.7)	45 (41.3)	18 (40.9)
Daily physical activity				
Affected	96 (28.2)	46 (24.6)	34 (31.2)	16 (36.4)
Not affected	117 (34.4)	141 (75.4)	75 (68.8)	28 (63.6)
Reduced concentration				
Affected	111 (32.6)	61 (32.6)	38 (34.9)	16 (36.4)
Not affected	229 (67.4)	126 (67.4)	71 (65.1)	28 (63.6)
School absenteeism				
Yes	50 (14.7)	27 (14.4)	16 (14.7)	7 (15.9)
No	290 (85.3)	160 (85.6)	93 (85.3)	37 (84.1)
Emotional instability				
Yes	98 (28.8)	50 (26.7)	31 (28.4)	17 (38.6)
No	242 (71.2)	137 (73.3)	78 (71.6)	27 (61.4)

Data presented as frequency (percent)

analogue and numeric scales (VAS) [30]. Furthermore, the differences in pain severity may also be due to individual differences with regards to pain perception and their threshold [10]. Pain is an immensely subjective symptom and grading the severity of dysmenorrhea had been very difficult.

The findings also showed that, as the degree of menstrual pain increases from mild, moderate to severe, interpersonal relationship with family and friends, emotional instability, reduced daily physical activity and concentration during class

hours were all affected. These findings are in accordance with other findings [12,20,21,26] which found that individuals who experience painful menstruation had poor interpersonal relationships with family and friends, reduced daily physical activity, and poor examination grades due to decreased class concentration.

A previous study in Ghana reported 9.2% of young girls absent from school [8]. About one-seventh (14.7%) of girls with dysmenorrhea in this study had been absent from school for at least 1 day due to painful menstruation in the

Table 4. Information on awareness stratified by dysmenorrhea status

Variable	Dysmenorrhea, n(%)	No dysmenorrhea, n(%)
At what point did you hear of dysmenorrhea?		
Before Menarche	28 (87.5)	4 (12.5)
Immediately after Menarche	67 (90.5)	7 (9.5)
Months after Menarche	70 (88.6)	9 (11.4)
Years after Menarche	150 (82.0)	33 (18.0)
Information source		
Family (mother)	131 (82.9)	27 (17.1)
Friends	65 (86.7)	10 (13.3)
Teacher	75 (91.5)	7 (8.5)
Media	9 (69.2)	4 (30.8)
Textbooks	35 (87.5)	5 (12.5)
Remedies		
Exercise	38 (92.7)	3 (7.3)
Warm compress	36 (97.3)	1 (2.7)
Over the counter drugs	61 (100.0)	0 (0.0)
Visiting a doctor	51 (94.4)	3 (5.6)
Herbs	32 (100.0)	0 (0.0)
Relaxation/rest	108 (100.0)	0 (0.0)
Lower fat and sugar foods	95 (100.0)	0 (0.0)
None	54 (96.4)	2 (3.6)
Who prescribes the remedies for you?		
Self	120 (97.6)	3 (2.4)
Mother	127 (100.0)	0 (0.0)
Father	4 (100.0)	0 (0.0)
Doctor	40 (95.2)	2 (4.8)
Friends	24 (100.0)	0 (0.0)
How effective is your management		
Effective	282 (100.0)	0 (0.0)
Not effective	33 (100.0)	0 (0.0)

Data presented as frequency (percent)

past 6 months. Although the 14.7% absenteeism reported was lower than that obtained by 46.3% in Farotimi, Esike, Nwozichi et al., [3] in Nigeria, 57% in Kumar, Konjengbam and Devi [12], Manipur and 80.6% in Tangchai, Titapant and Boriboonhirunsarn [31] in Thailand, it was found that absenteeism increases with increased severity of dysmenorrhea. Dysmenorrhea severity resulted in reduced class concentration and subsequently truancy which has direct negative effects on academic performance and consequently may lead to school drop-outs after junior high school (JHS).

Participants reported menarche at various ages with 77.8% of respondents with dysmenorrhea having their first menstruation between the ages 9-11 years. This is consistent with studies [12, 32] which reported 75.7% and 84.2% at menarche age of 9-11 years respectively. However, the result was different from that obtained by Nidhi, Benjamin, Balamma et al., [33] who reported that 47.0% of respondents with

dysmenorrhea experienced their first menstruation between the ages <12 years.

In this study, a total of 10% of participants had a family history of dysmenorrhea of which majority (90%) out of the total had experienced dysmenorrhea at least once in their lifetime. These findings are in line with previous studies [14,19,34] which found that the majority of the participants with dysmenorrhea had a history of dysmenorrhea in the family. Family history of dysmenorrhea appears to be an important risk factor for generational dysmenorrhea. The possible explanation was that daughters or mothers who complain of menstrual pains also experienced menstrual distress, and this was associated with behaviour learned from mothers [35].

In addition, this study found a relationship between dysmenorrhea and risk factors such as respondents who had menarche between the age 9-11 years, those who experienced short




Fig. 2. Prevalence of awareness of dysmenorrhea

(<21 days) menstrual cyclic length and those who experienced short (<2 days) and menstruation days. This agrees with studies [10, 12,18,36] who identified early age of menstruations, short (<21 days) menstrual cycle length and the short length of the menstrual period as possible risk factors for dysmenorrhea.

Information on awareness of dysmenorrhea was high, however, only 8.0% of the adolescent girls out of a total of 400 participants had heard of dysmenorrhea before menarche. This implies that the majority (92.0%) of the study participants were aware of dysmenorrhea after the first menstruations. The reason may be that many cultures in the Northern part of Ghana frown against the subject of menstruation and often see women in their menstrual period as not being clean. As a result, menstrual issues are covered in secrecy and are thought to be personal to only women. Similarly, 92% of Pakistan females reported needing more information about menstrual hygiene before menarche [37]. Again, 71% of girls in India did not have information about menstruation or dysmenorrhea at all [4].

Amongst those who had experienced dysmenorrhea, information on dysmenorrhea was sourced from family members (39.5%), teachers (20.5%), and friends (18.8%) with the least (3.3%) from the social media. In a similar study conducted by Ogunfowokan and Babatunde [24] in Nigeria, many adolescents (64%) received information on dysmenorrhea from their mothers. Furthermore, Chhabra, Gokhale and Yadav [4] in a study of pre-menarche information and dysmenorrhea among

young girls found that, amongst those with dysmenorrhea, 45% were given knowledge by their mothers, 16% received information from teachers in school, 36% from friends.

About 75.5% of the participants felt treatment was necessary for painful menstruation but only 10.5% consulted doctors with 30.8% resorting to self-medication. Involvement in self-medication for treatment of menstrual pains are in accordance with the 28.9% in rural Akwa Ibom in Nigeria [38], 34.7% in an Egyptian study [11], but lower compared with 50% [39] and 75% [9] amongst Nigerian secondary school girls. Apart from the negative side effects of self-treatment, indulgence in self-medication may have been the reason for low desire to contact medical doctors.

The strength of the study was that; respondents were selected from five (5) different districts in the Upper East region which do not limit the generalization of the results of the study setting. However, the limitations of the study were; the use of self-reported questionnaires and recall of painful menstruation for past 6 months could lead to false information being reported. Another limitation was that further histological investigation was not done to differentiate between primary (normal ovulatory cycle) and secondary (pelvic pathology) dysmenorrhea.

5. CONCLUSION

Dysmenorrhea is a common complaint among adolescent girls. The prevalence of dysmenorrhea in this study was high (85%) with girls who had experienced menarche between 9-

11years, short (<21 days) menstrual cyclic length and short (<2 days) menstrual days being at risk. Some of the negative effects included; poor interpersonal relationship with family and friends, depression, low classroom concentration, school absenteeism, reduced daily physical activity and self-medication. Information on awareness was high (92.0%) but girls become more aware years after menarche. Therefore, the study recommends that; reproductive health should be included in school health education programs early enough to improve menstrual health. Also, education should be extended to parents in order to address the reproductive health needs of daughters.

CONSENT

Informed consent was sought from the school authorities, parents as well as each participant before included in the study. Subjects who did not give their consent were excluded from the study.

ETHICAL APPROVAL

Ethical clearance was sought from the Ethical review board of the School of Allied Health Sciences, University for Development Studies and Upper East Regional Directorate of Ghana Education Service.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Aziato L, Dedey F, Clegg-Lampsey JNA, The experience of dysmenorrhoea among Ghanaian senior high and university students: Pain characteristics and effects. *Reproductive Health*. 2014;11(58):1- 8.
2. Pembe A, Ndolele N. Dysmenorrhoea and coping strategies among secondary school adolescents in Ilala District, Tanzania. *East African Journal of Public Health*. 2011;8(3): 233-237.
3. Farotimi AA, Esike J, Nwozichi CU, Ojedian TD, Ojewole FO. Knowledge, attitude and healthcare-seeking behavior towards dysmenorrhea among female students of a private university in Ogun State, Nigeria. *Journal of Basic and Clinical Reproductive Sciences*. 2015;4(1): 33-38.
4. Chhabra S, Gokhale S, Yadav S. Premenarche Information and Dysmenorrhea in Young Girls. *Insights Reprod Med*. 2017;1(2):6.
5. Sanyal S, Ray S. Variation in the menstrual characteristics in adolescents of West Bengal. *Singapore Medical Journal*, 2008;49(7): 542.
6. Harel Z. Dysmenorrhea in adolescents and young adults: from pathophysiology to pharmacological treatments and management strategies. *Expert Opinion on Pharmacotherapy*. 2008;9(15):2661-2672.
7. Ameade EPK, Amalba A, Mohammed BS. Prevalence of dysmenorrhea among University students in Northern Ghana; its impact and management strategies. *BMC Women's Health*. 2018;18(1):39.
8. Gumanga S, Kwame-Aryee R. Prevalence and severity of dysmenorrhoea among some adolescent girls in a secondary school in Accra, Ghana. *Postgrad Med J Ghana*. 2012;1(1):1-6.
9. Adinma ED, Adinma J. Perceptions and practices on menstruation amongst Nigerian secondary school girls. *African Journal of Reproductive Health*. 2008;12(1) :74-83.
10. Sidi I, Hounkpatin B, Obossou AAA, Salifou K, Vodouhe M. Primary dysmenorrhea in the schools of Parakou: Prevalence, Impact and Therapeutic Approach. *Gynecol Obstet (Sunnyvale)* 2016;6:376.
11. El Gilany, Badawi AK, El Fedawy S. Epidemiology of dysmenorrhoea among adolescent students in Mansoura, Egypt; 2005.
12. Kumar KS, Konjengbam S, Devi HS. Dysmenorrhea among higher secondary schoolgirls of Imphal West district, Manipur: A cross-sectional study. *Journal of Medical Society*. 2016;30(1):38.
13. Agarwal AK, Agarwal A. A study of dysmenorrhea during menstruation in adolescent girls. *Indian Journal of Community Medicine: Official Publication of Indian Association of Preventive & Social Medicine*. 2010;35(1):159.
14. Kural M, Noor NN, Pandit D, Joshi T, Patil A. Menstrual characteristics and prevalence of dysmenorrhea in college going girls. *Journal of Family Medicine and Primary Care*. 2015;4(3):426.

15. Harlow SD, Campbell OM. Epidemiology of menstrual disorders in developing countries: a systematic review. *BJOG: An International Journal of Obstetrics & Gynaecology: Review*. 2004;111(1):6-16.
16. Grandi G, Ferrari S, Xholli A, Cannoletta M, Palma F, Romani AC. Cagnacci, prevalence of menstrual pain in young women: What is dysmenorrhea? *Journal of Pain Research*. 2012;5:169.
17. Patel V, Tanksale V, Sahasrabhojane M, Gupte S, Nevrekar P. The burden and determinants of dysmenorrhoea: A population-based survey of 2262 women in Goa, India. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2006;113(4): 453-463.
18. Lee LK, Chen P, Lee K, Kaur J. Menstruation among adolescent girls in Malaysia: A cross-sectional school survey. *Singapore Medical Journal*. 2006;47(10): 869-874.
19. Gagua T, Tkeshelashvili B, Gagua D. Primary dysmenorrhea: prevalence in adolescent population of Tbilisi, Georgia and risk factors. *Journal of the Turkish German Gynecological Association*. 2012;13(3):162.
20. Al-Kindi R, Al-Bulushi A. Prevalence and impact of dysmenorrhoea among Omani high school students. *Sultan Qaboos University Medical Journal*. 2011;11(4): 485-491.
21. Mohamed EM. Epidemiology of dysmenorrhea among adolescent students in Assiut City, Egypt. *Life Sci J*. 2012;9(1): 348-353.
22. Wong LP. Attitudes towards dysmenorrhoea, impact and treatment seeking among adolescent girls: A rural school-based survey. *Australian Journal of Rural Health*. 2011;19(4):218-223.
23. Cochran W, Snedecor G. Analysis of variance: The random effects model. *Statistical Methods*, Iowa State University Press, Ames, IA. 1989;237-252.
24. Ogunfowokan AA, Babatunde OA. Management of primary dysmenorrhea by school adolescents in ILE-IFE, Nigeria. *The Journal of School Nursing*. 2010; 26(2):131-136.
25. Harlow SD, Campbell B. Ethnic differences in the duration and amount of menstrual bleeding during the postmenarcheal period. *American Journal of Epidemiology*. 1996;144(10):980-988.
26. Avasarala AK, Panchangam S. Dysmenorrhoea in different settings: Are the rural and urban adolescent girls perceiving and managing the dysmenorrhoea problem differently? *Indian Journal of Community Medicine: Official Publication of Indian Association of Preventive & Social Medicine*. 2008;33(4): 246-249.
27. Ortiz MI. Primary dysmenorrhea among Mexican university students: prevalence, impact and treatment. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2010;152(1):73-77.
28. Baidya S, Debnath M, Das R. A study of reproductive health problems among rural adolescent girls of Mohanpur block of West Tripura district. *Al Ameen J Med Sci*. 2014; 7(1):78-82.
29. McDowell I. *Measuring health: A guide to rating scales and questionnaires*. Oxford University Press, USA; 2006.
30. Takala EP, Pehkonen I, Forsman M, Hansson GÅ, Mathiassen SE, Neumann WP, Winkel J. Systematic evaluation of observational methods assessing biomechanical exposures at work. *Scandinavian Journal of Work, Environment & Health*. 2010;3-24.
31. Tangchai K, Titapant V, Boriboonhirunsarn D. Dysmenorrhea in Thai adolescents: Prevalence, impact and knowledge of treatment. *Journal-medical Association of Thailand*. 2004;87:S69-S73.
32. George NS, Priyadarshini S, Shetty S. Dysmenorrhoea among adolescent girls-characteristics and symptoms experienced during menstruation. *Nitte University Journal of Health Science*. 2014;4(3):45.
33. Nidhi S, Benjamin MS, Balamma S, Abdul Rahman AM, Abirami M, Adhil MHK, Ajantha P. Prevalence of dysmenorrhoea in college students. *International Journal of Scientific and Research Publications*, 2014;4(10):1-6.
34. Charu S, Amita R, Sujoy R, Thomas GA. Menstrual characteristics and prevalence and effects of dysmenorrhea on quality of life of medical students. *International Journal of Collaborative Research on Internal Medicine & Public Health*. 2012; 4(4):276.
35. Polat A, Celik H, Gurates B, Kaya D, Nalbant M, Kavak E, Hanay F. Prevalence of primary dysmenorrhea in young adult

- female university students. Archives of Gynecology and Obstetrics. 2009;279(4): 527-532.
36. Di Cintio E, Parazzini F, Tozzi L, Luchini L, Messopane R, Marchini M, Fedele L. Dietary habits, reproductive and menstrual factors and risk of dysmenorrhoea. European Journal of Epidemiology. 1997; 13(8):925-930.
37. House S, Mahon T, Cavill S. Menstrual hygiene matters: A resource for improving menstrual hygiene around the world. Reproductive Health Matters. 2013; 21(41):257-259.
38. Ekong I, Udofia E, Johnson O, Ekanem US, Okojie O. Menstrual problems and their prevalence among adolescents in Akwa Ibom. International Journal of Current Research and Academic Review. 2015;3(8):96-105.
39. Sule ST, Ukwenya JE. Menstrual experiences of adolescents in a secondary school. Turkish-German Gynecol Assoc. 2007;8(1):7-14.

© 2019 Ziba et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/49911>