



Effect of Blanching Pre-Treatments on Sensory Quality of Dragon Fruit Peel Pickle

Gohil Mehul M. ^{a++*}, Dev Raj ^{a#}, H. G. Suthar ^{at}
and N. V. Patel ^{at}

^a Department of Post Harvest Technology, ASPEE College of Horticulture, Navsari Agricultural University, Navsari, Gujarat, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i203855

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107174>

Original Research Article

Received: 21/07/2023

Accepted: 23/09/2023

Published: 30/09/2023

ABSTRACT

Present investigation entitled effect of blanching pre-treatments on quality of dragon fruit peel pickle was aimed to evaluate sensory quality of dragon fruit peel pickle during storage. For preparation of dragon fruit peel pickle, an experiment was laid out with six blanching pre-treatments [without blanching (T1), water blanching (T2), 0.5% citric acid blanching (T3), 1.0% citric acid blanching (T4), 0.5% alum (potassium aluminium sulphate) blanching (T5) and 1.0% alum (potassium aluminium sulphate) blanching (T6)]. The prepared pickle was stored for a period of 6 months to analyse the sensory quality attributes at three month intervals. The results of the investigation revealed that dragon fruit peel pickle prepared from peel blanched in 1 per cent citric acid (T₄) observed to have maximum colour, taste, flavour, texture and overall acceptability score. Overall findings of investigation revealed that best sensory quality dragon fruit peel pickle can be

⁺⁺Ph. D. Scholar;

[#]Professor and Head;

[†]Assistant Professor;

^{*}Corresponding author: E-mail: mehulgohil447@gmail.com;

prepared from peel when blanched in 1 per cent citric acid followed by curing in 8 per cent salt and addition of ingredients. The dragon fruit peel pickle can be successfully stored for a period of six months with minimum changes in sensoryquality.

Keywords: Dragon fruit peel; citric acid; alum; sensory.

1. INTRODUCTION

Dragon fruit, known as pitaya or pitahaya belongs to the Cactaceae family and is prevalent in two separate genera namely, 'Hylocereus' and 'Selenicereus'. The most common commercially cultivated varieties belongs to Hylocereus genus covering around 16 different species. It is also called as strawberry pear, thangloy (Vietnamese), pitayaroja (Spanish), la pitahaya rouge (French) andkamalam (Hindi). Dragon fruit has great potential as a new crop for mediterranean growers due to the requirement of little water and well adaption to the high temperatures [1]. The cultivation is prominent in about 20 countries including Thailand, Indonesia, Taiwan, Vietnam, India, Srilanka, Bangladesh, Japan, Malaysia, Philippines, Australia, United States, and China. There are three varieties that are grown commercially includes, *Hylocereus undatus* (white dragon fruit), *Hylocereus polyrhizus* (red dragon fruit) and *Selenicereus megalanthus* (yellow dragon fruit) [2]. It contributes to the food security of millions of people in the developing world. Gazing at the recent dragon fruit production scenario of 2018-2019, India is the producer of dragon fruit with annual production of 4200 ton from an area of 400 ha with 10.5 t/ha productivity [3].

In India, the indigenous fruits are processed into a number of value added products like, jam, juice, jelly, cheese, preserves etc. But the exotic dragon fruit processed products are rarely available in our markets as well as very little work has been done on processing of dragon fruit in India. So the scope for utilizing dragon fruit remains bright in India. A number of locally processed fruit products are now available in the market. If quality products from dragon fruit are developed, it might be welcomed by the consumers, because of this dragon fruit has gained much interest in the society due to its exotic features attractive colours, nutritional value and pleasant taste. Development of pickle by utilizing local produce is critically important for expansion of food industries. Therefore this study was planned keeping in view on medicinal and nutritional importance of dragon fruit peel, to

utilize peel for preparation of pickle for commercial exploitation.

Physico-chemical properties of the discarded dragon fruit peel can be determined in order to evaluate its potential for recovery of value-added materials. The moisture content of the peel reported approximately 92.7per cent and found low in total soluble solids, protein, ash and fat content. Betacyanin pigment (150.46 ± 2.19 mg/100 g) and pectin (10.8%) reported high in the peel. Glucose, maltose and fructose detected in the peel but not sucrose and galactose. The peel also had very high insoluble and soluble dietary fibre which had exhibited a good ratio of insoluble dietary fibre to soluble dietary fibre (3.8:1.0). The peel of dragon fruit possess higher radical scavenging activity than the pulp and thus could be a good source of antioxidants. The compounds detected in the peel are chlorogenic acid, gallic acid, and quercetin.

Dragon fruit peels account for approximately 25 per cent of the fruit's weight and contain a large variety of bioactive compounds (e.g., phenolics, betacyanin, vitamins, etc.). They have been proven to produce higher amounts of phenolic compounds (1.4 times in red-fleshed and 9.6 times in white-fleshed species) and antioxidant capacities than the pulp. Several studies have also reported that dragon fruit peels have higher cancer cell antiproliferation than their pulps [4]. The antioxidants in dragon fruit peel may include chlorogenic acid, gallic acid and quercetin [5]. Peel also possessa natural appealing red color due to the presence of betacyanin compounds that have a high stability under appropriate storage conditions [6]. Fresh and dried dragon fruit skin found rich in pectins and betalains making itnatural food thickener and natural colouring agent.

Pickles are preserved products prepared with salt alone or in combination with chilli and other spices and stored at room temperature for off-season utilization. The maintenance of dragon fruit peel colour is major problems in preparation of pickle. So, there is urgent need to standardize the pre-treatments for preparation of pickle from dragon fruit peel. Further, the quality of dragon

fruit peel pickle also needs the evaluation before going for commercialization of products.

2. MATERIALS AND METHOD

The fresh ripe dragon fruit were selected and washed with running tap water followed by trimming and peeling. The peel was cut into 12×17 mm±2 mm pieces. After that, the peel pieces were blanched as per treatment (control, blanching, 0.5% CA, 1% CA, 0.5% alum and 1% alum) at 95°C temperature for 5 min followed by draining to remove excess surface moisture. After pre-treatments, 4 per cent salt (NaCl) was added into the peel pieces, followed by mixing and then kept for 6 hours. After 6 hours, again additional 4 per cent salt (NaCl) was mixed and kept for 6 hours. The salt solution obtained due to mass transfer out from the peel pieces after 12 hours was discarded. Pickle of different treatments were prepared by addition of ingredients as shown in recipe. All the ingredients were taken in stainless steel vessels and added into the warm oil. After that blanched dragon fruit peel pieces were added into the oil containing salt and spices and mixed thoroughly with slow heating for 4-5 min. The prepared samples were then packed in glass jar and stored at room temperature for further analysis and to record observations. Principal steps used for preparation of dragon fruit peel pickle has been presented in Fig. 1.




Fig. 1. Flow chart for preparation of dragon fruit peel pickle

Recipe for Making Pickle: The recipe for the preparation of dragon fruit peel based pickle comprised of 1 kg peel pieces, 350 ml mustard oil, 25 g fenugreek, 22 g chilli powder, 10 g cumin, 6 g turmeric powder, 2 g asafoetida and 10 g coriander, 10 g fennel, 5 ml acetic acid and 1 g sodium benzoate.

3. RESULTS AND DISCUSSION

3.1 Colourscore

Perusal of data pertaining to effect of blanching pre-treatments on colourscore of dragon fruit peel pickle during six months storage has been presented in Table 1.

Effect of Pre-Treatments: Data showed that among different blanching pre-treatments, the mean colourscore of dragon fruit peel pickle (T) significantly varied between 6.49 and 7.36, with maximum colourscore in pickle which was prepared from peel which was blanched along with 1.0 per cent citric acid (T₄) and minimum colourscore in pickle prepared from unblanched dragon fruit peel (T₁). Alam et al. [7] reported similar findings with significant effect in colourscore of carrot powder.

Effect of Storage: Data depicted that storage of dragon fruit peel pickle resulted significant decrease in mean colourscore (S) from 7.30 to 6.48 (S₀ to S₂) during six months. Sangeeta et al. [8] also reported gradual decrease in colourscore of litchi aril during storage. Decrease in colourscore during storage might be due to degradation of pigments.

3.2 Taste Score

Perusal of data pertaining to effect of blanching pre-treatments on taste score of dragon fruit peel pickle during six months storage has been presented in Table 2.

Effect of Pre-Treatments: Data showed that among different blanching pre-treatments, the mean taste score of dragon fruit peel pickle (T) significantly varied between 6.69 and 6.97, with maximum taste score in pickle which was prepared from peel which was blanched along with 1.0 per cent citric acid (T₄) and minimum taste score in pickle prepared from water blanched peel only (T₂). Similar observation was recorded by Desayi [9] for partial dehydrated mushroom.

Effect of Storage: Data depicted that storage of dragon fruit peel pickle resulted significant increase in mean taste score (S) from 6.33 to 7.11 (S₀ to S₂) during six months. Haokip [10] also reported gradual increase in taste score of mango pickle during storage.

3.3 Flavour Score

Perusal of data pertaining to effect of blanching pre-treatments on flavour score of dragon fruit peel pickle during six months storage has been presented in Table 3.

Effect of Pre-Treatments: Data showed that among different blanching pre-treatments, the mean flavour score of dragon fruit peel pickle (T) varied between 6.62 and 6.77, with maximum flavour score in pickle which was prepared from peel which was blanched along with 1.0 per cent alum (T₆) and minimum flavour score in pickle prepared from unblanched dragon fruit peel (T₁) and water blanched peel only (T₂). Similar observation was recorded by Bairwa et al. [11] for pretreated aonla powder added in ready to serve beverage.

Effect of Storage: Data depicted that storage of dragon fruit peel pickle resulted significant increase in mean flavour score (S) from 6.44 to 6.98 (S₀ to S₂) during six months. Haokip [10] also reported gradual increase in flavour score of mango pickle during storage. Similar significant increase in flavour score was also reported by Bairwa et al. [11] in pretreated aonla powder added in ready to serve beverage.

3.4 Texture Score

Perusal of data pertaining to effect of blanching pre-treatments on texture score of dragon fruit peel pickle during six months storage has been presented in Table 4.

Effect of Pre-Treatments: Data showed that among different blanching pre-treatments, the mean texture score of dragon fruit peel pickle (T) varied between 7.32 and 7.50, with maximum texture score in pickle which was prepared from peel which was blanched along with 1.0 per cent citric acid (T₄) and 1.0 per cent alum (T₆) and minimum texture score in pickle prepared from water blanched peel only (T₂). However, effect of

Table 1. Effect of different treatments on colour score of dragon fruit peel pickle

Treatment (T)	Colour score (9 point Hedonic scale)			Mean (T)
	Storage (S)			
	S ₀ - Initial	S ₁ - 3 Month	S ₂ - 6 Month	
T ₁ (C)	7.00	6.47	6.00	6.49
T ₂ (B)	7.17	6.67	6.00	6.61
T ₃ (BC 0.5 %)	7.33	7.00	6.67	7.00
T ₄ (BC 1.0 %)	7.67	7.40	7.00	7.36
T ₅ (BA 0.5 %)	7.20	6.83	6.33	6.79
T ₆ (BA 1.0 %)	7.40	7.17	6.87	7.15
Mean (S)	7.30	6.92	6.48	6.90
S.Em.±	T	S	T×S	CV %
	0.040	0.033	0.070	T:1.75
CD_{0.05}	0.116	0.131	0.201	S:2.06

Table 2. Effect of different treatments on taste score of dragon fruit peel pickle

Treatment (T)	Taste score (9 point Hedonic scale)			Mean (T)
	Storage (S)			
	S ₀ - Initial	S ₁ - 3 Month	S ₂ - 6 Month	
T ₁ (C)	6.25	6.89	7.00	6.71
T ₂ (B)	6.25	6.83	7.00	6.69
T ₃ (BC 0.5 %)	6.33	6.83	7.00	6.72
T ₄ (BC 1.0 %)	6.42	7.17	7.33	6.97
T ₅ (BA 0.5 %)	6.33	7.00	7.17	6.83
T ₆ (BA 1.0 %)	6.42	7.00	7.17	6.86
Mean (S)	6.33	6.95	7.11	6.80
S.Em.±	T	S	T×S	CV %
	0.052	0.024	0.090	T:2.29
CD_{0.05}	0.149	0.094	NS	S:1.50

Table 3. Effect of different treatments on flavour score of dragon fruit peel pickle

Treatment (T)	Flavour score (9 point Hedonic scale)			Mean (T)
	Storage (S)			
	S ₀ - Initial	S ₁ - 3 Month	S ₂ - 6 Month	
T ₁ (C)	6.33	6.61	6.92	6.62
T ₂ (B)	6.33	6.61	6.92	6.62
T ₃ (BC 0.5 %)	6.50	6.72	7.00	6.74
T ₄ (BC 1.0 %)	6.50	6.77	7.00	6.76
T ₅ (BA 0.5 %)	6.50	6.72	7.00	6.74
T ₆ (BA 1.0 %)	6.50	6.77	7.05	6.77
Mean (S)	6.44	6.70	6.98	6.71
S.Em.±	T	S	T×S	CV %
	0.053	0.058	0.091	T:2.36
CD_{0.05}	NS	0.226	NS	S:3.64

treatment was found to have non-significant effect. Similar observation was recorded by Arendse and Jideani [12] for moringa powder.

Effect of Storage: Data depicted that storage of dragon fruit peel pickle resulted significant increase in mean texture score (S) from 7.12 to 7.70 (S₀ to S₂) during six months. Haokip [10] also reported gradual increase in texture score of mango pickle during storage. Similar significant increase in texture score was also reported by Singh [13] in cauliflower pickle. Increase in texture score during storage might be due to softening of pickle as reported by Sharma [14] in lime pickle during storage.

3.5 Overall Acceptability Score

Perusal of data pertaining to effect of blanching pre-treatments on overall acceptability score of dragon fruit peel pickle during six months storage has been presented in Table 5.

Effect of Pre-Treatments: Data showed that among different blanching pre-treatments, the

mean overall acceptability score of dragon fruit peel pickle (T) significantly varied between 6.80 and 7.15, with maximum overall acceptability score in pickle which was prepared from peel which was blanched along with 1.0 per cent citric acid (T₄) and minimum overall acceptability score in pickle prepared from unblanched dragon fruit peel (T₁) and water blanched peel only (T₁). Similar observation was recorded by Jyoti et al. [15] for mushroom pickle.

Effect of Storage: Data depicted that storage of dragon fruit peel pickle resulted significant increase in mean overall acceptability score (S) from 6.80 to 7.07 (S₀ to S₂) during six months. Haokip [10] also reported gradual increase in overall acceptability score of mango pickle during storage. Similar significant increase in overall acceptability score was also reported by Singh [13] in cauliflower pickle. Increase in overall acceptability score during storage might be due to use of many raw materials which were soluble in water as reported by Azzawi and Abdullah [16] in pickle during six months storage [17-19].

Table 4. Effect of different treatments on texture score of dragon fruit peel pickle

Treatment (T)	Texture score (9 point Hedonic scale)			Mean (T)
	Storage (S)			
	S ₀ - Initial	S ₁ - 3 Month	S ₂ - 6 Month	
T ₁ (C)	7.17	7.33	7.60	7.37
T ₂ (B)	7.00	7.37	7.60	7.32
T ₃ (BC 0.5 %)	7.11	7.43	7.67	7.40
T ₄ (BC 1.0 %)	7.17	7.50	7.83	7.50
T ₅ (BA 0.5 %)	7.11	7.43	7.67	7.40
T ₆ (BA 1.0 %)	7.17	7.50	7.83	7.50
Mean (S)	7.12	7.43	7.70	7.42
S.Em.±	T	S	T×S	CV %
	0.049	0.046	0.085	T:1.99
CD_{0.05}	NS	0.180	NS	S:2.62

Table 5. Effect of different treatments on overall acceptability score of dragon fruit peel pickle

Treatment (T)	Overall acceptability score(9 point Hedonic scale)			Mean (T)
	Storage (S)			
	S ₀ - Initial	S ₁ - 3 Month	S ₂ - 6 Month	
T ₁ (C)	6.69	6.83	6.88	6.80
T ₂ (B)	6.69	6.87	6.88	6.80
T ₃ (BC 0.5 %)	6.82	7.00	7.09	6.97
T ₄ (BC 1.0 %)	6.94	7.21	7.29	7.15
T ₅ (BA 0.5 %)	6.78	7.00	7.04	6.94
T ₆ (BA 1.0 %)	6.87	7.11	7.23	7.07
Mean (S)	6.80	7.00	7.07	6.96
S.Em.±	T	S	T×S	CV %
	0.027	0.018	0.047	T:1.17
CD_{0.05}	0.078	0.071	NS	S:1.11

4. CONCLUSION

Overall from the findings of investigation it can be concluded dragon fruit peel pickle can be prepared from peel when blanched in 1 per cent citric acid followed by curing in 8 per cent salt and addition of ingredients. The pickle prepared from the above treatment, possess higher sensory quality attributes during storage. Thus, the developed technology can commercially be adopted by food processing industry for the production of pickle by utilizing dragon fruit peel.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Trivellini A, Lucchesini M, Ferrante A, Massa D, Orlando M, Incrocci L, Mensuali Sodi A. Pitaya, an attractive alternative crop for mediterranean region. *J. Agron.* 2020;10(8):1065.
- Jalgaonkar K, Mahawar MK, Bibwe B, Kannaujia P. Postharvest profile, processing and waste utilization of dragon fruit (*Hylocereus Spp.*): A review. *Food Reviews International.* 2022;38(4):733-759.
- Wakchaure GC, Kumar S, Meena KK, Rane J, Pathak H. Dragon Fruit Cultivation in India: Scope, Constraints and Policy Issues. *Technical Bulletin.* 2021;27:47.
- Luo H, Cai Y, Peng Z, Liu T, Yang S. Chemical composition and in vitro evaluation of the cytotoxic and antioxidant activities of supercritical carbon dioxide extracts of pitaya (Dragon fruit) peel. *Chem. Cent. J.* 2014;8(1):1-10.
- Lourith N, Kanlayavattanakul M. Antioxidant and stability of dragon fruit peel colour. *Agro. Food Ind. Hi Tech.* 2013; 24(3):56-58.
- Woo KK, Ngou FH, Ngo LS, Soong WK, Tang PY. Stability of betalain pigment from red dragon fruit (*Hylocereus polyrhizus*). *Am. J. Food Technol.* 2011;6(2):140-148.
- Alam MS, Gupta K, Khaira H, Javed M. Quality of dried carrot pomace powder as affected by pretreatments and methods of drying. *Int. J. Agric. Eng.* 2013;15(4):236-243.
- Sangeeta S, Chopra CS, Sharma SK. Development of osmo-air dried litchi arils from fruits affected with pericarp browning using response surface methodology. *J. Hill Agric.* 2016;7(2):255-261.
- Desayi D. Development, recovery and sensory evaluation of fresh mushroom chips. *Int. J. Food Agr. Vet. Sci.* 2012; 2(2):190-193.
- Haokip H, Abonmai T, Singh A. Changes in physico-chemical properties of mango pickle during storage. *Int. J. Plant Soil Sci.* 2022;34(23):884-890.
- Bairwa KM, Bahadur V, Rolaniya MK, JayvanthKumar U. Study on changes in organoleptic quality of value added blended herbal Ready-to-serve (RTS) of aonla (*Emblica officinalis* Garetn.) CV. NA-6 during storage period. *J. Pharmacogn. Phytochem.* 2018;1:708-711.
- Arendse W, Jideani V. Storage stability and consumer acceptability of dried apple: Impact of citric acid, potassium sorbate and Moringa oleifera leaf extract powder. *Foods.* 2022;11(7):984.
- Singh SN, Patel VA, Varma MG, Patel SG, Savaliya MV. Effect of processing methods on physico-chemical properties, sensory

- evaluations and storage studies of cauliflower (*Brassica oleracea* L. var. botrytis) Pickle. Int. J. Curr. Microbiol. Appl. Sci. 2021;10(01):1413-1428.
14. Sharma A. Studies on preservation of products of sour lime (*Citrus aurantifolia* Swingle). M.Sc. Thesis, CCS Haryana University, Hisar; 2022.
 15. Jyoti S, Sindhu SC, Sindhu A. Development and evaluation of value added pickle from dehydrated shiitake (*Lentinus edodes*) Mushroom. Int. J. Food Sci. Nutr. 2016;1(1):24-26.
 16. Azzawi AA, Abdullah BY. Study of physico-chemical and nutritional properties of some processed pickles. J. Agric. Sci. 2019; 19(2):45-54.
 17. Ajibola OO, Thomas R, Bakare BF. Selected fermented indigenous vegetables and fruits from Malaysia as potential sources of natural probiotics for improving gut health. Food science and human wellness. 2023;12(5):1493-509.
 18. Deepa J, Rajkumar P. Mathematical modeling on drying of tamarind fruits. Mathematical modeling on drying of tamarind fruits. Journal of scientific and industrial research (JSIR). 2023;82(06): 661-70.
 19. Uppaluri RV, Das C, Goud VV. Anandalakshmi R, editors. Agro and food processing technologies: Proceedings of NERC 2022. Springer Nature; 2023.

© 2023 Mehul et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107174>