



# Growth Factors for a Strawberry Crop are Affected by Microalgae and Vermicompost

Sachin Kumar Singh <sup>a\*</sup>, Jasveer Singh <sup>a</sup>, Rajveer <sup>b</sup>,  
Sobita Simon <sup>a</sup> and Naseeb Choudhary <sup>c</sup>

<sup>a</sup> Department of Plant Pathology, Sam Higginbottom University of Agriculture, Technology, and Sciences, Prayagraj, Uttar Pradesh, India.

<sup>b</sup> SMS Soil science, Swami Keshwan and Rajasthan Agricultural University, Bikaner, Rajasthan, India.

<sup>c</sup> Department of Agricultural Economics, CCSHAU, Hisar, Haryana, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/JEAI/2023/v45i92189

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104274>

Original Research Article

Received: 28/05/2023

Accepted: 01/08/2023

Published: 05/08/2023

## ABSTRACT

A Randomized Block Design (RBD) field experiment was conducted in Central Research Farm (CRF) at Department of Plant Pathology, SHUATS, Prayagraj, U.P. during the Rabi season of 2019-2020 in pots. Growth factors for a strawberry crop are affected by microalgae and vermicompost different dose of microalgae and vermicompost on number of leaves and height per plant of strawberry at 30, 60 and 90 DAT under field condition.

The maximum number of leaves (per plant) was recorded T<sub>9</sub> @ Microalgae 8g + vermicompost (16, 18 and 23.50 per plant, respectively) followed by T<sub>6</sub> @ Microalgae @ 5g/pot + Vermicompost (15.75, 20.00 and 21.50 per plant, respectively) at 30, 60 and 90 DAT, respectively. The treatment T<sub>9</sub> @ Microalgae 8g + vermicompost recorded maximum plant height (cm) (14.80, 23.03 and 28.58

\*Corresponding author: E-mail: 18msspp040@shiats.edu.in;

cm, respectively) followed by T<sub>8</sub> @ Microalgae @ 7g/pot + Vermicompost (14.13, 22.23 and 26.43cm, respectively) at 30, 60 and 90 DAT, respectively. Shows significant increment in the number of leaves per plant, at different intervals and all the other treatments were non-significant among each other. And all the treatments showed significant result in comparison to control treatment.

**Keywords:** *Alternaria Spp.*; microalgae; vermicompost; plant height and leaves.

## 1. INTRODUCTION

The use of organic amendments, such as traditional thermophilic composts, has long been recognized as an effective means of improving soil structure, enhancing soil fertility [1], Strawberry is very susceptible to a number of fungal diseases, which make the defense management practices fundamental to prevent severe economic losses. Fungal pathogens can attack the crop in all cultivation systems and can cause strawberry fruit decay in post-harvest. Among soil-borne fungi that can infect the plant in soil cultivation, *Sclerotinia sclerotiorum*, *Rhizoctonia solani*, *Phytophthora spp.*, *Pythium spp.*, *Fusarium oxysporum*, *Verticillium dahliae*, and *Macrophomina phaseolina* are the most important. They affect the root system, hindering the nutrient and water absorption from soil increasing microbial diversity and populations [2], microbial activity [3], improving the moisture-holding capacity of soils and increasing crop yields. These pathogens cause severe symptoms such as yellowing, wilting, damping-off, root rot, and collar rot and, in conventional cultivation systems, they are controlled by synthetic compound applications. Effects on microorganisms have also been associated with their capability to suppress soil-borne plant diseases [4] plant parasitic nematode populations and increased crop yields [5].

Among foliar pathogens, *Podosphaera aphanis* is responsible for powdery mildew, a severe disease that causes important economic losses requiring several fungicidal treatments [6].

## 2. MATERIALS AND METHODS

Nine treatments were applied to the plots namely vermicompost (T<sub>1</sub>), Microalgae @ 1g/pot + vermicompost (T<sub>2</sub>), Microalgae @ 2g/pot + Vermicompost (T<sub>3</sub>), Microalgae @ 3g/pot + Vermicompost (T<sub>4</sub>), Microalgae @ 4g/pot + Vermicompost (T<sub>5</sub>), Microalgae @ 5g/pot + Vermicompost (T<sub>6</sub>), Microalgae @ 6g/pot + Vermicompost (T<sub>7</sub>), Microalgae @ 7g/pot + Vermicompost (T<sub>8</sub>), Microalgae @ 8g/pot + Vermicompost (T<sub>9</sub>). T<sub>0</sub> was kept as untreated control.

### 2.1 Methods

**Pot filling:** The soil for pot experiments was collected from the university field and brought to the Bio-net. The pots of 15 cm diameter were filled with 250cc of soil and vermicompost used for various experiments.

**Application of vermicompost:** 4kg vermicompost was applied in my total 40 pots at the @ of 100g per pot, vermicompost was mixed with the soil present in the pot manually and then the pot was ready for transplant.



**Fig. 1. Transplanted plants**



**Fig. 2. Application of micro algae**

## 2.2 Application of Microalgae

Microalgae was applied after 15 days of transplanting on the rhizosphere plants according to the Treatment given in concentration of 1g per pot to 8g per pot, second application of micro algae was done at 30 days after transplanting.

## 3. RESULTS AND DISCUSSION

Different dose of microalgae and vermicompost on number of leaves per plant of strawberry at 30, 60 and 90 DAT under field condition.

The results presented among all the treatments T<sub>9</sub> shows significant increment in the number of leaves per plant, at different intervals and all the other treatments were non-significant among each other. And all the treatments showed significant result in comparison to control treatment.

The data presented in table different dose of microalgae and vermicompost on height of per plant of strawberry at 30, 60 and 90 DAT under Bio-net condition.

The results presented in indicates that among all the treatments T<sub>9</sub> shows significant increment in the number of leaves per plant, at different intervals and all the treatment were non-significant among each other but significant with control treatment.

This result is in agreement with Edra et al. [7] who found that the brown-alga *Lessonia trabeculata* inhibited bacterial growth and reduced both the number and size of the necrotic lesions in tomato leaves following infection with *Botrytis cinerea*. Marine algae represent a great source of a vast variety of complex natural products and could be a promising source of a novel bioactive compound that can help plant survival by offering protection against stress imposed by pathogens. Aqueous and ethanolic extracts from the red-alga *Gracillaria chilensis* prevent the growth of *Phytophthora cinnamomi*.

Wafaa et al. [8] also reported that seeds and foliar application of Blue-green Cyanobacteria *Oscillatoria agardhii* showed significant effective net blotch, powdery mildew, rust and spot blotch management. Blue-green Algae applied treatments increased antioxidant enzymes as catalase (CAT), peroxidase (POD) and superoxide dismutase (SOD), as well as grain yield, kernel weight in most cases and improved barley yield quality parameters and grain protein. Burjus et al. [9] also observed that biomass obtained from cultivation of *N.commune*, *Anabaena circinalis* in bioreactor to inoculated mature compost was the best way of cultivation of nitrogen fixing cyanobacteria with low cost. The application of cyanobacteria in growth and yield of chickpea plant reduces the need of chemical fertilizers about 30%-50% and subsequently reduces environmental pollution compared with other mineral chemical fertilizers.

**Table 1. Effect of micro algae and vermicompost on number of leaves perplant of strawberry at different days of intervals**

Treatments	30 DAT	60 DAT	90 DAT
T <sub>0</sub>	12.25	17.00	18.00
T <sub>1</sub>	15.25	18.00	20.75
T <sub>2</sub>	14.25	18.00	18.50
T <sub>3</sub>	14.50	18.50	20.50
T <sub>4</sub>	15.25	17.75	18.75
T <sub>5</sub>	13.75	18.25	21.00
T <sub>6</sub>	15.75	19.00	21.50
T <sub>7</sub>	14.75	17.75	19.25
T <sub>8</sub>	14.50	15.50	20.75
T <sub>9</sub>	16.00	20.00	23.50
<b>F-test</b>	<b>S</b>	<b>S</b>	<b>S</b>
<b>C.D. at 0.5%</b>	<b>1.088</b>	<b>3.285</b>	<b>3.776</b>
<b>S.Ed. (+)</b>	<b>0.530</b>	<b>1.601</b>	<b>1.840</b>

**Table 2. Effect of microalgae and vermicompost on plant height (cm) of strawberry at different days of intervals**

Treatment S	30 DAT	60 DAT	90 DAT
T <sub>0</sub>	10.43	18.48	23.18
T <sub>1</sub>	11.23	18.83	23.73
T <sub>2</sub>	11.75	19.00	23.55
T <sub>3</sub>	12.00	18.83	23.85
T <sub>4</sub>	12.35	20.03	25.05
T <sub>5</sub>	12.90	19.73	25.30
T <sub>6</sub>	12.15	19.95	25.98
T <sub>7</sub>	13.48	22.15	27.20
T <sub>8</sub>	14.13	22.23	26.43
T <sub>9</sub>	14.80	23.03	28.58
<b>F-test</b>	<b>S</b>	<b>S</b>	<b>S</b>
<b>C.D. at 0.5%</b>	<b>1.279</b>	<b>1.428</b>	<b>1.437</b>
<b>S.Ed. (+)</b>	<b>0.623</b>	<b>0.696</b>	<b>0.700</b>

### 3.1 Discussion

Vermicompost and microalgae additions gave considerably improved outcomes, as evidenced by the strawberry crop's maximum number of leaves and plant height. Vermicompost and microalgae applications significantly increased strawberry growth and yields, as evidenced by increases of up to 37% in leaf area, 37% in plant shoot biomass, 40% in the number of flowers, 36% in the number of plant runners, and 35% in the weight of marketable fruit. Not nutrients, but the availability of plant development-influencing substances like plant growth regulators and humic acids, created by the much expanded microbial populations as a result of earthworm activity, was what made vermicompost beneficial for plant growth and yield. The availability of macronutrients in soils treated with vermicomposts in our current experiment could not account for the gains in strawberry

development and production because they were equalized at transplanting in comparison to the plots treated with microalgae. Analysis of the soil revealed no appreciable differences in the amounts of macronutrients in any of the plots after the addition of vermicomposts and microalgae. It was confirmed in a later greenhouse experiment that small concentrations of humic acids, extracted from vermicompost and substituted into container media, increased the growth of plants in a similar pattern independent of nutrient supply [10]. It has been shown that microorganisms can produce materials that may affect plant growth such as substances acting as plant hormone analogues or growth regulators [11,12].

### 4. CONCLUSION

A trial was conducted to check the effect of micro algae and vermicompost on leaf spot (*Alternaria*

sp.) disease of strawberry. The experiment was analysed by using RBD (randomized block design) with 4 replication, 9 treatments along with the control. Observations recorded were disease intensity at 30, 60 and 90 (days after transplanting), plant growth parameters such as plant height at 30,60 and 90 DAT. Among all the treatments T<sub>9</sub> (Microalgae 8g + vermicompost) shows significant increment in the number of leaves, Plant height at different intervals and all the other treatments were non-significant among each other. And all the treatments showed significant result in comparison to control treatment. And treatment T<sub>9</sub> shows least percent disease infestation.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Follet R, Donahue R, Murphy L. Soil and Soil Amendments. Prentice-Hall, Inc., New Jersey; 1981.
2. Barakan FN, Salem SH, Heggo AM, Bin-Shiha MA. Activities of rhizosphere microorganisms as affected by application of organic amendments in a calcareous loamy soil. 2. Nitrogen transformation. Arid Soil Research and Rehabilitation. 1995; 9(4):467– 480.
3. Zink TA, Allen MF. The effects of organic amendments on the restoration of a disturbed coastal sage scrub habitat. Restoration-Ecology. 1998;6(1): 52–58.
4. Hoitink HA, Fahy P. Basis for the control of soilborne plant pathogens with composts. Annual Review Phytopathology. 1986; 24:93– 114.
5. Johnston AM, Janzen HH, Smith EG. Long-term spring wheat response to summer fallow frequency and organic amendment in southern Alberta. Canadian Journal of Plant Science. 1995;75(2): 347–354.
6. Romero AJ, Carvajal SC. Adolescent Bicultural Stress And Its Impact On Mental Well-Being Among Latinos, Asian Americans, And European Americans. Journal of Community Psychology. 2007; 35(4):519–534.
7. Edra J, Dorta F, Medina C, Ramírez A, Ramírez I and Peña-Cortés H. Anti-Phytopathogenic Activities of Macro-Algae Extracts Mar Drugs. 2011;9:739-756.
8. Haggag M, Wafaa MM, Hussein Mekhail E, El-Habbash. Res. J. Pharmaceutical, Biolo.and Chem. Sci. 2014;3(11):1530.
9. Burjus, Sanaa J, Jawad, Abdul latif M, Al-Ani, Nabeel K. Effect of Two Species of Cyanobacteria as Biofertilizers on Characteristics and Yield of Chickpea Plant. Iraqi Journal of Science. 2014; 55(2B):685-696
10. Atiyeh RM, Arancon N, Edwards CA, Metzger JD. Influence of earthworm-processed pig manure on the growth and yield of greenhouse tomatoes. Bioresource Technology 2000a;75(3):175– 180.
11. Frankenberger Jr. WT, Arshad M. Phytohormones in Soils: Microbial Production and Function. Marcel and Deckker, New York. 1995;503.
12. Brown GG. How do earthworms affect microfloral and faunal community diversity? Plant and Soil. 1995;170: 209–231.

© 2023 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/104274>