



Assessing the Nutrient Status and Soil Fertility Using Nutrient Indexed of Farmer's Fields in Chikkaballapura District, Karnataka

Kollalu Sandhya ^{a*}, B. Gayathri ^b, M. Papireddy ^a,
R. Praveenkumar ^a, Vishwanath ^a, B. Swathi ^a
and D. V. Naveen ^c

^a ICAR-KVK, Chintamani, Chikkaballapura, 563125, University of Agricultural Sciences, GKVK, Bangalore, India.

^b Department of Soil Science, University of Agricultural Sciences, GKVK, Bangalore, India.

^c College of Sericulture, Chintamani 563125, University of Agricultural Sciences, GKVK, Bangalore, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i234283

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111114>

Original Research Article

Received: 15/10/2023

Accepted: 21/12/2023

Published: 23/12/2023

ABSTRACT

The soil samples were collected by farmers in their own cultivable land analysed at soil testing laboratory at KVK, Chikkaballapura. From last decade (2012 to 2022), the samples were analysed following standard procedures. Nutrient status and nutrient Index was estimated to know the fertility status of the soils. The pH of soils of Chikkaballapura district was found be neutral (65.15 %) followed by alkaline (24.33%) and acidic (10.52 %) and low (68.09%) to medium (31.65%) range in

*Corresponding author: E-mail: sandhyak739@gmail.com;

EC values. Organic carbon content was found lower (89.64%) in majority of the soils and medium range in 10.02 % soils. Likewise OC values, available N status was in lower limits (63.63 kg ha⁻¹) and 36.37 % soils are in medium range. Available P₂O₅ status was recorded lower in 91.63 % and medium range in 8.19 % of the soils. Majority of the soils of the Chikkaballapura district is in medium range of available K₂O according to critical limits. With respect to Nutrient Index (NI) values for OC, available N, P₂O₅ and K₂O more than 90% soil samples found in lower category with exception to Chikkaballapura taluk for N and K₂O. In this study, it was found that most of the examined soils are in low in important parameters such as OC, N, P₂O₅ and K₂O. In this context, the adoption of appropriate soil management strategies would be helpful to improve the low and medium soil nutrient index of the soil.

Keywords: Soil fertility; nutrient index; organic carbon; soil nitrogen; phosphorus; potassium.

1. INTRODUCTION

Soil fertility evaluation is a critical aspect of soil management in agriculture. Achieving and maintaining appropriate levels of soil fertility, especially plant nutrient availability, is of paramount importance if agricultural land is to remain capable of sustaining crop production at an acceptable level [1,2]. Soil testing is often used to determine the nutrient status of crops and it also helps in develop the cost effective nutrient management practices to recommend to the farmers [3-6]. Soil fertility plays an important role in sustaining crop productivity of an area, particularly in situations where input of nutrients application differs and the information on the nutritional status can go a long way to develop economically viable alternatives for management of deficient nutrients in the soil [7].

Chikkaballapura district comes under eastern dry zone of Karnataka (Zone 5) and has basically agriculture oriented economy. The district consisting of six taluks viz., Bagepalli, Chikkaballapur, Chintamani, Gawribidanur, Gudibande and Shidlaghatta. The district lies between 13° 26' 3" North, 77° 43' 27" East. The total geographical area of the district is 190.50 lakh hectares with 107 lakh hectares under cultivation and only 24 percent of land is under irrigation and remaining 76 percent under rainfed farming. The geological formation of the Chikkaballapura district is made up of cretaceous red loamy soils, red sandy loam soils, sandy clay loam, loamy soils, sandy soils etc. The major crops cultivated in Chikkaballapura district are finger millet, groundnut, maize, pigeon pea, field bean, sunflower and horticulture crops like plantations and orchards of guava, grapes, pomegranate, sapota, vegetables crops like tomato, cabbage, chilli, capsicum, cucumber and cut flowers etc. The district is surrounded by Andhra Pradesh in north, Bangalore Rural district

by south, Kolar district and Tumkur district in east and west respectively. Entire district is principally plain region and having a typical rainfed situation with an average rainfall of 722 mm with 46 rainy days. Districts average minimum temperature is 19.33°C and maximum temperature is 31.33°C [8].

The district is characterized by low, scanty and uneven distributed rainfall with shallow and poor soils. The status of agricultural productivity in the district indicates that there is very good scope to increase the productivity of many crops. But, the farmers lag behind in adopting new technologies, in this respect, the extension machinery needs to be strengthened to reach the farmers in order to convey and demonstrate the latest technologies. In general, the fertility status of soils needs to be improved so that crop yields are stabilized [9]. In this point of view, this article comprise of compilation of the data for soil samples brought since from 2012 to 2022 for analysis by farmer themselves to Krishi Vignana Kendra (KVK), Chikkaballapura District to deliberate the nutrient status and soil fertility.

2. METHODOLOGY

The soil samples are collected by farmers in their own field and brought to the laboratory for soil testing at KVK, Chikkaballapura to know the fertility status. From last decade (2012 to 2022) the samples were analysed following standard procedures (Table 1). The received soil samples from farmers were processed by shade drying, powdering, and passed through 2 mm sieve and stored in the plastic containers. Later these samples were analyzed for pH (Soil reaction (1:2.5)), EC (Electrical conductivity (dSm⁻¹)), O.C. (Organic carbon (%)), N (Available nitrogen (kg ha⁻¹)), P₂O₅ (Available phosphorus (kg ha⁻¹)) and K₂O (Available potassium (kg ha⁻¹)) (Table 1).

Table 1. Number of soil samples analysed and per cent of samples falling in each category for different soil properties in Chikkaballapura District

Parameters	Critical limits *	Percentage of soil samples in each category (%)	Number of soil samples	*References
pH (1:2.5 soil: water)	Acidic (<6.50)	10.52	1148	[1]
	Neutral (6.5.-7.50)	65.15	7110	
	Alkaline (>7.50)	24.33	2655	
EC (dS m ⁻¹)	Low (<0.80)	68.09	7431	[1]
	Medium (0.80-1.60)	31.65	3454	
	High (>1.60)	0.26	28	
O.C (%)	Low (<0.50)	89.64	9157	[2]
	Medium (0.50-0.75)	10.02	1024	
	High (>0.75)	0.33	34	
N(kg ha ⁻¹)	Low (<280)	63.63	6944	[3]
	Medium (280-560)	36.37	3969	
	High (>560)	0.00	0	
P ₂ O ₅ (kg ha ⁻¹)	Low (<22.90)	91.63	10000	[4]
	Medium (22.90-56.33)	8.19	894	
	High (>56.33)	0.12	13	
K ₂ O(kg ha ⁻¹)	Low (<114)	37.35	4076	[5]
	Medium (114-336)	62.19	6787	
	High (>336)	0.39	43	

(EC – Electrical Conductivity; O.C – Organic Carbon; Available nitrogen as N (kg ha⁻¹); Available phosphorus as P₂O₅ (kg ha⁻¹); Available potassium as K₂O (kg ha⁻¹))

2.1 Statistical Analysis and Nutrient Index

Mean and standard deviation for the number of soil samples collected from each taluks were counted for decade (from 2012 to 2022) and calculated by using excel software. Nutrient Index introduced by [10] and modified by [11] was used to compare the soil fertility level (Table 3).

$$\text{Nutrient Index} = \{(1 \times A) + (2 \times B) + (3 \times C)\} / \text{NS}$$

A= Number of samples in low category
 B= Number of samples in medium category
 C= Number of samples in high category
 NS = Total number of samples

3. RESULTS AND DISCUSSION

3.1 Soil Details of Chikkaballapura District

The soil of Chikkaballapura district can be categorised into red clay, lateritic clay, red clayey and alluvial soils based on the dominating size of

the particles within the soil (Fig. 1a). Among six taluks of the district, red clayey soil was found to be dominated in Shidlagatta, Chintamani and part of Bagepalli and lateritic clayey soil was dominated in Chikkaballapura taluk whereas, Gudibande and Gauribidanuru taluks were dominated with red clay and alluvial soil. Red soils are typically difficult for crop cultivation because high leaching leads to low water holding capacity, low nutrients, low organic matter (humus), and acidification.

Lateritic soils are most highly weathered soils in the classification system and significant features of the lateritic soils are their unique color, poor fertility, and high clay content and lower cation exchange capacity [12] Soil depth (Fig. 1b) was found to be moderate shallow to deep in all the taluks whereas, sandy soils were found in Chintamani, Shidlagatta and Bagepalli taluk and loamy textured soils in Gudibande and Gowribidanuru (Fig. 1c). Land capability was found to be good in Gouribidanuru, Gudibande, Chikkaballapura and part of bagepalli taluk (Fig. 1d).

Table 2. Descriptive statistics of the measured soil properties in Chikkaballapura district covering the six taluks

Taluk /Properties	Bagepalli Mean± SD	Shidlagatta Mean± SD	Chikkaballapura Mean± SD	Gudibande Mean± SD	Gowribidanuru Mean± SD	Chintamani Mean± SD
pH (1:2.5 soil: water)	7.58±0.75	7.12±1.64	6.89±1.90	7.30±0.68	7.43±0.63	7.40±1.37
EC(dS m ⁻¹)	0.22±0.21	0.16±0.65	0.22±0.99	0.22±0.20	0.20±0.20	0.16±0.21
O.C (%)	0.25±0.15	0.36±2.95	0.55±1.97	0.25±0.72	0.22±0.58	0.34±1.31
N(kg ha ⁻¹)	49.12±37.79	57.27±53.06	48.85±40.54	52.22±37.04	45.55±19.10	61.69±58.31
P ₂ O ₅ (kg ha ⁻¹)	15.39±5.82	17.91±6.58	18.15±6.96	14.97±5.99	13.37±5.84	17.81±11.89
K ₂ O(kg ha ⁻¹)	109.32±46.16	112.94±34.89	111.77±28.90	104.40±20.66	108.01±70.79	117.82±43.40

(EC – Electrical Conductivity; O.C – Organic Carbon; Available nitrogen as N (kg ha⁻¹); Available phosphorus as P₂O₅ (kg ha⁻¹); Available potassium as K₂O (kg ha⁻¹))

Table 3. Nutrient Index rating

Nutrient Index	Range	Remarks
A	<1.67	Low
B	1.67-2.33	Medium
C	>2.33	High




Fig. 1. A) Traditional nomenclature of Chikkaballapura district soils B) Soil depth of Chikkaballapura district soils C) Surface soil texture of Chikkaballapura district soils D) Land capability classification of Chikkaballapura district soil

3.2 Nutrient Status in Soils of Chikkaballapura District

3.2.1 Soil pH

On an average, the pH of soils of Chikkaballapura district was found be neutral

(65.15 %) followed by alkaline (24.33%) and acidic (10.52 %) (Table 1 and Fig. 2a). Acidic soil pH were found in Chikkaballapura and Shidlagatta taluk and thereby mean value of soil pH (6.89 and 7.12 respectively) of both taluks were lower compared to other taluks (Table 2). The soil pH of the district is adequate for crop

production [13]. Lower pH values could be as a result of the litter falls that releases organic acids after decomposition and red soil as they are formed by weathering of acid crystalline rocks and due to application of acidic fertilizers (Urea and DAP), the content of essential nutrients like nitrogen, phosphorus, lime is very less in red soils which results in slightly acidic behaviour [14]. The lowest values of pH under the cultivated land also may be due to the depletion of basic cations in crop harvest and drainage to streams in runoff generated from accelerated erosions as reported by Foth and Ellis [15]. Overall, soil pH is adequate to crop cultivation with exception to part of Chikkaballapura and Shidlagatta taluk.

3.2.2 Electrical Conductivity (EC) (dSm^{-1})

The soils of Chikkaballapura district were recorded low (68.09%) to medium (31.65%) range in EC values (Table 1 and Fig. 2b). EC values were ranged from 0.16 to 0.22 dSm^{-1} and lowest was found in Shidlagatta and Chintamani taluk and higher values were found in other four taluks (Table 2). Thus, it is inferred that the soils of cultivated fields of Chikkaballapura district do not suffer from any salt problem as of now. The district has neither major rivers of perennial nature nor any irrigation canals to influence the soluble salt content of the soils in general. The normal EC also may be ascribed to leaching of salts to lower horizons due the light textured nature of the soils [7].

3.2.3 Organic Carbon (OC) (%)

Organic carbon content was found lower (89.64%) in majority soils of the Chikkaballapura district and medium range only in 10.02 % soils (Table 1 and Fig. 2c). Lowest OC was found in Gowribidanuru taluk (0.22%) followed by Bagepalli and Gudibande (0.25%) and Chintamani (0.34%) and medium OC value was found in Chikkaballapura taluk (0.55%) (Table 2). The mean results for 20 samples analysed during 2012-13 and 2022-23 from each taluks were plotted and found that OC values were higher in recent years (Fig. 3), it was recorded 0.15-0.18% in 2012-13 and increased to 0.42-0.51% during 2022-23. Low-level soil OC is a significant challenge for sustainable soil health. The inappropriate cultivation practices, low input of organic manure and crop residues, and the rapid decomposition rate of organic matter might be responsible for inferior organic carbon content in the soils [16]. High temperature and good

aeration in the soil increased the rate of oxidation of organic matter resulting reduction of organic carbon content. The high temperature prevailing in the area was responsible for the rapid burning of organic matter, thus resulting in medium organic carbon content of these soils. These results are in confirmatory with results reported by Sathish et al [17] and [7]. Incorporation of different organic matter in adequate amounts, crop residue management, organic matter management and a balanced fertilizer application will be helpful for the improvement of organic carbon for sustained productivity [18]. Maintaining a certain level of organic carbon is extremely important for soil fertility.

3.2.4 Available Nitrogen (N) (kg ha^{-1})

Likewise OC values, available N status in Chikkaballapura soils were in lower limits (63.63 kg ha^{-1}) and medium in 36.37 % soils (Table 1 and Fig. 2d). Descriptive statistical values for N (Table 2) were found lower in all the taluks due to the available N was found very low during 2012-13 and 2013-14 and later on gradually increased and mean results for 20 samples analysed during 2022-23 from each taluks were clearly shown higher N values (Fig. 3) compared to 2012-13. Soil N values were found 59.58-71.25 kg ha^{-1} in 2012-13 and increased to 62.48-263.24 kg ha^{-1} during 2022-23. However, soil N values for Gudibande and Gowribidanuru taluks were not shown much difference in both years (Fig. 3) but it is increasing trend. The low available nitrogen in most of the soils might be due to the higher temperature in semi arid climate of Chikkaballapura as a result more volatilization which resulted in low status of available nitrogen. Similar results were also reported by [19] in the soils of Krishna district, Andhra Pradesh and [20] in Dumka and Lachinpur series of Jharkhand.

Soil OC and soil Nitrogen have long been identified as factors that are important to soil fertility in both management and natural ecosystem. In addition, soil OC is the main source of energy for soil microbes and therefore, the amount of soil OC will influence the availability of essential plant nutrients. One of these essential nutrients is nitrogen is required in relatively large concentration for plant growth and its availability can limit vegetation distribution [21]. Thereby maintaining the organic carbon status and inclusion of organic resources in agriculture could be supportive in maintaining the soil fertility of Chikkaballapura soil.




Fig. 2. Nutrient status in percentage for the measured soil properties in Chikkaballapura district covering the six taluks (a) pH- 1:2.5 soil : water; b)EC – Electrical conductivity; c)O.C – Organic carbon; d) Available nitrogen as N kg ha⁻¹); e)Available phosphorus as P₂O₅(kg ha⁻¹); f)Available potassium as K₂O(kg ha⁻¹))




Fig. 3. Difference in nutrient status in six taluks of Chikkaballapura district during 2012-13 and 2022-23 (Value are mean of 20 locations samples of each taluks for its respective nutrient parameters of both years)

Table 4. Frequency of soil samples falling in the indices categories, percent (%) of soil samples under each category, and Nutrient index value as well as rating in the Chikkaballapura district covering six taluks based on soil pH, EC, Organic carbon, Available N, P₂O₅, and K₂O

Parameters/Taluk	Number of samples falling in the index category			% of soil sample under each category			Nutrient Index	
	Low	Medium	High	Low	Medium	High	Value	Rating
pH (1:2.5 water ratio)								
Bagepalli	22	164	215	5.49	40.90	53.62	2.48	High
Shidlagatta	467	2891	777	11.29	69.92	18.79	2.07	Medium
Chikkaballapura	231	845	145	18.92	69.21	11.88	1.93	Low
Gudibande	20	82	60	12.35	50.62	37.04	2.25	Medium
Gowribidanuru	7	75	73	4.52	48.39	47.10	2.43	High
Chintamani	401	3053	1385	8.29	63.09	28.62	2.20	Medium
EC(dS m⁻¹)								
Bagepalli	377	23	1	94.01	5.74	0.25	1.06	Low
Shidlagatta	766	3356	13	18.52	81.16	0.31	1.82	Medium
Chikkaballapura	1178	36	7	96.48	2.95	0.57	1.04	Low
Gudibande	158	3	1	97.53	1.85	0.62	1.03	Low
Gowribidanuru	147	7	1	94.84	4.52	0.65	1.06	Low
Chintamani	4805	29	5	99.30	0.60	0.10	1.01	Low
O.C (%)								
Bagepalli	322	21	1	93.60	6.10	0.29	1.07	Low
Shidlagatta	3532	369	18	90.13	9.42	0.46	1.10	Low
Chikkaballapura	1021	96	3	91.16	8.57	0.27	1.09	Low
Gudibande	104	11	0	90.43	9.57	0.00	1.10	Low
Gowribidanuru	119	8	0	93.70	6.30	0.00	1.06	Low
Chintamani	4059	519	12	88.43	11.31	0.26	1.12	Low
N(kg ha⁻¹)								
Bagepalli	391	10	0	97.51	2.49	0.00	1.02	Low
Shidlagatta	351	3784	0	8.49	91.51	0.00	1.92	Medium
Chikkaballapura	108	133	0	89.11	10.89	0.00	1.11	Low
Gudibande	158	4	0	97.53	2.47	0.00	1.02	Low
Gowribidanuru	149	6	0	96.13	3.87	0.00	1.04	Low
Chintamani	149	6	0	99.34	0.66	0.00	1.01	Low
P₂O₅(kg ha⁻¹)								
Bagepalli	378	23	0	94.26	5.74	0.00	1.06	Low
Shidlagatta	3800	335	0	91.90	8.10	0.00	1.08	Low
Chikkaballapura	1111	105	5	90.99	8.60	0.41	1.09	Low
Gudibande	156	0	0	96.30	0.00	0.00	0.96	Low
Gowribidanuru	148	6	1	95.48	3.87	0.65	1.05	Low
Chintamani	4407	425	7	91.07	8.78	0.14	1.09	Low
K₂O(kg ha⁻¹)								
Bagepalli	361	38	2	90.02	9.48	0.50	1.10	Low
Shidlagatta	13	4122	0	0.31	99.69	0.00	2.00	Medium
Chikkaballapura	1111	108	2	90.99	8.85	0.16	1.09	Low
Gudibande	156	0	0	95.68	0.00	0.00	0.96	Low
Gowribidanuru	141	12	2	90.97	7.74	1.29	1.10	Low
Chintamani	2295	2507	37	47.43	51.81	0.76	1.53	Low

(EC – Electrical Conductivity; O.C – Organic Carbon; Available nitrogen as N (kg ha⁻¹); Available phosphorus as P₂O₅ (kg ha⁻¹); Available potassium as K₂O (kg ha⁻¹))

3.2.5 Available phosphorus (P₂O₅) (kg ha⁻¹)

Available P₂O₅ status in Chikkaballapura soils were found lower in 91.63 % and medium range in 8.19 % of the soils (Table 1 and Fig. 2e). As per the critical limits for P₂O₅ content, lower values in all six taluks of the district found in descriptive statistics

(Table 2). Lowest P₂O₅ content was found in Gowridinanuru taluk followed by Gudibande, Bagepalli, Chikkaballapura, Shidlagatta and Chintamani taluk. Trend graph derived for mean values of 20 samples analysed during a decade ago and 2022-23 shown that, P₂O₅ values are higher during recent years and it was ranged from 9.10 – 12.84 kg ha⁻¹ in 2012-13 and

increased to 18.81 – 21.43 kg ha⁻¹ during 2022-23. High and continuous application of phosphatic fertilizers might have resulted in occurrence of high phosphorus soils in the district and such build up in available phosphorus was also noticed in the soils of Amritsar district of Punjab, Haveri district of Karnataka and Coimbatore of Tamil Nadu during fertility mapping by Sharma et al [22] and Padmavathi et al [23], respectively. It is also reported by Chandrakala et al. [24].

3.2.6 Available potassium (K₂O) (kg ha⁻¹)

Majority of the soils of the Chikkaballapura district is in medium range of available K₂O according to critical limits (Table 1 and Fig. 2f). Available K₂O content was found medium in 62.19 % soils and lower in 37.35 % soils and higher during recent years and it was ranged from 68.56 – 90.49 kg ha⁻¹ in 2012-13 and increased to 110.81 – 123.71 kg ha⁻¹ during 2022-23 (Fig. 3). Among the six taluks, lowest K₂O content was found in Gudibande and highest in Chintamani taluk (Table 2). Adequate available K in these soils may be attributed to the prevalence of potassium-rich minerals like Illite and Feldspars [22] and also the variation in available potassium across the soils of different districts was noticed by several workers [25] and was attributed to variation in mineralogical compositions. The low content of K₂O in some areas under consideration might be due to the loose texture of the soil, inappropriate field management practices, and the low use of K fertilizers [26]. It has been observed that the significantly higher amount of K₂O in some soils might be due to the well-practiced fertilizer management schedule [27]. However, the taluks were showing K deficiency needs special care in terms of K fertilizer management.

3.2.7 Assessment of soil fertility based on the Nutrient Index (NI)

Overall nutrient indexes concerning pH, EC, OC and available N, P, K are tabulated in Table 4. Nutrient index analysis for the six taluks of the Chikkaballapura district indicated that the higher and lower under the soil pH category in Bagepalli (53.62%) and Gowribidanuru (4.52%) taluks respectively. The higher per cent of soil samples under medium category is in Shidlagatta taluk (69.92%). The low nutrient index of pH was found in Chikkaballapura taluk, medium in Shidlagatta, Gudibande, Chintamani and high in

Bagepalli and Gowribidanuru taluk. In case of EC values, the higher and lower percent of soils in low category found Chintamani (99.30%) and Shidlagatta (18.52%) respectively. In medium and high per cent category higher EC recorded in Shidlagatta (81.16%) and Gowribidanuru (0.65%) respectively. The NI of EC found medium only in Shidlagatta taluk and low in other five taluks of Chikkaballapura district. However, all the six taluks indicated that lower NI for organic carbon, nitrogen and phosphorus and potassium. The NI of N and K₂O were found medium in Chikkaballapura taluk. With respect to per cent of soil samples category for OC, N, P and K more than 90% soil samples found in lower category with exception to Chikkaballapura taluk for N and K₂O. The soil nutrient index value is a measure of the capacity of soil to supply nutrients to plants which also assesses the long-term impact of different crop-grown systems on changes in nutrient patterns [28,29]. In this study, it was found that most of the examined soils are in low in important parameters such as OC, N, P₂O₅ and K₂O. In this context, the adoption of appropriate soil management strategies would be helpful to improve the low and medium soil nutrient index of the soil.

4. CONCLUSION

Chikkaballapura district soils were detected with neutral in soil pH and normal range in EC values and results depicted that soil is adequate for crop production. Although the results shown improvement over the years, organic carbon status was indicated that requirement for more organic management practices in the soils of all the taluks of the Chikkaballapura district. The fertility assessment makes it possible to understand the major nutrient deficiency in the soils, to overcome this it would be wise for farmers to adopt integrated management approach to sustain the agriculture.

ACKNOWLEDGEMENT

Authors would like to thank all the farmers who were interested in soil testing of their cultivable land. And also extend our acknowledgement to NBSS and LUP for providing maps of Chikkaballapura district which includes soil texture, depth and land capability.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Jackson ML. Soil chemical analysis. Prentice. Hall of India Pvt. Ltd.: New Delhi; 1973.
2. Walkley A, Black CA. An examination of digestion method for determining soil organic matter and the proposed modification of the chromic acid titration method. *Soil Science*. 1934; 37:29–38.
3. Subbiah AY, Asija GK. Available nitrogen: Alkaline permanganate method. *Current Science*. 1949;25:254 – 260.
4. Bray RH, Kurtz LT. Determination of total organic and available forms of phosphorus in soils. *Soil Science*. 1945;59:39-45
5. Baruah TC., Barthakur HP. A Textbook of Soil Analysis. Vikas Publishing House Pvt. Ltd.: New Delhi. 1997:334.
6. Ganeshamurthy AN, Satisha GC, Prakash Patil. Potassium nutrition on yield and quality of fruit crops with special emphasis on banana and grapes. *Karnataka Journal of Agricultural Science*. 2011;24(1):29-38.
7. Siva Prasad PN, Subbarayappa CT, Sathish A. Spatial distribution and mapping of available micro nutrient status in Relation to Soil Properties of Tomato Grown Soils of Chikkaballapura District, Karnataka. 2020;32(10):1-10.
8. Ramachandra TV, Vinay S, Bhargavi R, Bharath H. Integrated watershed management for water and food security in Kolar and Chikkaballapura district, Karnataka. Technical report. 2017:1-74.
9. Available:<https://e-krishiuasb.karnataka.gov.in>
10. Parker FW, Nelson WL, Winters E, Miles JE. The broad interpretation and application of soil test summaries. *Agronomy Journal*. 1951;43(3):103-112.
11. Kumar P, Kumar A, Dyani BP, Shadi UP, Singh SP, Kumar R, Kumar Y, Sumith R, Soil fertility status of some soils of Muzaffarnagar district of Uttar Pradesh, India along with Ganga canal command area. *African Journal of Agricultural Research*. 2013;8(14):1209-1217.
12. Ko TH., Nature and properties of lateritic soil derived from different parameters in Taiwan. *Scientific World Journal*. 2014:2-4.
13. Onwudike SU, Uzoho BU, Ihem EE, Ahukaemere CM, Nkwopara U., Irokwe IF, Echeanyanwu GI. Evaluation of the fertility status of selected soils in Mbaise, Imo State Southeastern Nigeria using nutrient index method. *Agro. Search*. 2016;16(1):79 - 90.
14. Naveen Dv, Nalina CN, Manjunath Gowda KS, Venkataravana P, Venkatachlapathi V, Manjunatha R. Nutrient index approach - A key assessment of soil fertility status of chintamani campus, UAS, Bangalore. *Mysore J. of Agricul. Sci*. 2022:194-202.
15. Foth HD, Ellis BG. *Soil Fertility*, 2nd Edition Lewis CRC Press LLC., USA 1997: 290p.
16. Basak N, Mandal B., Datta A., Kundu NC., Rai AK., Basak P. Stock and stability of organic carbon in soils under major agro-ecological zones and cropping systems of sub-tropical India. *Agricul. Eco. Environ*. 2021:107317
17. Sathish A, Ramachandrappa BK, Devaraja K, Savitha MS, Thimme Gowda MN, Prashanth KM. Assessment of spatial variability in fertility status and nutrient recommendation in Alanatha cluster villages, Kanakapura taluk, Ramanagara district, Karnataka Using GIS Techniques. *Int. J. Curr. Microbiol. App. Sci*. 2017;6(5):211-224.
18. Kanyenji GM, Oluoch-Kosura W, Onyango CM, Ngang'a SK. Prospects and constraints in smallholder farmers' adoption of multiple soil carbon enhancing practices in Western Kenya. *Heliyon* 2020;6(3):e03226.
19. Rajeswar M, Rao CS, Balaguravaiah D, Khan MAA. Distribution of available macro and micronutrients in soils garikapadu of krishna district of Andhra pradesh. *J. Indian Soc. Soil Sci*. 2009;57(2):210-213.
20. Kumar R, Sarkar AS, Singh KP, Agarwal BK, Karmakar S. Appraisal of available nutrients status in Santhal Paraganas region of Jharkhand. *J. Indian Soc. Soil Sci*, 2009;57(3):366-369.
21. Emmanuel A, Louis H, Akakuru OU, Adeola OA, Timothy FT, Amos PI, Banu HD, Ilu D. Assessment of Organic Carbon and Available Nitrogen in the Soil of Some Selected Farmlands Located at Modibbo Adama University of Technology, Adamawa State, Nigeria. *J. Environ. Anal. Chem*. 2018;5:239
22. Sharma PK, Sood A, Setia RK, Tur NS, Mehra D, Singh H. Mapping of macronutrients in soils of Amritsar district (Punjab) A GIS approach. *J. Indian Soc. Soil Sci*. 2008; 56(1):34-41.
23. Padmavathi T, Muthukrishnan R, Mani, Sivasamy R. Assessment of soil physicchemical properties and

- macronutrients status in coimbatore district of Tamilnadu using GIS techniques. Trends in Bio Sci. 2014;7(19):2879-2881.
24. Chandrakala M, Srinivasamurthy CA, Sanjeev K, Bhaskar S, Parama VRR, Naveen, DV. Phosphorus Status in Soils of Eastern Dry Zone, Karnataka, India. 2017: 310-324
 25. Reza SK, Baruah U, Sarkar D. Spatial variability of soil properties in Brahmaputra plains of north-eastern india: A geo statistical approach. J. Indian Soc. Soil Sci. 2012;60(2):108-115
 26. Bagyalakshmi B, Ponmurugan P, Balamurugan A. Potassium solubilization, plant growth promoting substances by potassium solubilising bacteria (KSB) from southern Indian tea plantation soil. Biocatalysis. Agric. Biotechnol. 2017: 12:116–124.
 27. Hu X, Chen J, Guo J. Two phosphate and potassium solubilising bacteria isolated from tianmu mountain, zhejiang, China. World J. Microbiol. Biotechnol. 2006: 22:83–99.
 28. Amara DM, Patil PL, Kamara AM., Saidu, H. Assessment of soil fertility status using nutrient index approach. Academia. J. Agric. Res. 2017:5(2):28–38.
 29. Singh SR, Yadav P, Singh D, Bahadur L, Singh SP, Yadav AS. Impact of different crop grown systems on land nutrient index, microbial diversity and soil quality. Land. Degradation. Dev. 2021:32(14): 3973–3991.

© 2023 Sandhya et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/111114>