

RESEARCH ARTICLE

The relationship between wasting and stunting in Cambodian children: Secondary analysis of longitudinal data of children below 24 months of age followed up until the age of 59 months

Mueni Mutunga^{1*}, Alexandra Rutishauser-Perera², Arnaud Laillou³, Sophonneary Prak⁴, Jacques Berger⁵, Frank T. Wieringa⁵, Paluku Bahwere⁶

1 United Nations Children's Fund (UNICEF) East Asia Pacific Regional Office, Bangkok, Thailand, **2** Action Against Hunger UK, London, United Kingdom, **3** United Nations Children's Fund (UNICEF), Addis Ababa, Ethiopia, **4** National Nutrition Program, Maternal and Child Health Center, Phnom Penh, Cambodia, **5** Institut de Recherche pour le Développement, Montpellier, France, **6** Centre de Recherche en Épidémiologie, Biostatistique et Recherche Clinique, Ecole de santé publique, Université Libre de Bruxelles, Brussels, Belgium

* mmutunga@unicef.org



OPEN ACCESS

Citation: Mutunga M, Rutishauser-Perera A, Laillou A, Prak S, Berger J, Wieringa FT, et al. (2021) The relationship between wasting and stunting in Cambodian children: Secondary analysis of longitudinal data of children below 24 months of age followed up until the age of 59 months. *PLoS ONE* 16(11): e0259765. <https://doi.org/10.1371/journal.pone.0259765>

Editor: Srinivas Goli, University of Western Australia, AUSTRALIA

Received: December 9, 2020

Accepted: October 27, 2021

Published: November 18, 2021

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0259765>

Copyright: © 2021 Mutunga et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The interrelationship between wasting and stunting has been poorly investigated. We assessed the association between two indicators of linear growth, height-for-age Z-score (HAZ) change and occurrence of accelerated linear growth, and selected indicators of wasting and wasting reversal in 5,172 Cambodian children aged less than 24 months at enrolment in the 'MyHealth' study. The specific objectives were to evaluate the relationship between temporal changes in wasting and 1) change in HAZ and 2) episodes of accelerated linear growth. At enrolment, the stunting and wasting prevalence were 22.2 (21.0;23.3) % and 9.1 (8.1;10.1) %, respectively, and reached 41.4 (39.3;43.6) %, and 12.4 (11.5;13.3) % respectively, two years later. Between 14–19% of stunted children were also wasted throughout the whole study period. For each centimetre increase in Mid-Upper Arm Circumference (MUAC) from the previous assessment, the HAZ increased by 0.162 (0.150; 0.174) Z-score. We also observed a delayed positive association between the weight for height Z score (WHZ) unit increase and HAZ change of +0.10 to +0.22 units consistent with a positive relationship between linear growth and an increase in WHZ occurring with a lag of approximately three months. A similar positive correlation was observed for the occurrence of an episode of accelerated linear growth. These results show that interventions to prevent and treat wasting can contribute to stunting reduction and call for integrated wasting and stunting programming.

Data Availability Statement: All relevant data are within the manuscript and its [Supporting Information](#) files.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Wasting, defined by a low weight-for-height Z-score ($WHZ < -2z$) or mid-upper arm circumference ($MUAC < 12.5\text{cm}$), affected 47 million children globally in 2019. Stunting, defined by height-for-age Z-score ($HAZ < -2z$), affected 144 million children [1]. Many countries are not on track to meet the Sustainable Development Goal (SDG) 2.2 targets to reduce the wasting prevalence to below 5% and the number of stunted children by 40% by 2025 compared to 2012 [1–6]. These two forms of undernutrition often coexist in the affected countries and communities as they share similar risk factors [7–10]. The 2014 Cambodia Demographic Health Survey (CDHS) found a prevalence of wasting and stunting among children below five years of age of 10% and 32.5%, respectively [11]. These figures indicate that both conditions are of public health importance in Cambodia [6, 7]. Similar findings have been documented in other countries in the region [6, 9, 12, 13].

Despite the coexistence of wasting and stunting, the relationship between the two conditions has not attracted sufficient interest from the research community, program implementers or policymakers. Previous research focuses on the similarities of the determinants [7], associations of multiple nutritional deficits and mortality risk [7, 12, 14–16] and prevalence of concurrent wasting and stunting [12, 13, 16, 17].

We are aware of only one published peer-reviewed paper that checked whether wasting was a risk factor for stunting occurrence and vice-versa [17]. This study retrospectively analyzed data from well-run Gambian rural growth monitoring clinics collected from 1976 to 2016. It concluded that more attention on the interrelationship between wasting and stunting is required and proposed that both forms of undernutrition be addressed jointly [17].

Currently, wasting and stunting are addressed through separate programmes with limited integration [18–22]. Due to the separation between wasting and stunting programming, the effect of weight replenishment on linear growth has not been examined. This is despite hospital data since the 80's suggesting a link between the level of weight replenishment and linear catch-up growth in children recovering from wasting, with catch up growth only commencing when wasting has been addressed [23, 24]. Unfortunately, the separation of wasting and stunting programming is unlikely to change unless more evidence is produced. In this study, we analyzed longitudinal data from three provinces in Cambodia. The specific objectives of this study were to evaluate the relationship between temporal changes in wasting and 1) change in height-for-age and 2) episodes of accelerated linear growth. We aimed to answer two key research questions: 1) Are changes in indicators of wasting (WHZ and $MUAC$) associated with a change in HAZ velocity over time?, and, 2) Are the presence or absence of the same wasting parameters associated with the occurrence of accelerated linear catch-up growth in subsequent follow-up visits?

Methods

Study design and participants

This paper is based on a secondary analysis data from the “MyHealth” open cohort intervention study, which took place between February 2016 and August 2018 in six districts from three provinces in Cambodia; Russei Kaev in Phnom Penh, Chitr Borie and Krong Kratie in Kratie province, and Ou Chum, Krong Ban Lung, and Bar Kaev in Ratanakiri province. The provinces represent different population groups from Cambodia. The district in Phnom Penh concentrates a poor suburban population being from a diversity of ethnicities and religions. Kratie, a province crossed by the Mekong river, is inhabited by a population consisting of ethnic minorities that is highly reliant on agriculture. Ratanakiri province is concentrating ethnic

groups practicing subsistence agriculture and food collection from wild sources. Detailed methods of the study have been previously published [25–28]. The study enrolled all women from villages sampled from the six districts who were pregnant or lactating at any of the data collection rounds and surveyed them prospectively for over three years. In the same communities, children below three years were enrolled in the MyHealth study at the first round of data collection (baseline). For this paper, we have analyzed data of children who were less than 24 months at enrolment as they are the focus of most interventions aiming at tackling stunting. Hence, we excluded children aged ≥ 24 months at recruitment and those children who were surveyed only once. Subsequently, infants born from participating pregnant women were enrolled at the nearest data collection round after their birth. After recruitment, children were surveyed every 3 to 4 months during the first two years of the study and at six months intervals thereafter. Children exited the study when they reached five years or at the end of the study depending on whichever occurred first.

The measurements were taken by 8 teams of surveyors who received trained on anthropometric measurements for up to a maximum of 5 days for each round of data collection. Weight, length or height, and mid-upper arm circumference (MUAC) were measured in duplicates for each child, and the mean values were further used. The measurement tools were calibrated after each fifth measurement. A spot check team monitored the data collection teams' measurements techniques and calibration of the tools.

Source of data

Anonymized data were extracted from the database of the “MyHealth” study described above, including; study identifier (child, household and mother identities); administrative information (province, district and date of the survey for the different data collection points); household characteristics (head of household level of education, mother level of education, number of people in the household, number of children below five years in the household, number of children below 15 years, type of toilet facility, type of source used to get drinking water); child socio-demographic characteristics (sex, birth date, and age at the different data collection points); and child nutrition parameters (weight, height, MUAC, weight-for-age Z-score (WAZ), HAZ and WHZ for the different data collection points).

Analysis

Variable transformation and definitions. “MyHealth” study was a longitudinal open cohort design whereby children entered and exited the cohort at any of the follow up visits. Data were reorganized to ensure that the recruitment visit became ‘Visit 0’ for all children, regardless of the data collection round at which the children were recruited into the study. Subsequent visits were labelled as follow up visits.

The standard cut-off of -2 Z-score of the multiple countries' growth standards reference was used to define the nutrition status of children. Height-for-age Z-score (HAZ) < -2 were classified as stunting and weight-for-height Z-score (WHZ) < -2 as wasting [29].

Household and child characteristics were compared between children included and excluded from the analysis to evaluate potential selection bias.

Based on the age categories used by the Demographic Health Surveys (DHS), We defined the following three age categories: 0–5 months, 6 to 11 months and 12 to 23 months. We assumed that the level of education of both the mothers and heads of households (HHH) did not vary during the study period. Information from the follow-up visits (FV) were used to fill any missing variables data collected at recruitment. The following education level groups were defined: 0 to 6 years of education, 7 to 9 years and ≥ 10 years for the mothers, and 0 years, 1 to

6 years, 7 to 9 years and ≥ 10 years head of household (HHH). Household (HH) size was defined in four groups: < 4 people, 4 to 5 people, 6 to 10 people and ≥ 11 people.

We created the variables 'HAZ increment' (HAZd), 'WHZ increment' (WHZd) and 'MUAC increment' (MUACd) by calculating the absolute difference between two consecutive visits of HAZ, WHZ and MUAC, respectively. We also created the binary variables 'Ever had acute malnutrition' and 'Ever had an episode of accelerated linear growth' to distinguish those who experienced the condition at any data collection point. Accelerated linear growth was defined using the cut-off proposed in the literature of $+0.67$ HAZ increase between two data collection points [30–32]. Based on this cut-off, a child who had an HAZd ≥ 0.67 was considered as having experienced an accelerated linear growth [30].

We used the previous follow-up period to indicate the interval between the two last data collection points (visits) and the current follow-up period to indicate the previous and current FV interval period.

Outcomes of interest and sample size. The primary outcomes of interest were HAZ change and accelerated linear growth as defined by HAZd ≥ 0.67 . Secondary outcomes of interest were the prevalence of stunting, the prevalence of wasting, and the prevalence of concurrence of wasting and stunting (WaSt).

There was no specific calculation of sample size for all the outcomes of interest for this paper. The sample size was calculated based on the main objective of the original study. We assumed that a sample of 1200 children below two years of age per region was necessary for the study to have sufficient power to demonstrate at least a 6% reduction in stunting prevalence over a three-year period. Children aged ≥ 24 months at recruitment and children who were surveyed only once were excluded from our sample.

Data management and analysis. The data extracted from the "MyHealth" study database was exported into excel files. These files were then converted to STATA format and merged using STATA software version 14.1. Data cleaning, restructuring, missing values analysis, new variables creation and data analysis were performed using the same STAT 14.1 statistical package. The restructuring of the data consisted of arranging the data in order to have all recruitment (first assessment) data positioned at visit (V0) for children first surveyed at Follow up Visits one (FV1) to six (FV6). The analysis of missing values was done to determine the extent of missing information for each variable and identify time-varying variables for which data were not collected systematically in all the data collection time points. This analysis allowed the exclusion of variables not routinely collected and those associated with a considerable reduction in sample size in multivariate analyses. No missing values handling approach was used in our analysis as we assumed that for all the variables retained, the data were missing completely at random and opted for the complete cases analysis approach [33, 34].

We used standard statistics for quantitative analyses. Continuous variables were described using means and their 95% confidence interval (CI) and compared using the Student t-test. Categorical variables were described using proportion and their 95%CI and compared using Chi-square tests. To test for the linearity of the trend over time, we calculated a chi-square statistic for the trend using the STATA ptrend command [35]. We determined all the effect sizes, including prevalence, mean difference, univariate and adjusted odds ratio (OR) using multi-level linear or logistic mixed-effect modelling as appropriate to account for clustering at the regional level. The covariates included in the full initial multilevel linear or logistic regression mixed-effects models were selected based on their relevance according to the literature and availability in the received dataset]. We did not use stepwise methods to select the covariates in the multivariate analysis to determine the final models but excluded the non-significant covariates manually, as proposed by Greenland [36]. For fitting the mixed-effects models, we implemented the panel data analysis and used the STATA standard command Mixed and melogit as

appropriate [37, 38]. Based on the authors' observation from several countries on catch up growth after a hunger season or during recovery from an episode of wasting, linear catch-up growth occurs around three months after that of weight, starting only when the weight deficit is almost completely replenished. Therefore, we also conducted repeated cross-sectional modelling that allowed the introduction of one follow up the period lag in our models assessing the association between wasting and stunting [23, 39–41]. This corresponded to a time lag of approximately 4-months and meant that the covariates wasting and WHZ at FV1 were included in the model assessing the association between stunting and wasting at FV2. Such time lag was not introduced for MUAC as we did not have similar evidence. For this alternative modelling approach, we also used the standard stata command mixed and melogit. The results of this second modelling approach are mostly presented in S1 and S2 Tables.

Ethical considerations

Ethical approval for the “MyHealth” study was granted by the Cambodia National Ethics Committee for Health Research (NECHR), National Institute of Public Health, Ministry of Health, Cambodia (number 117/NECHR). All candidates were informed in their local languages about study objectives and procedures, the voluntary aspects of participation, the possibility to withdraw consent at any given points, the research team's obligation to preserve the participants' privacy, and the use of data for scientific data publications. All participants provided written informed consent at baseline before their inclusion in the original MyHealth study. Consent was obtained from adults primary caregivers (mostly mothers) for participating children. Community health volunteers witnessed the entire participation approval process. The authorization to use the anonymized “MyHealth” study data for secondary data analysis and publication was granted by the relevant authority from the Cambodian Ministry of Health. The need for a new ethical approval for the current study was waived by the NECHR of the Ministry of Health.

Results

Study participants

Of the 5172 children included in this analysis, 70.3% ($n = 3635$) were recruited at baseline, 5.8% ($n = 304$) at follow up visit 1 (FV1), 5.8% ($n = 298$) at FV2, 2.9% ($n = 151$) at FV3, 4.1% ($n = 210$) at FV4 and 11.1% (574) at FV5. The median number of survey rounds for which the children were surveyed was 5 (interquartile range:3–6); Only 13.3% ($n = 686$) were surveyed in all the seven rounds.

Table 1 presents households, demographic and nutritional and child characteristics of children included in this analysis, based on data collected during recruitment into the study. About 40% of children were recruited in the Kratie region and about 30% in the other two areas. Most mothers (64.5%) and heads of households (60.1%) had between zero and six years of formal education. The average household size was about six people per household. Over three quarters of the households had an improved source of drinking water, while improved sanitation was available for less than 60% of the households. The great majority of children used mosquito nets. Male and female were equally represented, and the vast majority were less than 12 months of age at recruitment. One out of 10 children had a birth weight of less than 2.5kg. The median duration of exclusive breastfeeding was two months only, and the maximum duration reported was four months. A significant proportion of the children were underweight, stunted and wasted at recruitment into the cohort (Table 1).

Overall, 5.8% (299/5172) of children had received treatment for Severe Acute Malnutrition (SAM) during the follow-up period as reported by their caregivers. The proportion of children

Table 1. Households and children characteristics.

Characteristics	n/N	%	Average
Households			
Region			
Phnom Penh	1448/5172	28.0	
Kratie	2051/5172	39.7	
Ratanakiri	1673/5172	32.3	
Mother formal education			
0–6 Years	3270/5069	64.5	
7–9 Years	1227/5069	24.2	
≥10 Years	572/5069	11.3	
HHH formal education			
0 Year	908/3878	23.4	
1–6 Years	1423/3878	36.7	
7–9 Years	950/3878	24.5	
≥10 Years	597/3878	15.4	
Household size (n people)			
Mean number of people (SD ¹)			5.8 (2.8)
<4	698/4219	16.5	
4–5	1695/4219	40.2	
6–10	1568/4219	37.2	
≥11	258/4219	6.1	
Drinking water source			
Improved	4070/5172	78.7	
Sanitation			
Improved	2911/5158	56.4	
Mosquito net use			
Yes	2256/3108	72.6	
Sometimes	632/3108	20.3	
No	220/3108	7.1	
Children			
Child gender			
Girls	2599/5172	50.2	
Boys	2573/5172	49.8	
Birth weight(kg)			
Mean (SD)			3.0(0.5)
<2.5 kg	326/3204	10.2	
≥2.5kg	2878/3204	89.8	
Age at recruitment			
Mean (SD)			9.7(6.8)
0–5 months kg	2004/5172	38.8	
6–11 months	1284/5172	24.8	
12–23 months	1884/5172	36.4	
≥ 24 months	0/5172		
EBF ² duration (months)			
Median (IQR ³)	5129		3(2–3)
Weight at recruitment			
Mean (SD)	5109		7.2(2.0)
Height at recruitment			

(Continued)

Table 1. (Continued)

Characteristics	n/N	%	Average
Mean (SD)	5093		67.7(9.3)
Weight-for-age Z-score			
Mean (SD)	5102		-1.1(1.1)
% Z-score < -2	1039/5102	20.4	
Height-for-age Z-score			
Mean (SD)	5075		-1.0(1.3)
% Z-score < -2	1026/5075	20.2	
Weight-for-height Z-score			
Mean (SD)	5074		-0.7(1.1)
% < -2 Z-score	5074		
MUAC ⁴ (cm)			
Mean (SD)	5051		13.4(1.3)
<11.5	367/5051	7.3	
11.5–12.4	736/5051	14.6	
≥12.5	3948/5051	78.1	

¹Wasted = weight-for-length/height Z-score < -2 (2006 WHO reference curves);

²Stunted = Length/height-for-age Z-score < -2 (2006 WHO reference curves);

³WaSt = Concurrently wasted and stunted;

⁴Mean = Standard deviations;

⁵CI = confidence interval;

⁶Visit 0 = recruitment into the cohort visit;

⁷FV = follow up visit;

⁸All age group cohorts combined;

⁹Both sex combined.

<https://doi.org/10.1371/journal.pone.0259765.t001>

for whom treatment of SAM between 2 follow-ups was reported was 1.8% (81/4489), 1.0% (33/3359), 2.0 (61/3105), 5.4% (107/1977) and 4.6% (69/1492) for FV1, FV2, FV3, FV5 and FV6, respectively. This also shows that some children had been on SAM treatment for more than one period.

Trends in the prevalence of wasting, stunting, and wasting and stunting concurrence

The regional weighted prevalence of wasting, stunting and wasting and stunting concurrence (WaSt) are presented in Table 2. Prevalence of wasting was always close or above 10%, with the highest prevalence observed at recruitment and the lowest at FV4. No linear trend was observed from recruitment to FV6 for wasting prevalence. For stunting, the prevalence was just over 20% at recruitment and linearly increased over time (p for linear trend < 0.001), with the highest prevalence being observed at FV5. For WaSt, the prevalence was below 5% at recruitment into the cohort but increased linearly over subsequent follow-up visits (p for linear trend < 0.001), crossing the 5% cut-off at FV 3.

Boys had a higher prevalence of wasting than girls at recruitment and at FV1 to FV4 but not at FV5 and FV6. The differences were statistically significant at recruitment [95% CI = 2.8 (1.0; 4.6) %; p = 0.002] and FV2 [95% CI = 4.0 (2.1; 6.0) %; p < 0.001]. Boys also had a higher prevalence of stunting than girls from recruitment to FV4, but the difference was statistically significant.

Table 2. Prevalence of wasting, stunting and concurrence of wasting and stunting at the different rounds of data collection by gender and age.

Data collection round	n	Age (m) ⁴	Wasted ¹ % (95%CI ⁵)	Stunted ² % (95%CI)	WaSt ³ % (95%CI)
All age groups & sex					
Visit 0 ⁶	5174	9.6 (6.7)	12.4 (11.5–13.3)	22.2 (21.0–23.3)	4.3 (3.8–4.9)
FV ⁷ 1	4447	14.2 (7.0)	10.0 (9.1–10.9)	27.8 (26.5–29.2)	4.5 (3.9–5.1)
FV2	3357	17.4 (6.7)	9.6 (8.6–10.6)	32.2 (30.6–33.8)	4.7 (4.0–5.4)
FV3	2955	21.8 (6.6)	11.2 (10.0–12.3)	35.6 (33.8–37.3)	6.4 (5.5–7.3)
FV4	3087	26.8 (6.6)	9.1 (8.1–10.1)	36.7 (35.0–38.4)	5.3 (4.5–6.1)
FV5	2041	38.0 (7.0)	10.4 (9.1–11.8)	41.4 (39.3–43.6)	5.9 (4.8–6.9)
FV6	1496	47.5 (6.7)	11.4 (9.7–13.0)	36.5 (34.1–39.0)	5.4 (4.2–6.5)
Boys⁸					
Visit 0	2573	9.7 (6.8)	13.4 (10.6–16.3)	22.6 (16.1–29.1+)	4.6 (3.4–5.8)
FV1	2195	14.3 (7.0)	10.5 (9.2–11.8)	30.7 (28.8–32.6)	5.1 (4.1–6.0)
FV2	1670	17.5 (6.7)	11.5 (10.0–13.1)	33.5 (31.3–35.8)	6.0 (4.8–7.1)
FV3	1456	21.8 (6.7)	11.6 (10.0–13.3)	37.2 (35.0–39.7)	7.1 (5.8–8.4)
FV4	1520	26.9 (6.7)	9.3 (7.8–10.7)	37.7 (35.3–40.2)	5.4 (4.3–6.6)
FV5	1010	37.9 (7.0)	10.1 (8.3–12.0)	40.9 (37.8–43.9)	5.7 (4.3–7.1)
FV6	733	47.5 (6.7)	10.2 (8.0–12.4)	35.9 (32.4–39.4)	5.2 (3.6–6.8)
Girls⁸					
Visit 0	2599	9.5(6.7)	10.6 (7.8–13.5)	17.6 (11.1–24.1)	3.3 (2.1–4.5)
FV1	2252	14.1 (7.0)	9.6 (8.4–10.8)	25.1 (23.3–26.9)	4.0 (3.1–4.8)
FV2	1687	17.4 (6.8)	7.7 (6.4–9.0)	30.9 (28.7–33.2)	3.4 (2.6–4.3)
FV3	1499	21.8 (6.7)	10.8 (9.2–12.4)	34.0 (31.6–36.4)	5.7 (4.5–6.9)
FV4	1567	26.7 (6.5)	8.9 (7.5–10.3)	35.7 (33.3–38.1)	5.2 (4.1–6.3)
FV5	1031	38.2 (7.1)	10.8 (8.8–12.7)	42.0 (38.9–45.0)	6.0 (4.6–7.5)
FV6	763	47.4(6.8)	12.5 (10.1–14.9)	37.2 (33.7–40.6)	5.6 (3.9–7.2)
Age 0–5 months⁹					
Visit 0	2004	2.9 (1.6)	7.4 (6.3–8.6)	13.0 (11.5–14.5)	0.7 (0.3–1.1)
FV1	1742	8.0 (3.6)	7.2 (6.0–8.5)	18.0 (16.2–19.8)	1.9 (1.2–2.5)
FV2	1386	12.0 (4.5)	9.2 (7.7–10.7)	26.4 (24.1–28.7)	3.6 (2.6–4.6)
FV3	1149	16.3 (4.5)	13.5 (11.5–15.4)	33.0 (30.2–35.7)	7.8 (6.2–9.3)
FV4	1061	21.7 (5.7)	10.2 (8.4–12.0)	37.0 (34.1–39.9)	6.7 (5.1–8.2)
FV5	641	31.3 (3.6)	13.5 (10.8–16.1)	44.7 (40.8–48.6)	8.4 (6.2–10.5)
FV6	417	39.5 (2.2)	14.9 (11.4–18.3)	41.5 (36.8–46.3)	8.1 (5.4–10.7)
Age 6–11 months⁹					
Visit 0	1284	8.8 (1.7)	11.4 (9.5–13.2)	20.8 (18.6–23.1)	3.0 (2.1–4.0)
FV1	1123	13.5 (3.6)	12.8 (10.9–14.8)	29.3 (26.6–31.9)	5.6 (4.2–6.9)
FV2	784	16.5 (3.2)	11.6 (9.3–13.8)	32.3 (29.0–35.6)	5.6(4.0–7.2)
FV3	712	20.2 (2.7)	11.3 (8.9–13.6)	36.4 (32.9–40.0)	7.2 (5.3–9.1)
FV4	764	24.0 (2.2)	8.4 (6.4–10.4)	35.9 (32.4–39.3)	5.4 (3.8–7.1)
FV5	508	35.8 (3.7)	8.8 (6.3–11.3)	41.8 (37.5–46.1)	5.1 (3.0–7.2)
FV6	402	45.2 (2.1)	10.5 (7.5–13.5)	37.1 (32.4–41.9)	5.3 (3.1–7.5)
Age 12–23 months⁹					
Visit 0	1884	17.4 (3.3)	18.5 (16.7–20.3)	33.0 (30.8–35.1)	9.1 (7.8–10.4)
FV1	1582	21.5 (4.2)	11.2 (9.6–12.8)	37.8 (35.4–40.2)	6,7 (5.4–7.9)
FV2	1187	24.4 (3.5)	8.8 (7.2–10.4)	39.2 (36.4–42.0)	5.4 (4.1–6.7)
FV3	1094	28.6 (3.5)	8.7 (7.1–10.4)	37.5 (34.8–40.6)	4.4 (3.2–5.7)
FV4	1262	32.5 (3.7)	8.5 (6.9–10.0)	36.9 (34.2–39.6)	4.0 (2.9–5.1)

(Continued)

Table 2. (Continued)

Data collection round	n	Age (m) ⁴	Wasted ¹ % (95%CI ⁵)	Stunted ² % (95%CI)	WaSt ³ % (95%CI)
FV5	892	44.3 (4.6)	9.1 (7.2–11.0)	38.8 (35.5–42.0)	4.4 (3.0–5.7)
FV6	677	53.7 (3.6)	9.6 (7.4–11.9)	33.1 (29.6–36.7)	3.7 (2.3–5.2)

¹Wasted = weight-for-length/height Z-score < -2 (2006 WHO reference curves);

²Stunted = Length/height-for-age Z-score < -2 (2006 WHO reference curves);

³WaSt = Concurrently wasted and stunted;

⁴Mean = Standard deviation,

⁵CI = confidence Interval;

⁶Visit 0 = recruitment into the cohort visit;

⁷FV = follow up visit;

⁸All age group cohorts combined;

⁹Both sex combined.

<https://doi.org/10.1371/journal.pone.0259765.t002>

The highest prevalence of wasting was observed at 39.5 (SD = 2.2) months of age for the 0 to 5 months age group cohort (FV6), at 13.5 (SD = 3.6) months for the age group cohort 6 to 11 months (FV1) and 17.4 (SD = 3.3) months for the age group cohort 11 to 23 months at recruitment (V0). For stunting, the highest prevalence was observed at the mean age of 31.3 (3.6) months for the age group cohort 0 to 5 months (FV5), 35.8 (3.7) months for the age group cohort 6 to 11 months (FV5) and 24.4 (3.5) months for the age group cohort 12 to 23 months (FV2).

The analysis of trends showed that the prevalence of wasting increased linearly for age group cohort 0 to 5 months [slope(SE) = 1.3 (0.1) %; $p < 0.001$] and age group cohort 6 to 11 months [slope(SE) = 0.3 (0.1) %; $p = 0.027$], while it decreased linearly for the age group cohort 12 to 23 months [slope(SE) = -1.4 (0.1) %; $p < 0.001$]. For stunting prevalence, there was a significant linear trend for the age group cohort 0 to 5 months [slope(SE) = 5.7 (0.2) %; $p < 0.001$] and 6 to 11 months age group cohort [slope(SE) = 3.1 (0.2) %; $p < 0.001$], but not for the 12 to 23 months age group cohort [slope(SE) = 0.2 (0.2) %; $p = 0.163$]. The proportion (95%CI) of stunted children who were WaSt did not show any linear trend.

Association between wasting parameters (presence of wasting, WHZ change, MUAC change) and HAZ change

At FV1, non-wasted children ($n = 3829$) had a smaller decrease in HAZ than wasted children ($n = 813$) at recruitment: [-0.23 (0.75) for the non-wasted versus -0.44 (0.90) for the wasted; [95% CI = 0.21 (0.14; 0.28); $P < 0.001$]. A similar relationship was observed at FV2 [-0.15 (0.57) for the non-wasted ($n = 2667$) at FV1 versus -0.23 (0.63) for the wasted ($n = 260$) at FV1 [95% CI = 0.08 (0.01, 0.15); $p = 0.031$]; FV3 [-0.10 (0.52) for the non-wasted ($n = 2199$) versus -0.15 (0.49) for the wasted ($n = 228$) at FV2 [95% CI = 0.05 (-0.01; 0.12); $p = 0.190$]; and FV4 [-0.06 (0.46) for the non-wasted ($n = 2143$) versus -0.11 (0.51) for the wasted ($n = 224$) at FV3 [95% CI = 0.05 (-0.01; 0.12); $p = 0.111$]. The observed difference was not significant for FV3 and FV4. For FV5 the relationship observed was inverse to that observed at FV1 to FV4 but not significant [-0.07 (0.57) for the non-wasted ($n = 1611$) versus -0.05 (0.40) for the wasted ($n = 162$); at FV4 [95% CI = -0.02 (-0.11; 0.07); $p = 0.697$]. At FV6, in both non-wasted and wasted at FV5, there was an increase rather than a decrease in HAZ suggesting a catch up in linear growth in both groups, with the magnitude being significantly higher in the non-wasted

children: [0.13 (0.41) for the non-wasted (n = 963) versus 0.04 (0.33) for the wasted (n = 110) at FV5 [95% CI = 0.09 (0.01; 0.16); p = 0.033].

Fig 1 below shows the change in HAZ over time for children categorized into four groups: no anthropometric deficit (good), only wasted, only stunted, and wasted and stunted. At all the FVs, the most significant decrease in HAZ was observed for children who were only wasted at the previous visit (Fig 1). In contrast, some catch up in linear growth (increase in HAZ) was observed for children with no anthropometric deficit, only stunted and wasted and stunted, with the largest catch up always observed for children only stunted (Fig 1). The difference in change of HAZ between those with no anthropometric deficit and children only wasted varied between -0.5 and -0.30 Z-score, with the smallest difference observed at FV3 and the largest difference observed at FV1. For the comparison between those with no anthropometric deficit and those who were only stunted at the previous follow up visit (FV), the observed HAZ differences varied between +0.15 Z-score at FV6 and +0.51 Z-score at FV1. The differences between children with no anthropometric deficit and those who had both wasting and stunting at the previous FV varied between 0.0 Z-score at FV6 and +0.31 Z-score observed at FV1. The observed differences were all significant (p<0.05) except for the comparison of those without the anthropometric deficit and wasted at FV3 and FV6, and for those without a deficit and the wasted and stunted children at FV6.

Further, multilevel multivariate longitudinal analysis adjusting for socio-demographic and economic status at baseline (age group, sex, region of residence, mother level of education, type of drinking water source, type of toilet facility), anthropometric characteristics at baseline *MUAC group, HAZ group and WHZ group) MUAC <12.5 cm or not, and the presence or absence of stunting at the previous visits showed that an increase in MUAC between two follow-up visits was associated with an increase in HAZ, while the inverse was observed for WHZ




Fig 1. Linear growth velocity for non-wasted/non-stunted, only wasted, only stunted and wasted and stunted children at the different follow-up visits.

<https://doi.org/10.1371/journal.pone.0259765.g001>

(Table 3). Also, having a MUAC \geq 12.5 cm was associated with a positive change in HAZ, while non-wasted children had, on average, a decrease in HAZ (Table 3).

The analysis per follow up visit (repeated cross-sectional analysis), allowing the introduction of the previous wave trend in WHZ in the multivariate model, showed a delayed positive association between WHZ change and HAZ change. The effect size of the WHZ change during the previous follow up period on change in HAZ in the subsequent follow period varied across the FVs from +0.10 to +0.22 HAZ units, consistent with a positive relationship between linear growth and an increase in WHZ (S1 Table). Similarly, the change in MUAC in the preceding 4 months was positively associated with linear growth across all the FVs, with the effect size varying between +0.04 and +0.14 HAZ units (S1 Table).

Table 3. Factors associated with height-for-age Z-score change overtime (N = 14226 and number of included children = 4629).

Variables	Univariate analysis ¹			Multivariate analysis ¹		
	Coef ²	(95%CI ³)	p	Coef ²	(95%CI ³)	p-value
Baseline variables						
Sex of the child						
Boys/Girls	0.026	(0.007; 0.046)	0.008	0.019	(0.01; 0.038)	0.041
Age at VO (months)						
0-5/12-23	-0.204	(-0.226; -0.181)	<0.001	-0.237	(-0.261; -0.214)	<0.001
6-11/12-23	-0.112	(-0.137; -0.087)	<0.001	-0.091	(-0.114; -0.067)	<0.001
Mother formal education						
7-9 years/0-6 years	-0.006	(-0.031; 0.018)	0.618	0.010	(-0.012; 0.033)	0.378
\geq 10 years/0-6 years	-0.001	(-0.032; 0.031)	0.963	0.034	(0.005; 0.064)	0.0023
Toilet type at V0						
Improved /Unimproved	-0.064	(-0.085; -0.082)	<0.001	-0.062	(-0.084; -0.040)	<0.001
Water source type at V0						
Improved/ Unimproved	-0.048	(-0.072; -0.024)	<0.001	-0.031	(-0.055; -0.008)	<0.001
MUAC⁶ category at V0						
MUAC \geq 12.5 cm/MUAC<12.5 cm	0.031	(0.007; 0.055)	0.012	0.008	(-0.018; 0.035)	0.539
Wasted at V0						
No WHZ ⁷ \geq -2/Yes: WHZ ⁷ < -2	-0.010	(-0.039; 0.019)	0.500	-0.063	(-0.094; -0.033)	<0.001
Stunted at V0						
No HAZ ⁸ \geq -2/Yes: HAZ ⁸ < -2	-0.230	(-0.250; -0.211)	<0.001	-0.167	(-0.187; -0.148)	<0.001
Time varying variables						
MUAC change between visits (cm)	0.073	(0.061; 0.085)	<0.001	0.162	(0.150; 0.174)	<0.001
WHZ change between visits	-0.189	(-0.203; -0.176)	<0.001	-0.279	(-0.293; -0.266)	<0.001
MUAC group						
MUAC \geq 12.5 cm/MUAC<12.5 cm	0.132	(0.089; 0.175)	<0.001	0.125	(0.081; 0.169)	<0.001
Wasted						
No Z-score \geq -2/Yes: Z-score < -2	-0.046	(-0.080; -0.013)	0.006	0.038	(0.003; 0.074)	0.034

¹Computed using multilevel regression analysis test (stata mixed command);

²Coef = coefficient;

³CI = confidence Interval;

⁴HAZ = Length/height-for-age Z-score (2006 WHO reference curves); stunted if HAZ < -2;

⁵V0 = recruitment into the cohort visit;

⁶MUAC = Mid-Upper Arm Circumference ⁷WHZ = weight-for-length/height Z-score (2006 WHO reference curves); wasted = yes if WHZ < -2; ⁸HAZ = Height/length-for-Age Z-score (2006 WHO reference curves); stunted = yes if WHZ < -2.

<https://doi.org/10.1371/journal.pone.0259765.t003>

Association between wasting parameters (presence of wasting, WHZ change, MUAC change) and accelerated linear growth

A total of 2965 episodes of accelerated linear growth, defined by an increase of ≥ 0.67 between two follow up visits, were observed during the study period representing 17.2% (2965/17212) of the measurements. This prevalence varied from 1.98% to 6.63% across FVs, with the highest observed at V0 and the lowest at FV4 (S2 Table). Close to half of the children had at least one episode of accelerated linear growth [95% CI = 47.4 (46.1; 48.8) %]. The proportion of children who experienced at least one episode of accelerated linear growth was higher among children already stunted at V0 (54.2% [51.0; 57.4%]) versus 45.6% [43.8; 47.5%] for those not stunted at recruitment; Δ (95%CI) = 8.6 (5.2–12.0) %; $p < 0.001$].

The univariate and multivariate associations between accelerated linear growth and the presence of wasting as measured by WHZ and MUAC, presence of stunting at V0, change in WHZ, and change in MUAC are shown in Table 4. A cm increase in MUAC between follow up visits more than doubled the likelihood of occurrence of accelerated linear growth, while an increase of 1 unit of WHZ more than halved this likelihood (Table 4). Overall, MUAC group and wasting status at FVs were not independently associated with the occurrence of accelerated linear growth (Table 4). Non-wasted children at baseline were less likely to experience an accelerated linear growth than those who were wasted (Table 4). Similarly, children not-stunted enrolment were less likely to have accelerated linear growth during the study period (Table 4). The cross-sectional analyses allowing the inclusion of ante-previous status showed that having a WHZ < -2 at the ante-previous follow-up visit was consistently associated with a reduced likelihood of experiencing an accelerated linear growth (S2 Table). For WHZ change after FV1, an increase of WHZ by one unit during the ante-previous was associated with a statistically significant two to three-fold increase in the likelihood of experiencing an episode of accelerated linear growth (S2 Table).

Discussion

Summary of findings

Firstly, the findings of this study confirm the high prevalence of stunting among Cambodian infants and young children. The prevalence was high at recruitment into the cohort and increased over time in both sexes. Secondly, our findings show a strong association between the presence of wasting, WHZ change and MUAC change on subsequent linear growth differences and the likelihood of experiencing accelerated linear growth. This finding shows that wasting parameters predicted linear growth.

Interpretation of the findings

Studies conducted in Africa have recently demonstrated a relationship between an episode of wasting and the occurrence of stunting a few months later [18, 25]. To the best of our knowledge, there has not been any study demonstrating an interrelationship between change in wasting parameters and linear growth during childhood. Our study is the first to demonstrate that correction of wasting may induce catch up linear growth a few months later. This conclusion is strengthened by the observed strong positive association between HAZ gain and the indicator of weight replenishment used in our analysis, WHZ and MUAC gain, which has not previously been reported among children followed up prospectively.

Despite experiencing, a slow decline in the prevalence of undernutrition in all its forms, Cambodia [42] is still classified as one of the countries with a very high level of stunting in the region [43, 44]. Our study showed an unchanged prevalence of stunting for the same age groups

Table 4. Association between stunting, wasting parameters and occurrence of accelerated linear growth in the 4630 children included in the analysis (n = 14340 & n episodes = 2965).

Variables	Univariate analysis ¹			Multivariate analysis ¹		
	OR ²	(95%CI ³)	p	OR ²	(95%CI ³)	p-value
Baseline variables						
Sex of the child						
Boys/Girls	1.22	(1.03; 1.45)	0.018	1.21	(1.08; 1.35)	0.001
Age at V0 (months)						
0-5/12-23	1.97	(1.67; 2.32)	<0.001	1.29	(1.10; 1.52)	0.001
6-11/12-23	1.10	(0.83; 1.45)	0.518	1.12	(0.76; 1.64)	0.573
Mother formal education						
7-9 years/0-6 years	0.97	(0.79; 1.18)	0.736	1.03	(0.82; 1.30)	0.786
≥10 years/0-6 years	0.97	(0.78; 1.21)	0.807	1.21	(0.92; 1.59)	0.180
Toilet type at V0						
Improved /Unimproved	1.30	(1.01; 1.68)	0.045	1.10	(0.83; 1.45)	0.494
Water source type at V0						
Improved/ Unimproved	1.21	(0.81; 1.82)	0.350	1.20	(0.64; 2.23)	0.570
MUAC⁶ group at V0						
MUAC≥12.5 cm/MUAC<12.5 cm	1.73	(1.46; 2.06)	<0.001	1.10	(0.84; 1.45)	0.471
Wasted at V0						
No WHZ ⁷ ≥-2/Yes: WHZ ⁷ <-2	0.73	(0.64; 0.84)	<0.001	0.91	0.82; 1.00)	0.058
Stunted at V0						
No HAZ ⁸ ≥-2/Yes: HAZ ⁷ <-2	0.52	(0.36; 0.74)	<0.001	0.54	(0.37; 0.80)	0.002
Time varying variables						
MUAC change between visits (cm)	1.80	(1.59; 2.04)	<0.001	2.15	(2.07; 2.22)	<0.001
WHZ change between visits	0.44	(0.40; 0.47)	<0.001	0.40	(0.37; 0.44)	<0.001
MUAC group						
MUAC≥12.5 cm/MUAC<12.5 cm	0.89	(0.56; 1.39)	0.603	0.72	(0.38; 1.36)	0.313
Wasted						
No Z-score ≥-2/Yes: Z-score <-2	1.31	(1.07; 1.62)	0.010	1.03	(0.91; 1.16)	0.637

¹Computed using multilevel regression analysis test (stata melogit command);

²OR = Odds Ratio;

³CI = confidence Interval;

⁴HAZ = Length/height-for-age Z-score (2006 WHO reference curves); stunted if HAZ<-2;

⁵V0 = recruitment into the cohort visit;

⁶MUAC = Mid-Upper Arm Circumference ⁷WHZ = weight-for-length/height Z-score (2006 WHO reference curves); wasted = yes if WHZ<-2; ⁸HAZ = Height/length-for-Age Z-score (2006 WHO reference curves); stunted = yes if WHZ<-2.

<https://doi.org/10.1371/journal.pone.0259765.t004>

since the 2014 Cambodia Demographic Health Survey (CDHS) [11, 45], confirming the results of a study that compared the rates found in successive CDHS [44]. This stunting decline rate of only around 1.1% per year only from 2010 to 2016 (39,2% to 32,5%) is insufficient for Cambodia to meet the World Health Assembly target of reducing the number of stunted by 40% by 2025 [1, 46]. Thus a review of current interventions and scaling up of interventions that can have an impact on accelerating the reduction of stunting are urgently required. A collaboration of private and public health sectors in delivering health and nutrition services, adoption of age-appropriate feeding practices at scale, and targeting of the poor with safety net programs have been suggested as effective strategies for reducing stunting in South East Asia, including Cambodia, even if the impact of such interventions for Cambodia is not clear today [18].

The detrimental short and long term consequences of undernutrition, including stunting, on survival and development during infancy and early childhood, are well documented [21, 47]. In our study, the prevalence of stunting was already high to very high at recruitment regardless of the age group cohort for children under the age of 24 months. This suggests that children are stunted at the most vulnerable period of their brain development, a piece of further evidence on the need to address stunting in Cambodia urgently [21, 47]. Some of the long-term health consequences of undernutrition include an increase in the risk of chronic non-communicable diseases (NCDs) such as obesity, diabetes and hypertension. The prevalence of these NCDs is on the rise in South East Asia, including Cambodia [44, 48]. From a human capital development perspective, a reduced potential in educational attainment due to undernutrition and countries' economic growth has been demonstrated as well [49–52].

The high prevalence of stunting at recruitment for children in the age group cohort of 0–5 months suggests that prenatal factors play an essential role in its occurrence. Parallely, the increase in the stunting prevalence over time indicates that postnatal factors also contribute to the maintenance and development of stunting. Analyses conducted using the same data used for this study by other authors have identified several postnatal determinants of stunting in our study population, including inappropriate feeding practices, increased exposure to animal faeces leading to *Giardia Duodenalis* infection, high incidence of infections and inadequate water and sanitation practices [26–28, 53]. Also, an interaction between wasting, infection and stunting in Cambodian under-five children has previously been documented [27].

Another study conducted in Cambodia showed that both feeding practices for mothers during pregnancy and lactation period and children below 14 months were inadequate in urban and rural settings of the country [54]. Interventions to address these determinants exist in Cambodia and are all scalable [20, 22]. Based on our findings, we advocate for their rapid scale-up to accelerate stunting reduction in Cambodia.

We observed a positive correlation between weight replenishment, measured by WHZ or MUAC change, and linear catch-up growth even though only a small proportion of wasted children in our study reported having received the recommended supplementary or therapeutic feeding treatment. It is thus likely that if all the wasted children had received the appropriate treatment, the increase in stunting prevalence over time might have been reduced. A study conducted in Burkina Faso showed that stunted children with wasting respond well to recommended treatment [55]. Increasing investment in the prevention and treatment of wasting in Cambodia will likely contribute to an accelerated reduction of stunting. Prevention and treatment of wasting should be part of the national package of critical interventions to address stunting. This recommendation is also supported by the fact that around 14 to 19% of the stunted children in this study were also wasted.

Our results support previously reported evidence suggesting that linear catch-up growth and weight catch up growth occurs with a lag of approximately three months. This suggests that the absence of effect of the treatment of wasting on linear growth reported in many earlier studies was because the height gain outcome was measured too early in the course of nutrition recovery [7, 17, 24, 56]. Our results, those of Isanaka et al., Lelijveld et al., and that of Schoenbuchner et al., who have examined this question recently, are in favour of introducing a systematic follow up after discharge at three and or 6 six months after the wasting correction criteria have been met and to collect data to measure the effect of wasting correction on catch up linear growth [17, 24, 56]. This will contribute to increasing the body of evidence on the need to link stunting and wasting programming. It will also create the possibility of including a new component in the guidelines such as a follow up multiple micronutrients supplementation to support the linear catch-up growth, the immune system recovery, address persisting iron deficiency and other micronutrient deficiencies [57, 58]. Such follow up is already

advocated for by the report of high relapse rate and mortality during the first three to six months following discharge from wasting correction programmes [59, 60].

Although both MUAC changes and WHZ changes are indicators of tissue deficit replenishment, in our study, we observed a difference in the time lags when both MUAC and WHZ change correlate to the same height change, with MUAC change having the shortest time lag. This finding has not been reported before and is thus difficult to interpret given that both MUAC and WHZ contribute to wasting correction and increase in body mass [61]. A plausible explanation is that this discrepancy is just a reflection of the weak correlation between WHZ change and MUAC change observed in our study, indicating that the two parameters selected different children (S1 Fig). A similar discrepancy between MUAC change and weight change in children recovering from wasting has been reported in the literature [62]. However, other authors have also reported a good correlation between MUAC and weight changes during recovery from severe wasting [62–64]. Nonetheless, this finding has a crucial programmatic implication. It indicates that MUAC change may be an early indicator of response to the stunting reduction interventions, although both MUAC change and weight change should always be used.

Currently, stunting is viewed as a chronic form of undernutrition that, once established, is largely irreversible, especially after two years of age, and that can only be addressed in developmental context by long-term preventive interventions [65, 66]. Our analysis showed that stunted children had accelerated linear growth suggests that many of the stunted children retained the potential to catch up on growth, with some achieving complete recovery. This finding calls into question the general thinking that stunting can be addressed only by preventive interventions. As demonstrated by our findings, interventions to support the potential of recovery from stunting are also critical. Studies that showed that recovered stunted children had similar cognitive capacity to those who never experienced stunting further support the need to add curative approaches to the package of interventions to reduce stunting [47, 67, 68].

Strengths and limitations

The results described in this paper need to be interpreted, taking into account the strengths and weaknesses of the data. The study's main strength is the longitudinal design that allowed follow-up of the same children prospectively and calculation of the estimates of interests for each follow-up visit. A second important strength is that the study was conducted in the three different livelihood contexts of Cambodia and in a "real world" environment; thus, the results can be generalized to the Cambodian context and other similar settings. The third strength is the cohort starting sample of over 5000 children, and a median number of data collection points per child of 5 out of the seven planned provided enough power for calculating key outcomes at all the follow-up visits.

However, this was a longitudinal open cohort study with many of those included in the analysis enrolled either at V0 or at FV5. This may have introduced a seasonal bias that we could not assess in the present study. It was also not a birth cohort; thus, the contribution of prenatal growth factors might be overstated. Also, we could not attach the growth pattern of a period to a particular health event or intervention. This hindered the adjustment of the observed associations between the wasting parameters and the linear growth outcomes. However, this is common in many community-based studies. Also, previous publications from the original analysis of these data addressed some of these aspects [27, 28, 53, 69]. These papers have suggested that infections may be contributing to the occurrence of stunting [27, 53, 69]. However, in our analysis for children below two years at enrolment, type of toilet and source of drinking water were not consistently associated with the linear growth parameters studied

departing from Manzoni et al. conclusion [53]. Finally, we could not adjust for wealth and household food security as these variables were not available. Despite these limitations, we consider that the results of our analyses and the derived conclusions are plausible, given that they are in line with the most recently published observations. The strength of the associations are high, and the adjustment with the variables available did not reduce the strength of the observed associations.

Conclusion

We provide evidence of a positive relationship between wasting parameters and linear growth. Our findings have underscored several points of programming and policy relevance, including 1) wasting was a contributing factor to stunting and prevention of wasting may contribute to reducing stunting in Cambodia; 2) WHZ and MUAC increases were positively correlated with accelerated linear growth suggesting wasting correction can contribute to preventing and reversing stunting; 3) Linear catch-up growth can be observed 3 to 4 months from the time of wasting correction, suggesting a need to include a post-wasting recovery follow up period in the guidelines for the management of wasting. Our findings support the few studies recommending a re-thinking of the current approach of wasting and stunting separation in terms of policy, programming, and financing and a move towards a more integrated approach to maximize the effectiveness of the interventions targeting both wasted and stunted children [17, 24, 65]. This will help accumulate evidence of the effect of SAM wasting treatment on linear catch-up growth, serve as a period for promoting catch up linear growth and corrections of persisting deficiencies.

The evidence in our study strongly supports the integration of both preventative and curative wasting and stunting programming, given the multiplicative effect they have on the prevalence and long-term effects of undernutrition and overnutrition and other metabolic syndrome-related diseases. Such integration has the potential of accelerating the attainment of World Health Assembly targets on undernutrition in a country where both wasting and stunting are of public importance, such as in Cambodia [6]. Such studies should also be repeated in other contexts such as South-East Asia.

Supporting information

S1 Fig. Overlap between weight-for-height Z-score (WHZ) and Mid-Upper Arm Circumference (MUAC) change categories during the different periods. WHZ change and MUAC change at visit 0 to follow up visit 6.
(PDF)

S1 Table. Factors associated with height-for-age Z-score (HAZ) change at the different follow-up visits. HAZ change from visit 0 to follow up visit 6.
(DOCX)

S2 Table. Association between stunting, wasting parameters and occurrence of accelerated linear growth at the different follow up visits. Change in wasting and stunting parameters from visit 0 to follow up visit 6.
(DOCX)

S1 File.
(XLS)

S2 File.
(XLS)

S3 File.
(XLSX)

S4 File.
(CSV)

S5 File.
(XLSX)

S6 File.
(CSV)

S7 File.
(CSV)

Acknowledgments

The authors would like to thank Roland Kupka for his insightful comments that improved the manuscript. We also thank Selamawit Negash and Sophonneary Prak for facilitating access permissions to the data sets. Thanks also to Gabriela Hondru for her support in extracting variables for this study from the MyHealth study database.

Disclaimer: Mueni Mutunga is a UNICEF staff member. The opinions and statements in this article are those of the author and may not reflect official UNICEF policies.

Author Contributions

Conceptualization: Mueni Mutunga, Alexandra Rutishauser-Perera, Sophonneary Prak, Paluku Bahwere.

Data curation: Paluku Bahwere.

Formal analysis: Paluku Bahwere.

Project administration: Alexandra Rutishauser-Perera.

Supervision: Mueni Mutunga.

Validation: Mueni Mutunga, Frank T. Wieringa.

Visualization: Mueni Mutunga, Paluku Bahwere.

Writing – original draft: Mueni Mutunga, Paluku Bahwere.

Writing – review & editing: Mueni Mutunga, Alexandra Rutishauser-Perera, Arnaud Laillou, Jacques Berger, Frank T. Wieringa, Paluku Bahwere.

References

1. Branca F, Grummer-Strawn L, Borghi E, Blössner Md, Onis Md. Extension of the WHO maternal, infant and young child nutrition targets to 2030. *SCN News*. 2015(41):55–8.
2. World Health Organization. Global nutrition targets 2025: Policy brief series. World Health Organization; 2014.
3. Haddad L, Achadi E, Bendech MA, Ahuja A, Bhatia K, Bhutta Z, et al. The Global Nutrition Report 2014: actions and accountability to accelerate the world's progress on nutrition. *J Nutr*. 2015; 145(4):663–71. <https://doi.org/10.3945/jn.114.206078> PMID: 25740908
4. Harding KL, Aguayo VM, Webb P. Factors associated with wasting among children under five years old in South Asia: Implications for action. *PLoS One*. 2018; 13(7):e0198749. <https://doi.org/10.1371/journal.pone.0198749> PMID: 29969457

5. Lozano R, Fullman N, Abate D, Abay SM, Abbafati C, Abbasi N, et al. Measuring progress from 1990 to 2017 and projecting attainment to 2030 of the health-related Sustainable Development Goals for 195 countries and territories: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet*. 2018; 392(10159):2091–138. [https://doi.org/10.1016/S0140-6736\(18\)32281-5](https://doi.org/10.1016/S0140-6736(18)32281-5) PMID: 30496107
6. Mutunga M, Frison S, Rava M, Bahwere P. The Forgotten Agenda of Wasting in Southeast Asia: Burden, Determinants and Overlap with Stunting: A Review of Nationally Representative Cross-Sectional Demographic and Health Surveys in Six Countries. *Nutrients*. 2020; 12(2). <https://doi.org/10.3390/nu12020559> PMID: 32093376
7. Briend A, Khara T, Dolan C. Wasting and stunting—similarities and differences: policy and programmatic implications. *Food Nutr Bull*. 2015; 36(1 Suppl):S15–S23.
8. Islam MR, Rahman MS, Rahman MM, Nomura S, de Silva A, Lanerolle P, et al. Reducing childhood malnutrition in Bangladesh: the importance of addressing socio-economic inequalities. *Public Health Nutr*. 2019;1–11. <https://doi.org/10.1017/S136898001900140X> PMID: 31203835
9. Kang Y, Kim J. Risk factors for undernutrition among children 0–59 months of age in Myanmar. *Matern Child Nutr*. 2019; 15(4):e12821. <https://doi.org/10.1111/mcn.12821> PMID: 30919554
10. Chaparro C, Oot L, Sethuraman K. Overview of the nutrition situation in seven countries in Southeast Asia. 2014.
11. National Institute of Statistics, Directorate-General for Health, ICF International. Cambodia Demographic and Health Survey, 2014. Phnom Penh, Cambodia, and Rockville, Maryland, USA: Institute of Statistics, Directorate-General for Health, and ICF International.; 2015.
12. Myatt M, Khara T, Schoenbuchner S, Pietzsch S, Dolan C, Lelijveld N, et al. Children who are both wasted and stunted are also underweight and have a high risk of death: a descriptive epidemiology of multiple anthropometric deficits using data from 51 countries. *Arch Public Health*. 2018; 76:28. <https://doi.org/10.1186/s13690-018-0277-1> PMID: 30026945
13. Khara T, Mwangome M, Ngari M, Dolan C. Children concurrently wasted and stunted: A meta-analysis of prevalence data of children 6–59 months from 84 countries. *Matern Child Nutr*. 2017. <https://doi.org/10.1111/mcn.12516> PMID: 28944990
14. McDonald CM, Olofin I, Flaxman S, Fawzi WW, Spiegelman D, Caulfield LE, et al. The effect of multiple anthropometric deficits on child mortality: a meta-analysis of individual data in 10 prospective studies from developing countries. *Am J Clin Nutr*. 2013; 97(4):896–901. <https://doi.org/10.3945/ajcn.112.047639> PMID: 23426036
15. Olofin I, McDonald CM, Ezzati M, Flaxman S, Black RE, Fawzi WW, et al. Associations of suboptimal growth with all-cause and cause-specific mortality in children under five years: a pooled analysis of ten prospective studies. *PLoS One*. 2013; 8(5):e64636. <https://doi.org/10.1371/journal.pone.0064636> PMID: 23734210
16. Garenne M, Myatt M, Khara T, Dolan C, Briend A. Concurrent wasting and stunting among under-five children in Niakhar, Senegal. *Matern Child Nutr*. 2019; 15(2):e12736. <https://doi.org/10.1111/mcn.12736> PMID: 30367556
17. Schoenbuchner SM, Dolan C, Mwangome M, Hall A, Richard SA, Wells JC, et al. The relationship between wasting and stunting: a retrospective cohort analysis of longitudinal data in Gambian children from 1976 to 2016. *Am J Clin Nutr*. 2019; 110(2):498–507. <https://doi.org/10.1093/ajcn/nqy326> PMID: 30753251
18. Bloem MW, de Pee S, Hop le T, Khan NC, Lailou A, Minarto, et al. Key strategies to further reduce stunting in Southeast Asia: lessons from the ASEAN countries workshop. *Food Nutr Bull*. 2013; 34(2 Suppl):S8–16. <https://doi.org/10.1177/15648265130342S103> PMID: 24049992
19. Isanaka S, Roederer T, Djibo A, Luquero FJ, Nombela N, Guerin PJ, et al. Reducing wasting in young children with preventive supplementation: a cohort study in Niger. *Pediatrics*. 2010; 126(2):e442–e50. <https://doi.org/10.1542/peds.2009-2814> PMID: 20660552
20. Bhutta ZA, Ahmed T, Black RE, Cousens S, Dewey K, Giugliani E, et al. What works? Interventions for maternal and child undernutrition and survival. *Lancet*. 2008; 371(9610):417–40. [https://doi.org/10.1016/S0140-6736\(07\)61693-6](https://doi.org/10.1016/S0140-6736(07)61693-6) PMID: 18206226
21. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet*. 2013; 382(9890):427–51. [https://doi.org/10.1016/S0140-6736\(13\)60937-X](https://doi.org/10.1016/S0140-6736(13)60937-X) PMID: 23746772
22. Bhutta ZA, Das JK, Rizvi A, Gaffey MF, Walker N, Horton S, et al. Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost? *Lancet*. 2013; 382(9890):452–77. [https://doi.org/10.1016/S0140-6736\(13\)60996-4](https://doi.org/10.1016/S0140-6736(13)60996-4) PMID: 23746776
23. Walker SP, Golden MH. Growth in length of children recovering from severe malnutrition. *Eur J Clin Nutr*. 1988; 42(5):395–404. PMID: 3135181

24. Isanaka S, Hitchings MDT, Berthe F, Briend A, Grais RF. Linear growth faltering and the role of weight attainment: Prospective analysis of young children recovering from severe wasting in Niger. *Matern Child Nutr.* 2019;e12817. <https://doi.org/10.1111/mcn.12817> PMID: 30903806
25. Wieringa FT, Gauthier L, Greffeuille V, Som SV, Dijkhuizen MA, Laillou A, et al. Identification of Acute Malnutrition in Children in Cambodia Requires Both Mid Upper Arm Circumference and Weight-For-Height to Offset Gender Bias of Each Indicator. *Nutrients.* 2018; 10(6). <https://doi.org/10.3390/nu10060786> PMID: 29921769
26. Hondru G, Laillou A, Wieringa FT, Poirot E, Berger J, Christensen DL, et al. Age-Appropriate Feeding Practices in Cambodia and the Possible Influence on the Growth of the Children: A Longitudinal Study. *Nutrients.* 2019; 12(1). <https://doi.org/10.3390/nu12010012> PMID: 31861580
27. Hondru G, Wieringa FT, Poirot E, Berger J, Som SV, Theary C, et al. The Interaction between Morbidity and Nutritional Status among Children under Five Years Old in Cambodia: A Longitudinal Study. *Nutrients.* 2019; 11(7). <https://doi.org/10.3390/nu11071527> PMID: 31284369
28. Caron Y, Hong R, Gauthier L, Laillou A, Wieringa FT, Berger J, et al. Stunting, Beyond Acute Diarrhoea: *Giardia Duodenalis*, in Cambodia. *Nutrients.* 2018; 10(10). <https://doi.org/10.3390/nu10101420> PMID: 30282952
29. WHO G, UNICEF. WHO Child Growth Standards and the Identification of Severe Acute Malnutrition in Infants and Children: A Joint Statement by the World Health Organization and the United Nations Children's Fund: World health organization (WHO); 2009 2009.
30. Monteiro PO, Victora CG. Rapid growth in infancy and childhood and obesity in later life—a systematic review. *Obes Rev.* 2005; 6(2):143–54. <https://doi.org/10.1111/j.1467-789X.2005.00183.x> PMID: 15836465
31. Cameron N. The human growth curve, canalization and catch-up growth. *Human growth and development: Elsevier*; 2012. p. 1–22.
32. Pradeilles R, Norris T, Ferguson E, Gazdar H, Mazhar S, Bux Mallah H, et al. Factors associated with catch-up growth in early infancy in rural Pakistan: A longitudinal analysis of the women's work and nutrition study. *Matern Child Nutr.* 2019; 15(2):e12733. <https://doi.org/10.1111/mcn.12733> PMID: 30345717
33. Marshall A, Altman DG, Royston P, Holder RL. Comparison of techniques for handling missing covariate data within prognostic modelling studies: a simulation study. *BMC Med Res Methodol.* 2010; 10:7. <https://doi.org/10.1186/1471-2288-10-7> PMID: 20085642
34. Ali AM, Dawson SJ, Blows FM, Provenzano E, Ellis IO, Baglietto L, et al. comparison of methods for handling missing data on immunohistochemical markers in survival analysis of breast cancer. *Br J Cancer.* 2011; 104(4):693–9. <https://doi.org/10.1038/sj.bjc.6606078> PMID: 21266980
35. Royston P. PTREND: Stata module for trend analysis for proportions. 2014.
36. Greenland S. Modeling and variable selection in epidemiologic analysis. *Am J Public Health.* 1989; 79(3):340–9. <https://doi.org/10.2105/ajph.79.3.340> PMID: 2916724
37. Hole AR, editor *Mixed logit modeling in Stata—an overview.* United Kingdom Stata Users' Group Meetings 2013; 2013: Stata Users Group.
38. StataCorp L. *Stata multilevel mixed-effects reference manual.* College Station, TX: StataCorp LP. 2013.
39. Maleta K, Virtanen SM, Espo M, Kulmala T, Ashorn P. Seasonality of growth and the relationship between weight and height gain in children under three years of age in rural Malawi. *Acta Paediatr.* 2003; 92(4):491–7. <https://doi.org/10.1111/j.1651-2227.2003.tb00584.x> PMID: 12801119
40. Brown KH, Black RE, Becker S. Seasonal changes in nutritional status and the prevalence of malnutrition in a longitudinal study of young children in rural Bangladesh. *Am J Clin Nutr.* 1982; 36(2):303–13. PMID: 6808822
41. Tomkins AM, Dunn DT, Hayes RJ, Bradley AK. Seasonal variations in the nutritional status of urban Gambian children. *Br J Nutr.* 1986; 56(3):533–43. <https://doi.org/10.1079/bjn19860134> PMID: 3676229
42. Rikimaru T. What Are the Current Situations and the Challenges of Maternal and Child Malnutrition in Asia? *J Nutr Sci Vitaminol (Tokyo).* 2015; 61 Suppl:S63–5. <https://doi.org/10.3177/jnsv.61.S63> PMID: 26598890
43. de Onis M, Borghi E, Arimond M, Webb P, Croft T, Saha K, et al. Prevalence thresholds for wasting, overweight and stunting in children under 5 years. *Public Health Nutr.* 2019; 22(1):175–9. <https://doi.org/10.1017/S1368980018002434> PMID: 30296964
44. Greffeuille V, Sophonneary P, Laillou A, Gauthier L, Hong R, Hong R, et al. Inequalities in Nutrition between Cambodian Women over the Last 15 Years (2000–2014). *Nutrients.* 2016; 8(4):224. <https://doi.org/10.3390/nu8040224> PMID: 27104556
45. National Institute of Statistics DGfHII. *Demographic and Health Survey 2014.* Phnom Penh, Cambodia, and Rockville, Maryland, USA: National Institute of Statistics, Directorate-General for Health, and ICF International; 2015.

46. Greffeuille V, Sophonneary P, Laillou A, Gauthier L, Hong R, Hong R, et al. Persistent Inequalities in Child Undernutrition in Cambodia from 2000 until today. *Nutrients*. 2016; 8(5).
47. Alderman H, Behrman JR, Glewwe P, Fernald L, Walker S. Evidence of Impact of Interventions on Growth and Development during Early and Middle Childhood. In: rd, Bundy DAP, Silva N, Horton S, Jamison DT, Patton GC, editors. *Child and Adolescent Health and Development*. Washington (DC) 2017.
48. Murtagh E NCD. Risk Factor Collaboration. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet*. 2017.
49. Borja JB. The impact of early nutrition on health: key findings from the Cebu Longitudinal Health and Nutrition Survey (CLHNS). *Malays J Nutr*. 2013; 19(1):1–8. PMID: 24800380
50. Berkman DS, Lescano AG, Gilman RH, Lopez SL, Black MM. Effects of stunting, diarrhoeal disease, and parasitic infection during infancy on cognition in late childhood: a follow-up study 11. *Lancet*. 2002; 359(9306):564–71. [https://doi.org/10.1016/S0140-6736\(02\)07744-9](https://doi.org/10.1016/S0140-6736(02)07744-9) PMID: 11867110
51. Moench-Pfanner R, Silo S, Laillou A, Wieringa F, Hong R, Hong R, et al. The Economic Burden of Malnutrition in Pregnant Women and Children under 5 Years of Age in Cambodia. *Nutrients*. 2016; 8(5). <https://doi.org/10.3390/nu8050292> PMID: 27187462
52. Fink G, Rockers PC. Childhood growth, schooling, and cognitive development: further evidence from the Young Lives study. *Am J Clin Nutr*. 2014; 100(1):182–8. <https://doi.org/10.3945/ajcn.113.080960> PMID: 24808488
53. Manzoni G, Laillou A, Samnang C, Hong R, Wieringa FT, Berger J, et al. Child-Sensitive WASH Composite Score and the Nutritional Status in Cambodian Children. *Nutrients*. 2019; 11(9). <https://doi.org/10.3390/nu11092142> PMID: 31500268
54. Som SV, Prak S, Laillou A, Gauthier L, Berger J, Poirot E, et al. Diets and Feeding Practices during the First 1000 Days Window in the Phnom Penh and North-Eastern Districts of Cambodia. *Nutrients*. 2018; 10(4). <https://doi.org/10.3390/nu10040500> PMID: 29670006
55. Fabiansen C, Phelan KP, Cichon B, Ritz C, Briend A, Michaelsen KF, et al. Short children with a low mid-upper arm circumference respond to food supplementation: an observational study from Burkina Faso. *Am J Clin Nutr*. 2016; 103(2):415–21. <https://doi.org/10.3945/ajcn.115.124644> PMID: 26739038
56. Lelijveld N, Seal A, Wells JC, Kirkby J, Opondo C, Chimwezi E, et al. Chronic disease outcomes after severe acute malnutrition in Malawian children (ChroSAM): a cohort study. *The Lancet Global health*. 2016. [https://doi.org/10.1016/S2214-109X\(16\)30133-4](https://doi.org/10.1016/S2214-109X(16)30133-4) PMID: 27470174
57. Bahwere P, James P, Abdissa A, Getu Y, Getnet Y, Sadler K, et al. Use of tuberculin skin test for assessment of immune recovery among previously malnourished children in Ethiopia. *BMC Res Notes*. 2017; 10(1):570. <https://doi.org/10.1186/s13104-017-2909-x> PMID: 29115985
58. Akomo P, Bahwere P, Murakami H, Banda C, Maganga E, Kathumba S, et al. Soya, maize and sorghum ready-to-use therapeutic foods are more effective in correcting anaemia and iron deficiency than the standard ready-to-use therapeutic food: a randomized controlled trial. *BMC Public Health*. 2019; 19(1):806. <https://doi.org/10.1186/s12889-019-7170-x> PMID: 31234806
59. Stobaugh HC, Mayberry A, McGrath M, Bahwere P, Zagre NM, Manary MJ, et al. Relapse after severe acute malnutrition: A systematic literature review and secondary data analysis. *Matern Child Nutr*. 2018:e12702. <https://doi.org/10.1111/mcn.12702> PMID: 30246929
60. Chang CY, Trehan I, Wang RJ, Thakwalakwa C, Maleta K, Deitchler M, et al. Children successfully treated for moderate acute malnutrition remain at risk for malnutrition and death in the subsequent year after recovery. *J Nutr*. 2013; 143(2):215–20. <https://doi.org/10.3945/jn.112.168047> PMID: 23256140
61. Chomtho S, Fewtrell MS, Jaffe A, Williams JE, Wells JC. Evaluation of arm anthropometry for assessing pediatric body composition: evidence from healthy and sick children. *Pediatr Res*. 2006; 59(6):860–5. <https://doi.org/10.1203/01.pdr.0000219395.83159.91> PMID: 16641219
62. Tadesse AW, Tadesse E, Berhane Y, Ekstrom EC. Choosing Anthropometric Indicators to Monitor the Response to Treatment for Severe Acute Malnutrition in Rural Southern Ethiopia-Empirical Evidence. *Nutrients*. 2017; 9(12). <https://doi.org/10.3390/nu9121339> PMID: 29292787
63. Binns P, Dale N, Hoq M, Banda C, Myatt M. Relationship between mid-upper arm circumference and weight changes in children aged 6–59 months. *Arch Public Health*. 2015; 73:54. <https://doi.org/10.1186/s13690-015-0103-y> PMID: 26693279
64. Mogendi JB, De Steur H, Gellynck X, Saeed HA, Makokha A. Efficacy of mid-upper arm circumference in identification, follow-up and discharge of malnourished children during nutrition rehabilitation. *Nutr Res Pract*. 2015; 9(3):268–77. <https://doi.org/10.4162/nrp.2015.9.3.268> PMID: 26060539
65. Khara T, Dolan C. Technical Briefing paper: Association between wasting and stunting, policy, programming and research implications. Oxford: Emergency Nutrition Network (ENN).

66. Golden MH. Is complete catch-up possible for stunted malnourished children? *European journal of clinical nutrition*. 1994; 48 Suppl 1:S58–70; discussion S1. PMID: [8005092](https://pubmed.ncbi.nlm.nih.gov/8005092/)
67. Crookston BT, Penny ME, Alder SC, Dickerson TT, Merrill RM, Stanford JB, et al. Children who recover from early stunting and children who are not stunted demonstrate similar levels of cognition. *The Journal of Nutrition*. 2010; 140(11):1996–2001. <https://doi.org/10.3945/jn.109.118927> PMID: [20844188](https://pubmed.ncbi.nlm.nih.gov/20844188/)
68. Crookston BT, Schott W, Cueto S, Dearden KA, Engle P, Georgiadis A, et al. Postinfancy growth, schooling, and cognitive achievement: Young Lives. *The American journal of clinical nutrition*. 2013; 98(6):1555–63. <https://doi.org/10.3945/ajcn.113.067561> PMID: [24067665](https://pubmed.ncbi.nlm.nih.gov/24067665/)
69. Forrer A, Khieu V, Schar F, Hattendorf J, Marti H, Neumayr A, et al. *Strongyloides stercoralis* is associated with significant morbidity in rural Cambodia, including stunting in children. *PLoS Negl Trop Dis*. 2017; 11(10):e0005685. <https://doi.org/10.1371/journal.pntd.0005685> PMID: [29059195](https://pubmed.ncbi.nlm.nih.gov/29059195/)