



Role of Gamma Glutamyl Transpeptidase to Platelet Ratio (GPR) in Predicting Advanced Liver Fibrosis in Patients with Chronic Hepatitis C Infection

**Raja Taha Yaseen Khan ^{a*}, Hina Ismail ^a, Muhammad Danish ^a,
Muhammad Qaiser Panezai ^a, Abdullah Nasir ^a, Kiran Bajaj ^a,
Ghulamullah Lail ^a, Rajesh Mandhwani ^a, Husnain Ali Metlo ^a, Danish Kumar ^a,
Munir Tareen ^a, Abbas Ali Tasneem ^a, Farina M. Hanif ^a, Syed Mudassir Laeeq ^a
and Nasir Hasan Luck ^a**

^a *Department of Hepatogastroenterology, Sindh Institute of Urology and Transplantation, Pakistan.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMMR/2022/v34i1931455

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/87730>

Original Research Article

Received 05 April 2022

Accepted 08 June 2022

Published 21 June 2022

ABSTRACT

The liver biopsy is the gold standard for the diagnosis of liver fibrosis, because of its invasiveness, high cost and lack of repeatability its use is limited. A new parameter widely used these days for evaluating the grade of hepatic fibrosis is the gamma-glutamyl transpeptidase (GGT)-to-Platelet ratio (GPR) and has shown great benefit in this regard. The aim of our study was to evaluate the role of GPR as a noninvasive predictor of liver fibrosis in patients with chronic hepatitis C in the study population.

Methods: All patients with chronic hepatitis C and compensated liver disease were included in the study after informed consent. Patient's baseline characteristics were recorded. Patient's baseline Complete blood count (CBC) and Liver function tests were also recorded. Patients then underwent shear wave elastography (SWE) to stratify the degree of fibrosis. These indices were used to calculate Gamma glutamyl transpeptidase (GGT) / platelet ratio. Results were presented as means

*Corresponding author: E-mail: raja_taha101488@hotmail.com;

± SD for quantitative data or as numbers with percentages for qualitative data. Continuous variables were analyzed using the Student's t-test; while categorical variables were analyzed using the Chi-square test. A p value of <0.05 was considered statistically significant.

Results: A total of 91 patients were included in the study. Out of 91 patients, 56(61.5%) were males. At baseline, 57(62.6%) patients had ≥F3 fibrosis (advanced fibrosis or cirrhosis). Mean GPR was 1.5±2.1. Area under ROC (Receiver Operating Curve) was obtained for GPR in predicting advanced liver fibrosis (≥F3) was 0.8 (p≤0.001). Higher GPR values were significantly associated with prediction of advanced liver fibrosis (≥F3) in patients with chronic hepatitis C with a sensitivity was of 94.74%, specificity of 62%, positive predictive value of 80.69%, negative predictive value was of 87.50% and diagnostic accuracy of 82.42%.

Conclusion: The GPR found to be significantly associated with liver fibrosis in HCV patients presented in our clinic. However, further studies are needed to validate the role of GPR in predicting liver fibrosis.

Keywords: HCV; shear wave elastography; liver fibrosis; platelets; gamma glutamyl transpeptidase; GPR.

1. INTRODUCTION

Globally, hepatitis C is the most common causes of cirrhosis and hepatocellular carcinoma associated with increased morbidity and mortality¹. In order to halt the disease progression, early diagnosis and prompt treatment are the corner stones of treatment of hepatitis C resulting in decreased number of patients with advanced fibrosis, cirrhosis and hepatocellular carcinoma (HCC) [1-3]. The gold standard method to assess the degree of fibrosis is liver biopsy. The use of liver biopsy is restricted due to its invasiveness, cost, complications and contraindications [3-6]. Currently, clinical practice needs non-invasive cost effective assessment tools that can easily predict liver inflammation and advanced liver fibrosis [7].

Recently, Shear Wave Elastography (SWE) has emerged as a reliable non-invasive tool in staging liver fibrosis which accurately assesses the liver fibrosis and has eliminated the need of liver biopsy. However, there are certain confounding factors that can alter the yield of SWE and have limited its utility including severe inflammation, increased bilirubin, cholestasis, presence of ascites, cost, and lack of skilled operators [8-9]. Lately, certain serum biomarkers and bedside scores have been proposed that are useful in predicting advanced liver fibrosis and are of particular importance as they do not require skilled staff and costly equipment and these scores have the advantage of comprising only two or three laboratory investigations [10].

Among these non-invasive tools, AST to Platelet Ratio Index (APRI) and Fibrosis-4 (FIB-4) are

the 2 widely used non-invasive models [11,12]. Several novel non-invasive tools to assess liver fibrosis include neutrophil-to-lymphocyte ratio (NLR), red cell distribution width (RDW)-to-platelet ratio (RPR) and AST/alanine aminotransferase (ALT) ratio (AAR) [13]. Recently, the gamma-glutamyl transpeptidase (GGT) to platelet ratio (GPR) had emerged as a novel marker in the estimation of liver fibrosis. Studies in patients with HBV infection have revealed the superior accuracy of GPR as compared to the classical biomarkers APRI and FIB-4 in this regard in west African cohorts while French studies found GPR to be non-superior to APRI and FIB-4 [14]. GPR also failed to show superior accuracy in Brazilian and Chinese cohort [15,16].

The aim our study was to evaluate the diagnostic accuracy of GPR in predicting advanced liver fibrosis in patients with chronic HCV infection in our population. There is little work done in this regard as GPR has not been studied in chronic HCV infection for the prediction of advanced liver fibrosis specifically in our population.

2. MATERIALS AND METHODS

This was a cross-sectional prospective study which was carried out in the department of hepato-gastroenterology, Sindh institute of urology and Transplantation from January 2019 to June 2019. All patient with age >18 years and chronic HCV infection were enrolled in the study. The patients with conditions other than chronic HCV infection that can result in enhanced liver injury and inflammation and liver fibrosis such as chronic HBV and HDV co-infection, metabolic conditions such as Non-alcoholic liver disease,

Wilson disease, hemochromatosis and patients with hepatocellular carcinoma were excluded from the study.

All the patients then underwent history taking and physical examination followed by non-invasive laboratory investigations such as complete blood count and liver function tests. These patients then underwent imaging studies including ultrasound abdomen for features of chronic liver disease and Shear Wave Elastography (SWE) to assess fibrosis.

GPR was calculated using the formula:

GPR=

$$\text{Gamma Glutamyl Transpeptidase} \left(\frac{\text{---}}{\text{ULN}} \right) \text{Platelet Count (109 /L)}$$

2.1 Assessment of Fibrosis Staging by SWE: [17]

Two-dimensional SWE was performed using a LOGIQ E9 (GE Healthcare, Milwaukee, WI, USA) and a convex probe (1–6 MHz) by experienced radiologist with expertise in SWE and have already performed 200 SWE. After the visualization of the right liver lobe through an intercostal space, and a region of interest (ROI) was marked at less than 5 cm below the liver capsule and avoided major vessels. Measurements were taken 10–15 times while the patients held their breath for 5–10 seconds. The result was considered reliable when 10 successful shots and an Inter-Quartile Range(IQR)/median ratio should be less than 30%.

2.2 SWE Fibrosis Grading: [17]

Normal-mild fibrosis (F0-F1) - <5.4 kPa

Significant fibrosis (F2) - 5.4-9.9kPa

Advanced fibrosis (F3) - 9.9-12.9kPa

Cirrhosis (F4) - >12.9 kPa

2.3 Statistical Analysis

The data was entered and analyzed using SPSS.v20. The baseline characteristics of all the patient were recorded. Continuous variables were expressed as Mean ±SD, while categorical variables were expressed as frequency and percentages. Categorical variables were analyzed using Chi square test while student t test was used to analyze continuous variables. GPR was calculated and area under the Receiver operating curve (AUROC) was obtained to evaluate the diagnostic performance of GPR in predicting advanced fibrosis. A p-value of ≤0.05 was considered statistically significant.

3. RESULTS

A total of 91 patients were included in the study. Out of 91 patients, 56 (61.5%) were males. Mean age was 43.51±12.5 years. The baseline characteristics of the patients are shown in Tables 1 and 2. Child Pugh Class (CTP) A was noted in 60(65.9%) at the time of presentation while CTP B was noted in 31(64.1%) patients. Most of the patients did not have any comorbidities. At baseline, 52(54%) patients had F4 fibrosis as measured by SWE, while 24(37.4%) patients had F2 fibrosis, 10(11%) patients had F1 fibrosis and 5(5.5%) had F3 fibrosis. 57(62.6%) patients had ≥F3 fibrosis (advanced fibrosis or cirrhosis). Mean platelet count was 85±47(x10⁹/L). Mean Gamma glutamyl Trans-peptidase was 74±59 (IU/L). Mean GPR was 1.5±2.1.

Comparison of continuous and categorical variables in terms of advanced fibrosis (Tables 2 and 3). Area under ROC was obtained for GPR in predicting advanced liver fibrosis (≥F3) was 0.8 (p<0.001) (Figs. 1, 2). At a GPR cut off of ≥0.6, the sensitivity was of 94.74%, specificity of 62%, positive predictive value of 80.69%, negative predictive value was of 87.50% and diagnostic accuracy of 82.42% in predicting advanced liver fibrosis.(≥F3) in patients with chronic hepatitis C Table 4.

Table 1. Baseline characteristics of the population included in the study (n-91)

Study population		n-91(%)
Mean age(years±S.D)		43.5±12.5
Gender	Male	56(61.5)
	Female	35(38.5)
Hemoglobin(g/dL)		10.8±2.1
Total Leucocyte Count(x10 ⁹ /L)		4.5±2.4

Study population		n-91(%)
Platelet Count(x10 ⁹ /L)		80±52
Total Bilirubin(mg/dl)		1.3±0.75
Alkaline Phosphatase(IU/L)		190±187
Aspartate Transaminase(AST)(IU/L)		59±52
Alanine Transaminase(ALT)(IU/L)		45±34
Gamma Glutamyl Transpeptidase(GGT)(IU/L)		74±59
Child Turcotte Pugh Score	A	60(65.9)
	B	31(34.1)
MELD score		10.9±3.4
Shear wave Elastography(SWE)		
F1 Fibrosis		10(11%)
F2 Fibrosis		24(37.4%)
F3 Fibrosis		5(5.5%)
F4 Fibrosis		52(54%)
Esophageal varices	Present	50(55)
	Absent	41(45)
Liver Fibrosis	Present	57(62.6)
	Absent	34(37.4)
GPR(Gamma Glutamyl Transpeptidase to Platelet Ratio)		1.5±2.1

Table 2. Comparison of continuous variables in terms of advanced liver fibrosis

Variable	No Fibrosis(≤F2)	Advanced Fibrosis(≥F3)(n-57)	p-value
	(n-34)	(n-57)	
	Mean ± SD	Mean ± SD	
Age	43.3 ± 10.75	43.65 ± 13.5	0.889
Hemoglobin(g/dL)	10.6 ± 2.5	10.9 ± 1.75	0.334
Total Leucocyte Count(x10 ⁹ /L)	5.3± 2.6	4.1 ± 2.6	0.024
Platelet Count(x10 ⁹ /L)	108.7 ± 66.6	62.9 ± 29.5	≤0.001
Total Bilirubin(mg/dl)	0.95± 0.62	14.5 ± 0.75	≤0.001
Alkaline Phosphatase(IU/L)	119±44	224± 29	0.005
Aspartate Transaminase(AST)(IU/L)	35± 19	73±60	0.001
Alanine Transaminase(ALT)(IU/L)	26±13	57±37	≤0.001
Gamma Glutamyl Transpeptidase(GGT)(IU/L)	48±21	90±69	0.001
CTP score	5.7± 1.2	6.4 ± 1.3	0.015
MELD Score	10.2±4.3	11.3 ± 3.6	0.208
GPR	1.0± 2	1.8± 2	0.04

Table 3. Comparison of categorical variables in terms of advanced liver fibrosis

Variable		Liver fibrosis (≥F3)(n-57)	No Fibrosis(≤F2)(n-34)	p-value
		n(%)	n(%)	
Gender	Male	39(68.4)	17(50)	0.08
	Female	18(31.6)	17(50)	
Esophageal varices	Yes	43(75.4)	23(67.6)	0.421
	No	14(24.6)	11(32.4)	
CTP score	A	32(56.1)	28(82.4)	0.01
	B	28(43.9)	6(17.6)	
GPR	≥0.6	54(94.7)	13(38.2)	≤0.001
	<0.6	3(5.3)	21(61.8)	




Fig. 1. Area under Receiver Operating curve for GPR in the prediction of significant hepatic fibrosis ($\geq F3$) in chronic hepatitis C is 0.8 (p-value ≤ 0.001)




Fig. 2. Box plot showing association of high Gamma Glutamyl Transpeptidase (GPR) with advanced liver fibrosis

Table 4. Diagnostic accuracy of Gamma Glutamyl Transpeptidase(GPR) in predicting advanced (\geq F3) liver fibrosis

Statistics	Percentage(95%CI)
Sensitivity	94.74%(85.3-98.9)
Specificity	62%(43.6-78.8)
Positive Predictive Value	80.60%(72.96-86.48)
Negative Predictive Value	87.50%(69.3-95.6)
Diagnostic Accuracy	82.42%(73-89.6)

4. DISCUSSION

Chronicity of liver disease can result in liver fibrosis and cirrhosis due to continued inflammation and non-avoidance of the etiological factor. In order to prevent disease progression, accurate and timely evaluation of degree of liver fibrosis is required. The gold standard test for this purpose is liver parenchymal biopsy but due to its invasiveness and technical expertise it is less often performed. Currently, non-invasive imaging methods such as fibroscan, SWE and TE are used to document the degree of liver fibrosis. However, high cost, the lack of availability of these techniques everywhere, presence of certain confounding factors along with lack of skilled operators can alter the interpretation of these imaging techniques. Currently, non-invasive tool are utilized for assessment of liver fibrosis in addition to these imaging techniques.

A model of gamma-glutamyl transpeptidase-to platelet ratio (GPR), as bedside available test useful in predicting the levels of liver fibrosis in chronic hepatitis B(CHB) patients in African population was proposed by Lemoine and his colleagues [14]. Ding R et al. [18] further validated this model in Chinese population. In our study, we have used the GPR ratio in predicting liver fibrosis in chronic HCV population using shear wave elastography to evaluate the degree of fibrosis.

Our study population included those patients who had chronic HCV infection and who were either under treatment for it with direct acting antivirals (Sofosbuvir/Daclatasvir) or already treated for it with the same regimen.

We categorized the patients into advanced fibrosis (\geq F3) and non-advanced fibrosis (\leq F2). The application of GPR in the two groups revealed the association of High GPR with advanced fibrosis (p -value \leq 0.001). Similar results for GPR were observed in the previous studies done by Vardar et al. [19] in a Turkish

cohort in 2009, lemoine et al. [14] in an African cohort with AUROC of 0.80 and Ding R et al. [18] in a Chinese cohort with AUROC 0.78 in predicting advanced (\geq F3) fibrosis in CHB patients [19]. Lui et al. [20] used the GPR cutoff values suggested by lemoine et al. [14] in both HBsAg positive and HbsAg negative population in evaluating the degree of fibrosis and found lower sensitivity and specificity of GPR in both the groups. Previously, Nada *and her colleagues* have shown the association of high GPR of \geq 0.31 with advanced fibrosis in chronic hepatitis C patients in an Egyptian population with a sensitivity of 92%, specificity of 88%, PPV of 86% and NPV of 91% [21]. In our study, GPR cut off of \geq 0.6 was obtained using AUROC. The sensitivity of GPR was 91%, which was comparable to that of the previous studies. However, when compared to the previous studies, the specificity, PPV and NPV of GPR was lower in our studied population with a good diagnostic accuracy of 80% in predicting advanced liver fibrosis (\geq F3) in patients with chronic hepatitis C.

There were certain limitations to our study. First of all the small sample size has affected the specificity of the GPR in our population. Secondly, for the assessment of liver fibrosis, SWE was used instead of liver biopsy which is the gold standard due to its invasiveness and high cost.

The strength of the study lies in the fact that this is the pioneer study from this region which has evaluated the role of GPR in HCV patients.

5. CONCLUSION

The diagnostic accuracy of GPR in predicting advanced fibrosis in our population was effective in 82.42% patients. However, studies comprising larger sample size are needed to validate this score before using it as an important non-invasive assessment tool in predicting advanced fibrosis.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT

An informed consent was taken from all the patients before enrollment in the study.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Alter HJ, Seeff LB. Recovery, persistence, and sequelae in hepatitis C virus infection: A perspective on long-term outcome. *Semin. Liver Dis.* 2000;20:17-35.
2. Hoofnagle JH, di Bisceglie AM. The treatment of chronic viral hepatitis. *N. Engl. J. Med.* 1997;336:347-356.
3. Shiha G, et al. Asian-Pacific Association for the Study of the Liver (APASL) consensus guidelines on invasive and non-invasive assessment of hepatic fibrosis: a 2016 update. *Hepatol. Int.* 11, 1–30.
4. Cadranet JF, Rufat P, Degos F. Practice of liver biopsy in France: results of a prospective nationwide survey. For the Group of epidemiology of the French Association for the study of the liver (AEF). *Hepatology.* 2000;32:477–481.
5. Bravo AA, Sheth SG, Chopra S. Liver biopsy. *N Engl J Med.* 2001;344:495–500.
6. Piccinino F, Sagnelli E, Pasquale G, Giusti G. Complications following percutaneous liver biopsy. A multicenter retrospective study on 68,276 biopsies. *J Hepatol.* 1986; 2:165–173.
7. Zhang Z, Wang G, Kang K, Wu G, Wang P. The diagnostic accuracy and clinical utility of three noninvasive models for predicting liver fibrosis in patients with HBV infection. *PLoS One.* 2016;11: e0152757.
8. Kim JH, Kim MN, Han KH, Kim SU. Clinical application of transient elastography in patients with chronic viral hepatitis receiving antiviral treatment. *Liver Int.* 2015;35:1103–1115.
9. Seo YS, et al. Accuracy of transient elastography in assessing liver fibrosis in chronic viral hepatitis: A multicentre, retrospective study. *Liver Int.* 2015;35: 2246–2255
10. World Health Organization. Guidelines for the Prevention, Care and Treatment of Persons with Chronic Hepatitis B Infection; 2015.
11. Wai CT, Greenson JK, Fontana RJ, Kalbfleisch JD, Marrero JA, Conjeevaram HS et al. A simple noninvasive index can predict both significant fibrosis and cirrhosis in patients with chronic hepatitis C. *Hepatology.* 2003;38(20):518-526.
12. Xiao G, Yang J, Yan L. Comparison of diagnostic accuracy of aspartate aminotransferase to platelet ratio index and fibrosis-4 index for detecting liver fibrosis in adult patients with chronic hepatitis B virus infection: a systemic review and meta-analysis. *Hepatology.* 2015;61:292-302
13. Chen B, Ye B, Zhang J, Ying L, Chen Y. RDW to platelet ratio: a novel noninvasive index for predicting hepatic fibrosis and cirrhosis in chronic hepatitis B. *PLoS One.* 8, e68780.
14. Lemoine M, et al. The gamma-glutamyl transpeptidase to platelet ratio (GPR) predicts significant liver fibrosis and cirrhosis in patients with chronic HBV infection in West Africa. *Gut.* 2016;65: 1369–1376. DOI: 10.1136/gutjnl-2015-309260
15. Li Q, et al. The Gamma-Glutamyl-Transpeptidase to Platelet Ratio Does Not Show Advantages than APRI and Fib-4 in Diagnosing Significant Fibrosis and Cirrhosis in Patients with Chronic Hepatitis B: A Retrospective Cohort Study in China. *Medicine.* 2016;95:e3372. DOI:10.1097/MD.0000000000003372
16. Schiavon LL, Narciso-Schiavon JL, Ferraz MLG, Silva AEB, Carvalho-Filho RJ. The γ -glutamyl transpeptidase to platelet ratio

- (GPR) in HBV patients: just adding up? Gut; 2016.
DOI:10.1136/gutjnl-2016-312658
17. Song P, et al. Performance of 2-dimensional ultrasound shear wave elastography in liver fibrosis detection using magnetic resonance elastography as the reference standard: A Pilot study. J Ultrasoun; 2016.
 18. Ding R, Lu W, Zhou X, Huang D, Wang Y, Li X, Yan L, Lin W, Song S, Zhang Z and Chen L. A Novel Non-invasive Model Based on GPR for the Prediction of Liver Fibrosis in Patients with Chronic Hepatitis B. Front. Med. 2021;8:727706.
 19. Vardar R, Vardar E, Demiri S, Sayhan SE, Bayol U, Yildiz C, et al. Is there any noninvasive marker replace the needle liver biopsy predictive for liver fibrosis, in patients with chronic hepatitis? Hepatogastroenterology. 2009;56:1459–65.
 20. Liu DP, Lu W, Zhang ZQ, Wang YB, Ding RR, Zhou XL, Huang D, Li XF. Comparative evaluation of GPR versus APRI and FIB-4 in predicting different levels of liver fibrosis of chronic hepatitis B. Journal of Viral Hepatitis. 2018;25(5): 581-9.
 21. Nada A, Sabry A, Elabd NS, Elbahr O. The Gamma-glutamyl Transpeptidase to Platelet Ratio (GPR) Predict Significant Liver Fibrosis and Cirrhosis in Chronic Egyptian HCV Patients. Afro-Egyptian Journal of Infectious and Endemic Diseases. 2020;10(2):226-32.

© 2022 Khan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/87730>*